

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
27 January 2005 (27.01.2005)

PCT

(10) International Publication Number
WO 2005/007647 A1

(51) International Patent Classification⁷: **C07D 401/12**,
401/14, 403/12, 405/14, 409/14, 413/14, 417/14, 239/52,
A61K 31/506, 31/5377, 31/505, A61P 3/10

LEHMANN, Juerg [CH/US]; 9710 Mesa Springs Way
#7, San Diego, CA 92126 (US). BRUCE, Marc, A.
[US/US]; 662 Corte Raquel, San Marcos, CA 92069 (US).

(21) International Application Number:
PCT/US2004/022327

(74) Agent: **GODDARD, Christine, A.**; Cozen O'Connor,
1900 Market Street, Philadelphia, PA 19103 (US).

(22) International Filing Date: 9 July 2004 (09.07.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/486,728 11 July 2003 (11.07.2003) US
60/487,370 14 July 2003 (14.07.2003) US

(71) Applicant (for all designated States except US): **ARENA
PHARMACEUTICALS, INC.** [US/US]; 6166 Nancy
Ridge Drive, San Diego, CA 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **JONES, Robert,
M.** [GB/US]; 10937 Corte luz del Sol, San Diego, CA
92130 (US). **SEMPLE, Graeme** [GB/US]; 15920 Camino
Codorniz, San Diego, CA 92127 (US). **XIONG, Yifeng**
[US/US]; 4711 Caminito Eva, San Diego, CA 92130
(US). **SHIN, Young-Jun** [KR/US]; 11752 Westview
Parkway, #3, San Diego, CA 92126 (US). **REN, Albert, S.**
[CN/US]; 4266 Corte de Sausilito, San Diego, CA 92130
(US). **CALDERON, Imelda** [MX/US]; 10971 Caminito
Arcada, San Diego, CA 92131 (US). **CHOI, Jin, Sun,
Karoline** [US/US]; 11120 Caminito Alares, San Diego,
CA 92126 (US). **FIORAVANTI, Beatriz** [BR/US]; 8540
Costa Verde Blvd., #4344, San Diego, CA 92122 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.


(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: TRISUBSTITUTED ARYL AND HETEROARYL DERIVATIVES AS MODULATORS OF METABOLISM AND THE
PROPHYLAXIS AND TREATMENT OF DISORDERS RELATED THERETO



(57) Abstract: The present invention relates to certain trisubstituted aryl and heteroaryl derivatives of Formula (I) that are modulators of metabolism. Accordingly, compounds of the present invention are useful in the prophylaxis or treatment of metabolic disorders and complications thereof, such as, diabetes and obesity.

**TRISUBSTITUTED ARYL AND HETEROARYL DERIVATIVES AS MODULATORS OF
METABOLISM AND THE PROPHYLAXIS AND TREATMENT OF DISORDERS
RELATED THERETO**

5

FIELD OF THE INVENTION

The present invention relates to certain trisubstituted aryl and heteroaryl derivatives that are modulators of glucose metabolism. Accordingly, compounds of the present invention are useful in the prophylaxis or treatment of metabolic disorders and complications thereof, such as, diabetes and obesity.

BACKGROUND OF THE INVENTION

Diabetes mellitus is a serious disease afflicting over 100 million people worldwide. In the United States, there are more than 12 million diabetics, with 600,000 new cases diagnosed each year.

Diabetes mellitus is a diagnostic term for a group of disorders characterized by abnormal glucose homeostasis resulting in elevated blood sugar. There are many types of diabetes, but the two most common are Type I (also referred to as insulin-dependent diabetes mellitus or IDDM) and Type II (also referred to as non-insulin-dependent diabetes mellitus or NIDDM).

The etiology of the different types of diabetes is not the same; however, everyone with diabetes has two things in common: overproduction of glucose by the liver and little or no ability to move glucose out of the blood into the cells where it becomes the body's primary fuel.

People who do not have diabetes rely on insulin, a hormone made in the pancreas, to move glucose from the blood into the cells of the body. However, people who have diabetes either don't produce insulin or can't efficiently use the insulin they produce; therefore, they can't move glucose into their cells. Glucose accumulates in the blood creating a condition called hyperglycemia, and over time, can cause serious health problems.

Diabetes is a syndrome with interrelated metabolic, vascular, and neuropathic components. The metabolic syndrome, generally characterized by hyperglycemia, comprises alterations in carbohydrate, fat and protein metabolism caused by absent or markedly reduced insulin secretion and/or ineffective insulin action. The vascular syndrome consists of abnormalities in the blood vessels leading to cardiovascular, retinal and renal complications. Abnormalities in the peripheral and autonomic nervous systems are also part of the diabetic syndrome.

People with IDDM, which accounts for about 5% to 10% of those who have diabetes, don't produce insulin and therefore must inject insulin to keep their blood glucose levels normal. IDDM is characterized by low or undetectable levels of endogenous insulin production caused by destruction of the insulin-producing β cells of the pancreas, the characteristic that most readily distinguishes IDDM from NIDDM. IDDM, once termed juvenile-onset diabetes, strikes young and older adults alike.

Approximately 90 to 95% of people with diabetes have Type II (or NIDDM). NIDDM subjects produce insulin, but the cells in their bodies are insulin resistant: the cells don't respond properly to the hormone, so glucose accumulates in their blood. NIDDM is characterized by a relative disparity between endogenous insulin production and insulin requirements, leading to elevated blood glucose levels. In contrast to IDDM, there is always some endogenous insulin production in NIDDM; many NIDDM patients have normal or even elevated blood insulin levels, while other NIDDM patients have inadequate insulin production (Rotwein, R. et al. *N. Engl. J. Med.* 308, 65-71 (1983)). Most people diagnosed with NIDDM are age 30 or older, and half of all new cases are age 55 and older. Compared with whites and Asians, NIDDM is more common among Native Americans, African-Americans, Latinos, and Hispanics. In addition, the onset can be insidious or even clinically inapparent, making diagnosis difficult.

The primary pathogenic lesion on NIDDM has remained elusive. Many have suggested that primary insulin resistance of the peripheral tissues is the initial event. Genetic epidemiological studies have supported this view. Similarly, insulin secretion abnormalities have been argued as the primary defect in NIDDM. It is likely that both phenomena are important contributors to the disease process (Rimoin, D. L., et al. Emery and Rimoin's Principles and Practice of Medical Genetics 3rd Ed. 1:1401-1402 (1996)).

Many people with NIDDM have sedentary lifestyles and are obese; they weigh approximately 20% more than the recommended weight for their height and build. Furthermore, obesity is characterized by hyperinsulinemia and insulin resistance, a feature shared with NIDDM, hypertension and atherosclerosis.

Obesity and diabetes are among the most common human health problems in industrialized societies. In industrialized countries a third of the population is at least 20% overweight. In the United States, the percentage of obese people has increased from 25% at the end of the 1970s, to 33% at the beginning the 1990s. Obesity is one of the most important risk factors for NIDDM. Definitions of obesity differ, but in general, a subject weighing at least 20% more than the recommended weight for his/her height and build is considered obese. The risk of developing NIDDM is tripled in subjects 30% overweight, and three-quarters with NIDDM are overweight.

Obesity, which is the result of an imbalance between caloric intake and energy expenditure, is highly correlated with insulin resistance and diabetes in experimental animals and human. However, the molecular mechanisms that are involved in obesity-diabetes syndromes are not clear. During early development of obesity, increase insulin secretion balances insulin resistance and protects patients from hyperglycemia (Le Stunff, et al. *Diabetes* 43, 696-702 (1989)). However, after several decades, β cell function deteriorates and non-insulin-dependent diabetes develops in about 20% of the obese population (Pederson, P. *Diab. Metab. Rev.* 5, 505-509 (1989)) and (Brancati, F. L., et al., *Arch. Intern. Med.* 159, 957-963 (1999)). Given its high prevalence in modern societies, obesity has thus become the leading risk factor for NIDDM (Hill, J. O., et al., *Science* 280, 1371-1374 (1998)). However, the factors which

predispose a fraction of patients to alteration of insulin secretion in response to fat accumulation remain unknown.

Whether someone is classified as overweight or obese is generally determined on the basis of their body mass index (BMI) which is calculated by dividing body weight (kg) by height squared (m²). Thus, the units of BMI are kg/m² and it is possible to calculate the BMI range associated with minimum mortality in each decade of life. Overweight is defined as a BMI in the range 25-30 kg/m², and obesity as a BMI greater than 30 kg/m² (see TABLE below). There are problems with this definition in that it does not take into account the proportion of body mass that is muscle in relation to fat (adipose tissue). To account for this, obesity can also be defined on the basis of body fat content: greater than 25% and 30% in males and females, respectively.

**CLASSIFICATION OF WEIGHT BY
BODY MASS INDEX (BMI)**

BMI	CLASSIFICATION
< 18.5	Underweight
18.5-24.9	Normal
25.0-29.9	Overweight
30.0-34.9	Obesity (Class I)
35.0-39.9	Obesity (Class II)
>40	Extreme Obesity (Class III)

As the BMI increases there is an increased risk of death from a variety of causes that is independent of other risk factors. The most common diseases with obesity are cardiovascular disease (particularly hypertension), diabetes (obesity aggravates the development of diabetes), gall bladder disease (particularly cancer) and diseases of reproduction. Research has shown that even a modest reduction in body weight can correspond to a significant reduction in the risk of developing coronary heart disease.

Compounds marketed as anti-obesity agents include Orlistat (XENICALTM) and Sibutramine. Orlistat (a lipase inhibitor) inhibits fat absorption directly and tends to produce a high incidence of unpleasant (though relatively harmless) side-effects such as diarrhea. Sibutramine (a mixed 5-HT/noradrenaline reuptake inhibitor) can increase blood pressure and heart rate in some patients. The serotonin releaser/reuptake inhibitors fenfluramine (PondiminTM) and dexfenfluramine (ReduxTM) have been reported to decrease food intake and body weight over a prolonged period (greater than 6 months). However, both products were withdrawn after reports of preliminary evidence of heart valve abnormalities associated with their use. Accordingly, there is a need for the development of a safer anti-obesity agent.


Obesity considerably increases the risk of developing cardiovascular diseases as well. Coronary insufficiency, atheromatous disease, and cardiac insufficiency are at the forefront of the cardiovascular complication induced by obesity. It is estimated that if the entire population had an ideal weight, the risk of coronary insufficiency would decrease by 25% and the risk of cardiac insufficiency and of cerebral vascular accidents by 35%. The incidence of coronary diseases is doubled in subjects less than 50 years of age who are 30% overweight. The diabetes patient faces a 30% reduced lifespan. After age 45, people with diabetes are about three times more likely than people without diabetes to have significant heart disease and up to five times more likely to have a stroke. These findings emphasize the inter-relations between risks factors for NIDDM and coronary heart disease and the potential value of an integrated approach to the prevention of these conditions based on the prevention of these conditions based on the prevention of obesity (Perry, I. J., et al., *BMJ* 310, 560-564 (1995)).

Diabetes has also been implicated in the development of kidney disease, eye diseases and nervous-system problems. Kidney disease, also called nephropathy, occurs when the kidney's "filter mechanism" is damaged and protein leaks into urine in excessive amounts and eventually the kidney fails. Diabetes is also a leading cause of damage to the retina at the back of the eye and increases risk of cataracts and glaucoma. Finally, diabetes is associated with nerve damage, especially in the legs and feet, which interferes with the ability to sense pain and contributes to serious infections. Taken together, diabetes complications are one of the nation's leading causes of death.

SUMMARY OF THE INVENTION

The present invention is drawn to compounds which bind to and modulate the activity of a GPCR, referred to herein as **RUP3**, and uses thereof. The term **RUP3** as used herein includes the human sequences found in GeneBank accession numbers XM_066873 and AY288416, and naturally-occurring allelic variants, mammalian orthologs, and recombinant mutants thereof. A preferred human **RUP3** for use in screening and testing of the compounds of the invention is provided in the nucleotide sequence of Seq. ID.No:1 and the corresponding amino acid sequence in Seq. ID.No:2.

One aspect of the present invention encompasses trisubstituted aryl and heteroaryl derivatives as shown in Formula (I):



or a pharmaceutically acceptable salt, hydrate or solvate, or N-oxide thereof;
wherein:

A and B are each independently C₁₋₃ alkylene optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₃ alkyl, C₁₋₄ alkoxy, carboxy, cyano, C₁₋₃ haloalkyl and halogen;

D is O, S, S(O), S(O)₂, CR₂R₃ or N-R₂;

5 E is N, C or CR₄;

-- is a single bond when E is N or CR₄, or a double bond when E is C;

V₁ is selected from the group consisting of C₁₋₃ alkylene, ethynylene and C₁₋₂ heteroalkylene optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₃ alkyl, C₁₋₄ alkoxy, carboxy, cyano, C₁₋₃ haloalkyl and halogen; or V₁ is a bond;

10 V₂ is C₃₋₆ cycloalkylene or C₁₋₃ alkylene wherein each are optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₃ alkyl, C₁₋₄ alkoxy, carboxy, cyano, C₁₋₃ haloalkyl and halogen; or V₂ is a bond;

W is NR₅, O, S, S(O) or S(O)₂; or W is absent;

Q is NR₆, O, S, S(O) or S(O)₂;


15 X is N or CR₇;

Y is N or CR₈;

Z is selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, C₁₋₂ alkylamino, C₂₋₄ dialkylamino, carbamimidoyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₄₋₈ diacylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₂₋₆ dialkylsulfonylamino, formyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylcarboxamide, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, halogen, aryl, heterocyclic, heteroaryl, hydroxyl, hydroxycarbamimidoyl, hydroxylamino, nitro and tetrazolyl, wherein C₁₋₈ alkyl, C₃₋₇ cycloalkyl, and heterocyclic are each optionally substituted with 1, 2, 3 or 4 groups selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, amino, C₁₋₂ alkylamino, C₂₋₄ dialkylamino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, formyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro, and wherein said C₁₋₇ alkyl is optionally substituted with amino; or

20
25
30

Z is a group of Formula (A):



wherein:


R₉ is H, C₁₋₈ alkyl or C₃₋₇ cycloalkyl; and

R₁₀ is H, nitro or nitrile;

Ar₁ is aryl or heteroaryl each optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅; wherein

R₁₁ is selected from the group consisting of C₁₋₅ acyl, C₁₋₆ acylsulfonamide, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfanyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, arylsulfonyl, carbamimidoyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, guanidinyl, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfanyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, heterocyclic, heterocyclic-oxy, heterocyclicsulfonyl, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, hydroxyl, nitro, C₄₋₇ oxo-cycloalkyl, phenoxy, phenyl, sulfonamide, sulfonic acid, and thiol, and wherein C₁₋₅ acyl, C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, arylsulfonyl, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, heteroaryl, phenoxy and phenyl are optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfanyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfanyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, heteroaryl, heterocyclic, hydroxyl, nitro, phenyl, and phosphonooxy, wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; or

R_{11} is a group of Formula (B):



(B)

wherein:

“p” and “r” are each independently 0, 1, 2 or 3; and R₁₆ is H, C₁₋₅ acyl, C₂₋₆ alkenyl, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, halogen, heteroaryl or phenyl, and wherein the heteroaryl or phenyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₄ alkoxy, amino, C₁₋₄ alkylamino, C₂₋₆ alkynyl, C₂₋₈ dialkylamino, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl and hydroxyl; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected form the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, carbo-C₁₋₆-

alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, hydroxyl and nitro; or


- two adjacent groups selected from the group consisting of R₁₂, R₁₃, R₁₄ and R₁₅ together with the atoms to which they are attached form a 5-, 6- or 7-membered cycloalkyl, cycloalkenyl or heterocyclic group fused with Ar₁, wherein the 5-, 6- or 7-membered group is optionally substituted with halogen;

- R₁, R₇ and R₈ are each independently selected from the group consisting of H, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, amino, C₁₋₄ alkylamino, C₂₋₈ dialkylamino, carboxamide, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylsulfonamide, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio and hydroxyl;

- R₂ is selected from the group consisting of C₁₋₈ alkyl, amino, aryl, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl and hydroxyl; and wherein C₁₋₈ alkyl, aryl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₃₋₆-cycloalkyl-C₁₋₃-heteroalkylene, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino and nitro; or

- R₂ is -Ar₂-Ar₃ wherein Ar₂ and Ar₃ are each independently aryl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of H, C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, C₁₋₄ alkylamino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, hydroxyl and nitro; or


- R₂ is a group of Formula (C):



wherein:

R₁₇ is H, C₁₋₈ alkyl, C₃₋₇ cycloalkyl, aryl, heteroaryl or OR₁₉; and R₁₈ is F, Cl, Br, CN or NR₂₀R₂₁; where R₁₉ is H, C₁₋₈ alkyl or C₃₋₇ cycloalkyl, and R₂₀ and R₂₁ are each independently H, C₁₋₈ alkyl, C₃₋₇ cycloalkyl, aryl or heteroaryl; or

R₂ is a group of Formula (D):



(D)

wherein:

G is:

- i) -C(O)-, -C(O)NR₂₃-, -C(O)O-, -OC(O)NR₂₃-, -NR₂₃C(O)O-, -OC(O)-, -C(S)-, -C(S)NR₂₃-, -C(S)O-,
 10 -OC(S)-, -CR₂₃R₂₄-, -O-, -S-, -S(O)- or -S(O)₂- when D is CR₂R₃, or
 ii) -CR₂₃R₂₄C(O)-, -C(O)-, -CR₂₃R₂₄C(O)NR₂₅-, -C(O)NR₂₃-, -C(O)O-, -C(S)-, -C(S)NR₂₃-, -C(S)O-, -CR₂₃R₂₄-, -S(O)₂-, or a bond when D is NR₂,
 wherein R₂₃, R₂₄ and R₂₅ are each independently H or C₁₋₈ alkyl; and R₂₂ is H, C₁₋₈ alkyl, C₂₋₆ alkynyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5
 15 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl,
 20 C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heteroaryl, heterocyclic, hydroxyl, hydroxylamino, nitro, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, heteroaryl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄
 25 alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino, and nitro;

30 R₃ is H, C₁₋₈ alkyl, C₁₋₄ alkoxy or hydroxyl; and

R₄, R₅ and R₆ are each independently H, C₁₋₈ alkyl or C₃₋₇ cycloalkyl, wherein said C₁₋₈ alkyl is optionally substituted with C₁₋₄ alkoxy, C₃₋₇ cycloalkyl, or heteroaryl.

One aspect of the present invention pertains to pharmaceutical compositions comprising at least one compound of the present invention and a pharmaceutically acceptable carrier.

One aspect of the present invention pertains to methods for the treatment of a metabolic-related disorder in an individual comprising administering to the individual in need of such treatment a therapeutically effective amount of a compound of the present invention or a pharmaceutical composition thereof.

5 One aspect of the present invention pertains to methods of decreasing food intake of an individual comprising administering to the individual in need thereof a therapeutically effective amount of a compound of the present invention or pharmaceutical composition thereof.

10 One aspect of the present invention pertains to methods of inducing satiety in an individual comprising administering to the individual in need thereof a therapeutically effective amount of a compound of the present invention or pharmaceutical composition thereof.

One aspect of the present invention pertains to methods of controlling or decreasing weight gain of an individual comprising administering to the individual in need thereof a therapeutically effective amount of a compound of the present invention or pharmaceutical composition thereof.

15 One aspect of the present invention pertains to methods of modulating a **RUP3** receptor in an individual comprising contacting the receptor with a compound of the present invention. In some embodiments, the compound is an agonist for the **RUP3** receptor. In some embodiments, the modulation of the **RUP3** receptor is the treatment of a metabolic-related disorder.

20 Some embodiments of the present invention include a method of modulating a **RUP3** receptor in an individual comprising contacting the receptor with a compound of the present invention wherein the modulation of the **RUP3** receptor reduces food intake of the individual.

Some embodiments of the present invention include a method of modulating a **RUP3** receptor in an individual comprising contacting the receptor with a compound of the present invention wherein the modulation of the **RUP3** receptor induces satiety in the individual.

25 Some embodiments of the present invention include a method of modulating a **RUP3** receptor in an individual comprising contacting the receptor with a compound of the present invention wherein the modulation of the **RUP3** receptor controls or reduces weight gain of the individual.

One aspect of the present invention pertains to use of a compound of the present invention for production of a medicament for use in the treatment of a metabolic-related disorder.

30 One aspect of the present invention pertains to use of a compound of the present invention for production of a medicament for use in decreasing food intake in an individual.

One aspect of the present invention pertains to use of a compound of the present invention for production of a medicament for use of inducing satiety in an individual.

One aspect of the present invention pertains to use of a compound of the present invention for production of a medicament for use in controlling or decreasing weight gain in an individual.

35 One aspect of the present invention pertains to a compound of the present invention for use in a method of treatment of the human or animal body by therapy.

One aspect of the present invention pertains to a compound of the present invention for use in a method of treatment of a metabolic-related disorder of the human or animal body by therapy.

One aspect of the present invention pertains to a compound of the present invention for use in a method of decreasing food intake of the human or animal body by therapy.

5 One aspect of the present invention pertains to a compound of the present invention for use in a method of inducing satiety of the human or animal body by therapy.

One aspect of the present invention pertains to a compound of the present invention for use in a method of controlling or decreasing weight gain of the human or animal body by therapy.

10 Some embodiments of the present invention pertain to methods wherein the human has a body mass index of about 18.5 to about 45. In some embodiments, the human has a body mass index of about 25 to about 45. In some embodiments, the human has a body mass index of about 30 to about 45. In some embodiments, the human has a body mass index of about 35 to about 45.

In some embodiments the individual is a mammal. In some embodiments the mammal is a human.

15 In some embodiments, the metabolic-related disorder is hyperlipidemia, type 1 diabetes, type 2 diabetes mellitus, idiopathic type 1 diabetes (Type 1b), latent autoimmune diabetes in adults (LADA), early-onset type 2 diabetes (EOD), youth-onset atypical diabetes (YOAD), maturity onset diabetes of the young (MODY), malnutrition-related diabetes, gestational diabetes, coronary heart disease, ischemic stroke, restenosis after angioplasty, peripheral vascular disease, intermittent claudication, myocardial infarction (e.g. necrosis and apoptosis), dyslipidemia, post-prandial lipemia, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose, metabolic acidosis, ketosis, arthritis, obesity, osteoporosis, hypertension, congestive heart failure, left ventricular hypertrophy, peripheral arterial disease, diabetic retinopathy, macular degeneration, cataract, diabetic nephropathy, glomerulosclerosis, chronic renal failure, diabetic neuropathy, 20 metabolic syndrome, syndrome X, premenstrual syndrome, coronary heart disease, angina pectoris, thrombosis, atherosclerosis, myocardial infarction, transient ischemic attacks, stroke, vascular restenosis, hyperglycemia, hyperinsulinemia, hyperlipidemia, hypertriglyceridemia, insulin resistance, impaired glucose metabolism, conditions of impaired glucose tolerance, conditions of impaired fasting plasma glucose, obesity, erectile dysfunction, skin and connective tissue disorders, foot 25 ulcerations and ulcerative colitis, endothelial dysfunction and impaired vascular compliance.

30 In some embodiments, the metabolic-related disorder is type I diabetes, type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia or syndrome X. In some embodiments, the metabolic-related disorder is type II diabetes. In some embodiments, the metabolic-related disorder is hyperglycemia. In some embodiments, the metabolic-related disorder is hyperlipidemia. In some 35 embodiments, the metabolic-related disorder is hypertriglyceridemia. In some embodiments, the

metabolic-related disorder is type I diabetes. In some embodiments, the metabolic-related disorder is dyslipidemia. In some embodiments, the metabolic-related disorder is syndrome X.

One aspect of the present invention pertains to a method of producing a pharmaceutical composition comprising admixing at least one compound, as described herein, and a pharmaceutically acceptable carrier.

This application is related to two US Provisional Patent Applications, Serial Nos. 60/486,728 filed July 11, 2003; and 60/487,370 filed July 14, 2003, both which are incorporated by reference in their entirety.

Applicant reserves the right to exclude any one or more of the compounds from any of the embodiments of the invention. Applicant additionally reserves the right to exclude any disease, condition or disorder from any of the embodiments of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A shows RT-PCR analysis of RUP3 expression in human tissues. A total of twenty-two (22) human tissues were analyzed.

Figure 1B shows the cDNA Dot-Blot analysis of RUP 3 expression in human tissues.

Figure 1C shows analysis of RUP3 by RT-PCR with isolated human pancreatic islets of Langerhans.

Figure 1D shows analysis of RUP3 expression with cDNAs of rat origin by RT-PCR.

Figure 2A shows a polyclonal anti-RUP3 antibody prepared in Rabbits.

Figure 2B shows the expression of RUP3 in insulin-producing β cells of pancreatic islets.

Figure 3 shows *in vitro* functional activities of RUP3.

Figure 4 shows a RUP3 RNA blot.

DETAILED DESCRIPTION OF THE INVENTION

DEFINITIONS

The scientific literature that has evolved around receptors has adopted a number of terms to refer to ligands having various effects on receptors. For clarity and consistency, the following definitions will be used throughout this patent document.

AGONISTS shall mean moieties that interact and activate the receptor, such as the RUP3 receptor and initiates a physiological or pharmacological response characteristic of that receptor. For example, when moieties activate the intracellular response upon binding to the receptor, or enhance GTP binding to membranes.

AMINO ACID ABBREVIATIONS used herein are set out in TABLE 1:

TABLE 1

ALANINE	ALA	A
ARGININE	ARG	R
ASPARAGINE	ASN	N
ASPARTIC ACID	ASP	D
CYSTEINE	CYS	C
GLUTAMIC ACID	GLU	E
GLUTAMINE	GLN	Q
GLYCINE	GLY	G
HISTIDINE	HIS	H
ISOLEUCINE	ILE	I
LEUCINE	LEU	L
LYSINE	LYS	K
METHIONINE	MET	M
PHENYLALANINE	PHE	F
PROLINE	PRO	P
SERINE	SER	S
THREONINE	THR	T
TRYPTOPHAN	TRP	W
TYROSINE	TYR	Y
VALINE	VAL	V
ALANINE	ALA	A

The term **ANTAGONISTS** is intended to mean moieties that competitively bind to the receptor at the same site as agonists (for example, the endogenous ligand), but which do not activate the intracellular response initiated by the active form of the receptor, and can thereby inhibit the intracellular responses by agonists or partial agonists. Antagonists do not diminish the baseline intracellular response in the absence of an agonist or partial agonist.


CHEMICAL GROUP, MOIETY OR RADICAL:

The term "**C₁₋₅ acyl**" denotes a C₁₋₅ alkyl radical attached to a carbonyl wherein the definition of alkyl has the same definition as described herein; some examples include but not

limited to, acetyl, propionyl, n-butanoyl, *iso*-butanoyl, sec-butanoyl, t-butanoyl (i.e., pivaloyl), pentanoyl and the like.

The term "**C₁₋₅ acyloxy**" denotes an acyl radical attached to an oxygen atom wherein acyl has the same definition as described herein; some examples include but not limited to acetyloxy, propionyloxy, butanoyloxy, *iso*-butanoyloxy, sec-butanoyloxy, t-butanoyloxy and the like.

The term "**C₁₋₆ acylsulfonamide**" refers to a C₁₋₆ acyl attached directly to the nitrogen of the sulfonamide, wherein the definitions for C₁₋₆ acyl and sulfonamide have the same meaning as described herein, and a C₁₋₆ acylsulfonamide can be represented by the following formula:





Some embodiments of the present invention are when acylsulfonamide is a C₁₋₅ acylsulfonamide, some embodiments are C₁₋₄ acylsulfonamide, some embodiments are C₁₋₃ acylsulfonamide, and some embodiments are C₁₋₂ acylsulfonamide. Examples of an acylsulfonamide include, but not limited to, acetylsulfamoyl [-S(=O)₂NHC(=O)Me], propionylsulfamoyl [-S(=O)₂NHC(=O)Et], isobutyrylsulfamoyl, butyrylsulfamoyl, 2-methylbutyrylsulfamoyl, 3-methylbutyrylsulfamoyl, 2,2-dimethylpropionylsulfamoyl, pentanoylsulfamoyl, 2-methylpentanoylsulfamoyl, 3-methylpentanoylsulfamoyl, 4-methylpentanoylsulfamoyl, and the like.

The term "**C₂₋₆ alkenyl**" denotes a radical containing 2 to 6 carbons wherein at least one carbon-carbon double bond is present, some embodiments are 2 to 4 carbons, some embodiments are 2 to 3 carbons, and some embodiments have 2 carbons. Both *E* and *Z* isomers are embraced by the term "**alkenyl**." Furthermore, the term "**alkenyl**" includes di- and tri-alkenyls. Accordingly, if more than one double bond is present then the bonds may be all *E* or *Z* or a mixture of *E* and *Z*. Examples of an alkenyl include vinyl, allyl, 2-butenyl, 3-butenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 2,4-hexadienyl and the like.

The term "**C₁₋₄ alkoxy**" as used herein denotes a radical alkyl, as defined herein, attached directly to an oxygen atom. Examples include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, t-butoxy, isobutoxy, sec-butoxy and the like.

The term "**alkyl**" denotes a straight or branched carbon radical containing 1 to 8 carbons, some embodiments are 1 to 7 carbons, some embodiments are 1 to 6 carbons, some embodiments are 1 to 3 carbons, and some embodiments are 1 or 2 carbons. Examples of an alkyl include, but not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, isobutyl, t-butyl, pentyl, iso-pentyl, t-pentyl, neo-pentyl, 1-methylbutyl [i.e., -CH(CH₃)CH₂CH₂CH₃], 2-methylbutyl [i.e., -CH₂CH(CH₃)CH₂CH₃], n-hexyl and the like.

The term C_{1-6} or C_{1-4} "alkylcarboxamido" or "alkylcarboxamide" denotes a single C_{1-6} or C_{1-4} alkyl group attached to the nitrogen or carbon of an amide group, wherein alkyl has the same definition as found herein. The alkylcarboxamide may be represented by the following:





Examples include, but not limited to, *N*-methylcarboxamide, *N*-ethylcarboxamide, *N*-n-propylcarboxamide, *N*-iso-propylcarboxamide, *N*-n-butylcarboxamide, *N*-sec-butylcarboxamide, *N*-iso-butylcarboxamide, *N*-t-butylcarboxamide and the like.

The term " C_{1-3} alkylene" refers to a C_{1-3} divalent straight carbon group. In some embodiments C_{1-3} alkylene refers to, for example, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, and the like. In some embodiments, C_{1-3} alkylene refers to $-\text{CH}-$, $-\text{CHCH}_2-$, $-\text{CHCH}_2\text{CH}_2-$, and the like wherein these examples relate generally to "A".

The term " C_{1-4} alkylsulfinyl" denotes a C_{1-4} alkyl radical attached to a sulfoxide radical of the formula: $-\text{S}(\text{O})-$ wherein the alkyl radical has the same definition as described herein. Examples include, but not limited to, methylsulfinyl, ethylsulfinyl, n-propylsulfinyl, iso-propylsulfinyl, n-butylsulfinyl, sec-butylsulfinyl, iso-butylsulfinyl, t-butyl, and the like.

The term " C_{1-4} alkylsulfonamide" refers to the groups





wherein C_{1-4} alkyl has the same definition as described herein.

The term " C_{1-4} alkylsulfonyl" denotes a C_{1-4} alkyl radical attached to a sulfone radical of the formula: $-\text{S}(\text{O})_2-$ wherein the alkyl radical has the same definition as described herein. Examples include, but not limited to, methylsulfonyl, ethylsulfonyl, n-propylsulfonyl, iso-propylsulfonyl, n-butylsulfonyl, sec-butylsulfonyl, iso-butylsulfonyl, t-butyl, and the like.

The term " C_{1-4} alkylthio" denotes a C_{1-4} alkyl radical attached to a sulfide of the formula: $-\text{S}-$ wherein the alkyl radical has the same definition as described herein. Examples include, but not limited to, methylsulfanyl (i.e., $\text{CH}_3\text{S}-$), ethylsulfanyl, n-propylsulfanyl, iso-propylsulfanyl, n-butylsulfanyl, sec-butylsulfanyl, iso-butylsulfanyl, t-butyl, and the like.

The term " C_{1-4} alkylthiocarboxamide" denotes a thioamide of the following formulae:



wherein C_{1-4} alkyl has the same definition as described herein.

The term " C_{1-4} alkylthioureyll" denotes the group of the formula:

-NC(S)N- wherein one are both of the nitrogens are substituted with the same or different C₁₋₄ alkyl groups and alkyl has the same definition as described herein. Examples of an alkylthioureyll include, but not limited to, CH₃NHC(S)NH-, NH₂C(S)NCH₃-, (CH₃)₂N(S)NH-, (CH₃)₂N(S)NCH₃-, CH₃CH₂NHC(S)NH-, CH₃CH₂NHC(S)NCH₃-, and the like.

The term "C₁₋₄ alkylureyl" denotes the group of the formula: -NC(O)N- wherein one are both of the nitrogens are substituted with the same or different C₁₋₄ alkyl group wherein alkyl has the same definition as described herein. Examples of an alkylureyl include, but not limited to, CH₃NHC(O)NH-, NH₂C(O)NCH₃-, (CH₃)₂N(O)NH-, (CH₃)₂N(O)NCH₃-, CH₃CH₂NHC(O)NH-, CH₃CH₂NHC(O)NCH₃-, and the like.

The term "C₂₋₆ alkynyl" denotes a radical containing 2 to 6 carbons and at least one carbon-carbon triple bond, some embodiments are 2 to 4 carbons, some embodiments are 2 to 3 carbons, and some embodiments have 2 carbons. Examples of an alkynyl include, but not limited to, ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-pentylnyl, 2-pentylnyl, 3-pentylnyl, 4-pentylnyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl and the like. The term "alkynyl" includes di- and tri-ynes.

The term "amino" denotes the group -NH₂.

The term "C₁₋₄ alkylamino" denotes one alkyl radical attached to an amino radical wherein the alkyl radical has the same meaning as described herein. Some examples include, but not limited to, methylamino, ethylamino, n-propylamino, iso-propylamino, n-butylamino, sec-butylamino, iso-butylamino, t-butylamino, and the like. Some embodiments are "C₁₋₂ alkylamino."

The term "aryl" denotes an aromatic ring radical containing 6 to 10 ring carbons. Examples include phenyl and naphthyl.


The term "arylalkyl" defines a C₁-C₄ alkylene, such as -CH₂-, -CH₂CH₂- and the like, which is further substituted with an aryl group. Examples of an "arylalkyl" include benzyl, phenethylene and the like.

The term "arylcarboxamido" denotes a single aryl group attached to the nitrogen of an amide group, wherein aryl has the same definition as found herein. The example is N-phenylcarboxamide.

The term "arylureyl" denotes the group -NC(O)N- where one of the nitrogens are substituted with an aryl.

The term "benzyl" denotes the group -CH₂C₆H₅.

The term "carbamimidoyl" refers to a group of the following chemical formula:



and in some embodiments, one or both hydrogens are replaced with another group. For example, one hydrogen can be replaced with a hydroxyl group to give a *N*-hydroxycarbamimidoyl group, or one hydrogen can be replaced with an alkyl group to give *N*-methylcarbamimidoyl, *N*-ethylcarbamimidoyl, *N*-propylcarbamimidoyl, *N*-butylcarbamimidoyl, and the like.

The term “**carbo-C₁₋₆-alkoxy**” refers to a C₁₋₆ alkyl ester of a carboxylic acid, wherein the alkyl group is as defined herein. Examples include, but not limited to, carbomethoxy, carboethoxy, carbopropoxy, carboisopropoxy, carbobutoxy, carbo-sec-butoxy, carbo-iso-butoxy, carbo-*t*-butoxy, carbo-*n*-pentoxy, carbo-iso-pentoxy, carbo-*t*-pentoxy, carbo-*neo*-pentoxy, carbo-*n*-hexyloxy, and the like.

The term “**carboxamide**” refers to the group –CONH₂.


The term “**carboxy**” or “**carboxyl**” denotes the group –CO₂H; also referred to as a carboxylic acid group.

The term “**cyano**” denotes the group –CN.

The term “**C₃₋₇ cycloalkenyl**” denotes a non-aromatic ring radical containing 3 to 6 ring carbons and at least one double bond; some embodiments contain 3 to 5 carbons; some embodiments contain 3 to 4 carbons. Examples include cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, and the like.


The term “**C₃₋₇ cycloalkyl**” denotes a saturated ring radical containing 3 to 6 carbons; some embodiments contain 3 to 5 carbons; some embodiments contain 3 to 4 carbons. Examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like.

The term “**C₃₋₆ cycloalkylene**” refers to a divalent cycloalkyl radical, where cycloalkyl is as defined herein, containing 3 to 6 carbons; some embodiments contain 3 to 5 carbons; some embodiments contain 3 to 4 carbons. In some embodiments, the two bonding groups are on the same carbon, for example:



In some embodiments, the two bonding groups are on different carbons.

The term “**C₄₋₈ diacylamino**” denotes an amino group bonded with two acyl groups defined herein wherein the acyl groups may be the same or different, such as:



Examples of C₄₋₈ diacylamino groups include, but limited to, diacetylamino, dipropionylamino, acetylpropionylamino and the like.

The term "**C₂₋₆ dialkylamino**" denotes an amino substituted with two of the same or different alkyl radicals wherein alkyl radical has the same definition as described herein. Some examples include, but not limited to, dimethylamino, methylethylamino, diethylamino, methylpropylamino, methylisopropylamino, ethylpropylamino, ethylisopropylamino, dipropylamino, propylisopropylamino and the like. Some embodiments are "**C₂₋₄ dialkylamino**."

The term "**C₁₋₄ dialkylcarboxamido**" or "**C₁₋₄ dialkylcarboxamide**" denotes two alkyl radicals, that are the same or different, attached to an amide group, wherein alkyl has the same definition as described herein. A C₁₋₄ dialkylcarboxamido may be represented by the following groups:




wherein C₁₋₄ has the same definition as described herein. Examples of a dialkylcarboxamide include, but not limited to, *N,N*-dimethylcarboxamide, *N*-methyl-*N*-ethylcarboxamide, *N,N*-diethylcarboxamide, *N*-methyl-*N*-isopropylcarboxamide, and the like.

The term "**C₂₋₆ dialkylsulfonamido**" refers to one of the following groups shown below:



wherein C₁₋₃ has the same definition as described herein, for example but not limited to, methyl, ethyl, *n*-propyl, isopropyl, and the like.

The term "**C₂₋₆ dialkylthiocarboxamido**" or "**C₂₋₆ dialkylthiocarboxamide**" denotes two alkyl radicals, that are the same or different, attached to a thioamide group, wherein alkyl has the same definition as described herein. A C₁₋₄ dialkylthiocarboxamido may be represented by the following groups:



Examples of a dialkylthiocarboxamide include, but not limited to, *N,N*-dimethylthiocarboxamide, *N*-methyl-*N*-ethylthiocarboxamide and the like.


The term "**C₂₋₆ dialkylsulfonylamino**" refers to an amino group bonded with two C₁₋₃ alkylsulfonyl groups as defined herein.

The term "**ethynylene**" refers to the carbon-carbon triple bond group as represented below:



The term “formyl” refers to the group ---CHO .

The term “guanidine” refers to a group of the following chemical formula:



5 The term “**C₁₋₄ haloalkoxy**” denotes a haloalkyl, as defined herein, which is directly attached to an oxygen atom. Examples include, but not limited to, difluoromethoxy, trifluoromethoxy, 2,2,2-trifluoroethoxy, pentafluoroethoxy and the like.

 The term “**C₁₋₄ haloalkyl**” denotes an C₁₋₄ alkyl group, defined herein, wherein the alkyl is substituted with one halogen up to fully substituted and a fully substituted C₁₋₄ haloalkyl can be represented by the formula C_nL_{2n+1} wherein L is a halogen and “n” is 1, 2, 3 or 4; when more than one halogen is present then they may be the same or different and selected from the group consisting of F, Cl, Br and I, preferably F. Examples of C₁₋₄ haloalkyl groups include, but not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, chlorodifluoromethyl, 2,2,2-trifluoroethyl, pentafluoroethyl and the like.

15 The term “**C₁₋₄ haloalkylcarboxamide**” denotes an alkylcarboxamide group, defined herein, wherein the alkyl is substituted with one halogen up to fully substituted represented by the formula C_nL_{2n+1} wherein L is a halogen and “n” is 1, 2, 3 or 4. When more than one halogen is present they may be the same or different and selected from the group consisting of F, Cl, Br and I, preferably F.


20 The term “**C₁₋₄ haloalkylsulfinyl**” denotes a haloalkyl radical attached to a sulfoxide group of the formula: ---S(O)--- wherein the haloalkyl radical has the same definition as described herein. Examples include, but not limited to, trifluoromethylsulfinyl, 2,2,2-trifluoroethylsulfinyl, 2,2-difluoroethylsulfinyl and the like.

 The term “**C₁₋₄ haloalkylsulfonyl**” denotes a haloalkyl radical attached to a sulfone group of the formula: $\text{---S(O)}_2\text{---}$ wherein haloalkyl has the same definition as described herein. Examples include, but not limited to, trifluoromethylsulfonyl, 2,2,2-trifluoroethylsulfonyl, 2,2-difluoroethylsulfonyl and the like.

 The term “**C₁₋₄ haloalkylthio**” denotes a haloalkyl radical directly attached to a sulfur wherein the haloalkyl has the same meaning as described herein. Examples include, but not limited to, trifluoromethylthio (i.e., CF₃S-), 1,1-difluoroethylthio, 2,2,2-trifluoroethylthio and the like.

 The term “**halogen**” or “**halo**” denotes to a fluoro, chloro, bromo or iodo group.

 The term “**C₁₋₂ heteroalkylene**” refers to a C₁₋₂ alkylene bonded to a heteroatom selected from O, S, S(O), S(O)₂ and NH. Some represented examples include, but not limited to, the groups of the following formulae:




and the like.

The term “**heteroaryl**” denotes an aromatic ring system that may be a single ring, two fused rings or three fused rings wherein at least one ring carbon is replaced with a heteroatom selected from, but not limited to, the group consisting of O, S and N wherein the N can be optionally substituted with H, C₁₋₄ acyl or C₁₋₄ alkyl. Examples of heteroaryl groups include, but not limited to, pyridyl, benzofuranyl, pyrazinyl, pyridazinyl, pyrimidinyl, triazinyl, quinoline, benzoxazole, benzothiazole, 1*H*-benzimidazole, isoquinoline, quinazoline, quinoxaline and the like. In some embodiments, the heteroaryl atom is O, S, NH, examples include, but not limited to, pyrrole, indole, and the like. Other examples include, but not limited to, those in TABLE 2A, TABLE 4, and the like.

The term “**heterocyclic**” denotes a non-aromatic carbon ring (i.e., cycloalkyl or cycloalkenyl as defined herein) wherein one, two or three ring carbons are replaced by a heteroatom selected from, but not limited to, the group consisting of O, S, N, wherein the N can be optionally substituted with H, C₁₋₄ acyl or C₁₋₄ alkyl, and ring carbon atoms optionally substituted with oxo or a thiooxo thus forming a carbonyl or thiocarbonyl group. The heterocyclic group is a 3-, 4-, 5-, 6- or 7-membered containing ring. Examples of a heterocyclic group include but not limited to aziridin-1-yl, aziridin-2-yl, azetidin-1-yl, azetidin-2-yl, azetidin-3-yl, piperidin-1-yl, piperidin-4-yl, morpholin-4-yl, piperzin-1-yl, piperzin-4-yl, pyrrolidin-1-yl, pyrrolidin-3-yl, [1,3]-dioxolan-2-yl and the like. Additional examples of heterocyclic groups are shown in TABLES 2B, 2C, 2D, 2E, 2F and 2G, *infra*.


The term “**heterocyclic-carbonyl**” denotes a heterocyclic group, as defined herein, directly bonded to the carbon of a carbonyl group (i.e., C=O). In some embodiments, a ring nitrogen of the heterocyclic group is bonded to the carbonyl group forming an amide.

Examples include, but not limited to,




and the like.


In some embodiments, a ring carbon is bonded to the carbonyl group forming a ketone group. Examples include, but not limited to,



The term “**heterocyclic-oxy**” refers to a heterocyclic group, as defined herein, that is directly bonded to an oxygen atom. Examples include the following:



The term “**heterocyclicsulfonyl**” denotes a heterocyclic group, as defined herein, with a ring nitrogen where the ring nitrogen is bonded directly to an SO_2 group forming a sulfonamide. Examples include, but not limited to,




The term “**hydroxyl**” refers to the group $-\text{OH}$.

The term “**hydroxylamino**” refers to the group $-\text{NHOH}$.

The term “**nitro**” refers to the group $-\text{NO}_2$.

The term “ **C_{4-7} oxo-cycloalkyl**” refers to a C_{4-7} cycloalkyl, as defined herein, wherein one of the ring carbons is replaced with a carbonyl. Examples of C_{4-7} oxo-cycloalkyl include, but are not limited to, 2-oxo-cyclobutyl, 3-oxo-cyclobutyl, 3-oxo-cyclopentyl, 4-oxo-cyclohexyl, and the like and represented by the following structures respectively:




The term “**perfluoroalkyl**” denotes the group of the formula $-\text{C}_n\text{F}_{2n+1}$; stated differently, a perfluoroalkyl is an alkyl as defined herein wherein the alkyl is fully substituted with fluorine atoms and is therefore considered a subset of haloalkyl. Examples of perfluoroalkyls include CF_3 , CF_2CF_3 , $\text{CF}_2\text{CF}_2\text{CF}_3$, $\text{CF}(\text{CF}_3)_2$, $\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_3$, $\text{CF}_2\text{CF}(\text{CF}_3)_2$, $\text{CF}(\text{CF}_3)\text{CF}_2\text{CF}_3$ and the like.

The term “**phenoxy**” refers to the group $\text{C}_6\text{H}_5\text{O}-$.

The term “phenyl” refers to the group C_6H_5 .


The term “phosphonoxy” refers to a group with the following chemical structure:



The term “sulfonamide” refers to the group $-SO_2NH_2$.

The term “sulfonic acid” refers to the group $-SO_3H$.

The term “tetrazolyl” refers to the five membered heteroaryl of the following formulae:



In some embodiments, the tetrazolyl group is further substituted at either the 1 or 5 position respectively with a group selected from the group consisting of C_{1-3} alkyl, C_{1-3} haloalkyl and C_{1-3} alkoxy.

The term “thiol” denotes the group $-SH$.

CODON shall mean a grouping of three nucleotides (or equivalents to nucleotides) which generally comprise a nucleoside (adenosine (A), guanosine (G), cytidine (C), uridine (U) and thymidine (T)) coupled to a phosphate group and which, when translated, encodes an amino acid.

COMPOSITION shall mean a material comprising at least two compounds or two components; for example, and without limitation, a Pharmaceutical Composition is a Composition comprising a compound of the present invention and a pharmaceutically acceptable carrier.

CONTACT or **CONTACTING** shall mean bringing the indicated moieties together, whether in an in vitro system or an in vivo system. Thus, “contacting” a **RUP3** receptor with a compound of the invention includes the administration of a compound of the present invention to an individual, preferably a human, having a **RUP3** receptor, as well as, for example, introducing a compound of the invention into a sample containing a cellular or more purified preparation containing a **RUP3** receptor.

IN NEED OF PROPHYLAXIS OR TREATMENT as used herein refers to a judgment made by a caregiver (e.g. physician, nurse, nurse practitioner, etc. in the case of humans; veterinarian in the case of animals, including non-human mammals) that an individual or animal requires or will benefit from prophylaxis or treatment. This judgment is made based on a variety of factors that are in the realm of a caregiver’s expertise, but that includes the knowledge that the individual or animal is ill, or will be ill, as the result of a disease, condition or disorder that is treatable by the compounds of the invention. In general, “in need of prophylaxis” refers to the judgment made by the caregiver that the individual will become ill. In this context, the compounds of the invention are used in a protective or preventive manner. However, “in need of treatment” refers to the judgment of the caregiver that

the individual is already ill, therefore, the compounds of the present invention are used to alleviate, inhibit or ameliorate the disease, condition or disorder.

INDIVIDUAL as used herein refers to any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans.

INHIBIT or **INHIBITING**, in relationship to the term "response" shall mean that a response is decreased or prevented in the presence of a compound as opposed to in the absence of the compound.

INVERSE AGONISTS shall mean moieties that bind the endogenous form of the receptor or to the constitutively activated form of the receptor, and which inhibit the baseline intracellular response initiated by the active form of the receptor below the normal base level of activity which is observed in the absence of agonists or partial agonists, or decrease GTP binding to membranes. Preferably, the baseline intracellular response is inhibited in the presence of the inverse agonist by at least 30%, more preferably by at least 50%, and most preferably by at least 75%, as compared with the baseline response in the absence of the inverse agonist.

LIGAND shall mean an endogenous, naturally occurring molecule specific for an endogenous, naturally occurring receptor.

As used herein, the terms **MODULATE** or **MODULATING** shall mean to refer to an increase or decrease in the amount, quality, response or effect of a particular activity, function or molecule.

PHARMACEUTICAL COMPOSITION shall mean a composition comprising at least one active ingredient, whereby the composition is amenable to investigation for a specified, efficacious outcome in a mammal (for example, without limitation, a human). Those of ordinary skill in the art will understand and appreciate the techniques appropriate for determining whether an active ingredient has a desired efficacious outcome based upon the needs of the artisan.

THERAPEUTICALLY EFFECTIVE AMOUNT as used herein refers to the amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal, individual or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes one or more of the following:

(1) Preventing the disease; for example, preventing a disease, condition or disorder in an individual that may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease,

(2) Inhibiting the disease; for example, inhibiting a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., arresting further development of the pathology and/or symptomatology),


and

Cmpd#	Structure	Chemical Name
A65		4-[5-Cyano-6-(2-fluoro-4-propoxy-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A66		4-[5-Cyano-6-(6-propyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A67		4-{5-Cyano-6-[4-(2-dimethylamino-ethylsulfanyl)-2-fluoro-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A68		4-{5-Cyano-6-[4-(2-dimethylamino-ethanesulfonyl)-2-fluoro-phenylamino]-3-oxypyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A69		4-{5-Cyano-6-[2-fluoro-4-(4-methyl-piperazin-1-yl)-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A70		4-{5-Cyano-6-[2-fluoro-4-(3-methyl-butylamino)-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A71		4-[5-Cyano-6-(2-fluoro-4-morpholin-4-yl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A72		4-{5-Cyano-6-[4-(2-dimethylamino-ethylamino)-2-fluoro-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

(3) Ameliorating the disease; for example, ameliorating a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., reversing the pathology and/or symptomatology).

5 Compounds of the Present Invention:

One aspect of the present invention encompasses trisubstituted aryl and heteroaryl derivatives as shown in Formula (I):



or a pharmaceutically acceptable salt, or N-oxide thereof; wherein Ar₁, V₁, V₂, W, Q, X, Y, Z, A, B, D, E, ---, and R₁ have the same definitions as described herein, *supra* and *infra*.

Some embodiments of the present invention encompass trisubstituted aryl and heteroaryl derivatives as shown in Formula (I) wherein:

A and B are each independently C₁₋₃ alkylene optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₃ alkyl, C₁₋₄ alkoxy, carboxy, cyano, C₁₋₃ haloalkyl and halogen;

D is O, S, S(O), S(O)₂, CR₂R₃ or N-R₂;

E is N, C or CR₄;

--- is a single bond when E is N or CR₄, or a double bond when E is C;

V₁ is selected from the group consisting of C₁₋₃ alkylene, ethynylene and C₁₋₂ heteroalkylene optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₃ alkyl, C₁₋₄ alkoxy, carboxy, cyano, C₁₋₃ haloalkyl and halogen; or V₁ is a bond;

V₂ is selected from the group consisting of C₁₋₃ alkylene optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₃ alkyl, C₁₋₄ alkoxy, carboxy, cyano, C₁₋₃ haloalkyl and halogen; or V₂ is a bond;

W is NR₅, O, S, S(O) or S(O)₂; or W is absent;

Q is NR₆, O, S, S(O) or S(O)₂;


X is N or CR₇;

Y is N or CR₈;

Z is selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, C₁₋₂ alkylamino, C₂₋₄ dialkylamino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₄₋₈ diacylamino, C₂₋₆

dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₂₋₆ dialkylsulfonylamino, formyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylcarboxamide, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, halogen, aryl, heterocyclic, heteroaryl, hydroxyl, hydroxylamino, nitro and tetrazolyl, wherein C₁₋₈ alkyl is optionally substituted with 1, 2, 3 or 4 groups selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, amino, C₁₋₂ alkylamino, C₂₋₄ dialkylamino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, formyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro; or

Z is a group of Formula (A):



wherein:


R₉ is H, C₁₋₈ alkyl or C₃₋₇ cycloalkyl; and

R₁₀ is H, nitro or nitrile;

Ar₁ is aryl or heteroaryl each optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅;

where R₁₁ is selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, amino, arylsulfonyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, heterocyclic, heterocyclicsulfonyl, heteroaryl, hydroxyl, nitro, C₄₋₇ oxo-cycloalkyl, phenoxy, phenyl, sulfonamide and sulfonic acid, and wherein C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylsulfonamide, alkylsulfonyl, arylsulfonyl, heteroaryl, phenoxy or phenyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, heteroaryl, heterocyclic, hydroxyl, nitro and phenyl; or

R₁₁ is a group of Formula (B):



wherein:

“p” and “r” are each independently 0, 1, 2 or 3; and

5 R₁₆ is H, C₁₋₅ acyl, C₂₋₆ alkenyl, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, halogen, heteroaryl or phenyl, and wherein the heteroaryl or phenyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₄ alkoxy, amino, C₁₋₄ alkylamino, C₂₋₆ alkynyl, C₂₋₈ dialkylamino, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl and hydroxyl;

10 R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, hydroxyl and nitro; or two adjacent groups selected from the group consisting of R₁₂, R₁₃, R₁₄ and R₁₅ form a 5, 6 or 7 membered cycloalkyl, cycloalkenyl or heterocyclic group with Ar₁ wherein the 5, 6 or 7 membered group is optionally substituted with halogen;


15 R₁, R₇ and R₈ are each independently selected from the group consisting of H, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, amino, C₁₋₄ alkylamino, C₂₋₈ dialkylamino, carboxamide, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylsulfonamide, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio and hydroxyl;

20 R₂ is selected from the group consisting of H, C₁₋₈ alkyl, amino, aryl, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl and hydroxyl; and wherein C₁₋₈ alkyl, aryl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₃₋₆-cycloalkyl-C₁₋₃-heteroalkylene, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino and nitro; or

25 R₂ is -Ar₂-Ar₃ wherein Ar₂ and Ar₃ are each independently aryl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of H, C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋

alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, C₁₋₄ alkylamino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, hydroxyl and nitro; or

5 R₂ is a group of Formula (C):




wherein:

R₁₇ is H, C₁₋₈ alkyl, C₃₋₇ cycloalkyl, aryl, heteroaryl or OR₁₉; and R₁₈ is F, Cl, Br, CN or NR₂₀R₂₁; where R₁₉ is H, C₁₋₈ alkyl or C₃₋₇ cycloalkyl, and R₂₀ and R₂₁ are each

10 independently H, C₁₋₈ alkyl, C₃₋₇ cycloalkyl, aryl or heteroaryl; or

R₂ is a group of Formula (D):



wherein:

G is:

15 i) C(O), C(O)NR₂₃, C(O)O, OC(O), C(S), C(S)NR₂₃, C(S)O, OC(S), CR₂₃R₂₄, O, S, S(O) or S(O)₂ when D is CR₂R₃, or

ii) C(O), C(O)NR₂₃, C(O)O, C(S), C(S)NR₂₃, C(S)O, CR₂₃R₂₄ or S(O)₂ when D is NR₂, where R₂₃ and R₂₄ are each independently H or C₁₋₈ alkyl; and

20 R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthioureyl, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthioureyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro;

R₃ is H, C₁₋₈ alkyl, C₁₋₄ alkoxy or hydroxyl; and

30 R₄, R₅ and R₆ are each independently H, or C₁₋₈ alkyl; or a pharmaceutically acceptable salt, hydrate or solvate thereof.

It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable subcombination.

5 As used herein, "substituted" indicates that at least one hydrogen atom of the chemical group is replaced by a non-hydrogen substituent or group, the non-hydrogen substituent or group can be monovalent or divalent. When the substituent or group is divalent, then it is understood that this group is further substituted with another substituent or group. When a chemical group herein is "substituted" it may have up to the full valance of substitution; for example, a methyl group can be
10 substituted by 1, 2, or 3 substituents, a methylene group can be substituted by 1 or 2 substituents, a phenyl group can be substituted by 1, 2, 3, 4, or 5 substituents, a naphthyl group can be substituted by 1, 2, 3, 4, 5, 6, or 7 substituents and the like. Likewise, "substituted with one or more substituents" refers to the substitution of a group with one substituent up to the total number of substituents physically allowed by the group. Further, when a group is substituted with more than one group they
15 can be identical or they can be different.

It is understood and appreciated that compounds of Formula (I) may have one or more chiral centers, and therefore can exist as enantiomers and/or diastereomers. The invention is understood to extend to and embrace all such enantiomers, diastereomers and mixtures thereof, including but not limited, to racemates. Accordingly, some embodiments of the present invention pertain to compounds
20 of Formula (I) and formulae used throughout this disclosure that are *R* enantiomers. Further, some embodiments of the present invention pertain to compounds of Formula (I) and formulae used throughout this disclosure that are *S* enantiomers. In examples where more than one chiral center is present, then, some embodiments of the present invention include compounds that are *RS* or *SR* enantiomers. In further embodiments, compounds of the present invention are *RR* or *SS* enantiomers.
25 It is understood that compounds of Formula (I) and formulae used throughout this disclosure are intended to represent all individual enantiomers and mixtures thereof, unless stated or shown otherwise.

Many geometric isomers of olefins, C=N double bonds, disubstituted cycloalkyl (i.e., 1,4-cyclohexyl groups), and the like can also be present in the compounds described herein,
30 and all such stable isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms.

Compounds of the invention can also include tautomeric forms, such as keto-enol tautomers, and the like. Tautomeric forms can be in equilibrium or sterically locked into one form by
35 appropriate substitution. It is understood that the various tautomeric forms are within the scope of the compounds of the present invention.


Compounds of the invention can also include all isotopes of atoms occurring in the intermediates and/or final compounds. Isotopes include those atoms having the same atomic number but different mass numbers. For example, isotopes of hydrogen include deuterium and tritium.

In some embodiments, X and Y are each independently N or CH, provided that if either X or Y is CH then the other is N.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein W is NR₅. In some embodiments, R₅ is H.

In some embodiments, W is NH.


In some embodiments compounds of the present invention can be represented by Formula (Ia) as illustrated below:



wherein each variable in Formula (Ia) has the same meaning as described herein, *supra* and *infra*. In some embodiments, V₁ is a bond. In still further embodiments, both V₁ and V₂ are bonds.

In some embodiments, W is O.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein W is O and can be represented by Formula (Ic) as illustrated below:



wherein each variable in Formula (Ic) has the same meaning as described herein, *supra* and *infra*.

Some embodiments of the present invention pertain to compounds of Formula (Ic) wherein V₁ is absent. In some embodiments, Q is an oxygen atom. In still further embodiments, Q is an oxygen atom and both V₁ and V₂ are bonds.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein W is S, S(O) or S(O)₂.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein W is absent.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Q is NR₆.

In some embodiments, R_6 is H.

In some embodiments, R_6 is C_{1-8} alkyl.

In some embodiments, R_6 is selected from the group consisting of methyl, ethyl, isopropyl, and *n*-propyl.

5 In some embodiments, R_6 is isopropyl.

In some embodiments, R_6 is C_{3-7} cycloalkyl.


In some embodiments, R_6 is selected from the group consisting of cyclopropyl, cyclobutyl, and cyclopentyl.

In some embodiments, R_6 is cyclopropyl

10 In some embodiments, Q is NH.

In some embodiments, R_6 is cyclopropylmethyl.


Some embodiments of the present invention pertain to compounds of Formula (I) wherein Q is NH and can be represented by Formula (Ie) as illustrated below:



15 wherein each variable in Formula (Ie) has the same meaning as described herein, *supra* and *infra*. In some embodiments, V_2 is a bond.

In some embodiments, Q is O.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Q is O (i.e., an oxygen atom) and can be represented by Formula (Ig) as illustrated below:



20 wherein each variable in Formula (Ig) has the same meaning as described herein, *supra* and *infra*. In some embodiments, V_2 is a bond. In some embodiments, V_2 is $-CH_2-$. In still further embodiments, V_2 is $-CH_2CH_2-$.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Q is S, S(O) or S(O)₂.

25 In some embodiments, Q is S.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein V_1 is a bond.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein V_2 is a bond.

5 In some embodiments, V_1 and V_2 are both a bond.


Some embodiments of the present invention pertain to compounds of Formula (I) wherein V_2 is $-\text{CH}_2-$.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein V_2 is $-\text{CH}_2\text{CH}_2-$.

10 Some embodiments of the present invention pertain to compounds of Formula (I) wherein A and B are independently C_{1-2} alkylene optionally substituted with 1 to 4 substituents selected from the group consisting of C_{1-3} alkyl, C_{1-4} alkoxy, carboxy, cyano, C_{1-3} haloalkyl and halogen.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein both A and B are C_1 alkylene groups wherein A and B are optionally substituted with 1 to 2 methyl groups.

15 In some embodiments, A and B are both $-\text{CH}_2-$. In some embodiments, compounds of the present invention can be represented by Formula (Ik) as shown below:



wherein each variable in Formula (Ik) has the same meaning as described herein, *supra* and *infra*.

In some embodiments, both A and B are $-\text{CH}_2-$ and E is CH.

20 In some embodiments, both A and B are $-\text{CH}_2-$, E is CH, and D is N-R_2 .


In some embodiments, A is $-\text{CH}_2\text{CH}_2-$ and B is $-\text{CH}_2-$.

In some embodiments, A is $-\text{CH}_2\text{CH}_2-$ and B is $-\text{CH}_2-$, and E is CH.

In some embodiments, A is $-\text{CH}_2\text{CH}_2-$ and B is $-\text{CH}_2-$, E is CH and D is N-R_2 .


25 Some embodiments of the present invention pertain to compounds of Formula (I) wherein A is a C_1 alkylene group and B is a C_2 alkylene group wherein A is optionally substituted with 1 to 2 methyl groups and B is optionally substituted with 1 to 4 methyl groups.

In some embodiments, compounds of the present invention can be represented by Formulae (Im) and (In) respectively as shown below:




wherein each variable in Formulae (Im) and (In) has the same meaning as described herein, *supra* and *infra*. In some embodiments, A is -CH₂- and B is -CH₂CH₂-. In further embodiments, A is -CH₂-, B is -CH₂CH₂-, and V₂ is -CH₂- or -CH₂CH₂-.

- 5 Some embodiments of the present invention pertain to compounds of Formula (I) wherein A is a C₁ alkylene group and B is a C₃ alkylene group wherein A is optionally substituted with 1 to 2 methyl groups and B is optionally substituted with 1 to 4 methyl groups. In some embodiments, A is -CH₂- or -CH- and B is -CH₂CH₂CH₂- and can be represented by Formulae (Ip) and (Iq) respectively as shown below:






- 10 wherein each variable in Formulae (Ip) and (Iq) has the same meaning as described herein, *supra* and *infra*.

- 15 Some embodiments of the present invention pertain to compounds of Formula (I) wherein A is a C₂ alkylene group and B is a C₁ alkylene group wherein A is optionally substituted with 1 to 4 methyl groups and B is optionally substituted with 1 to 2 methyl groups. In some embodiments, A is -CHCH₂- and B is -CH₂-; these embodiments can be represented by Formula (It) as shown below:




wherein each variable in Formula (It) has the same meaning as described herein, *supra* and *infra*.

- 20 Some embodiments of the present invention pertain to compounds of Formula (I) wherein A is CH₂ and B is -CH₂CH₂-, -CH₂CH(CH₃)-, -CH(CH₃)CH₂-, -CH₂CH(CF₃)- or -CH(CF₃)CH₂-. In some embodiments, compounds of the invention are represented by Formulae (Iv), (Iw) and (Ix) as shown below:



wherein each variable in Formulae (Iv), (Iw) and (Ix) has the same meaning as described herein, *supra* and *infra*. In some embodiments, D is $N-R_2$. In further embodiments, D is $N-R_2$ wherein R_2 is represented by Formula (D). In still further embodiments, D is $N-R_2$ wherein R_2 is $-C(O)OC_{1-8}$ alkyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein A is a C_3 alkylene group and B is a C_1 alkylene group wherein A is optionally substituted with 1 to 4 methyl groups and B is optionally substituted with 1 to 2 methyl groups. In some embodiments, A is $-CHCH_2CH_2-$ and B is $-CH_2-$ and represented by Formulae (IIa) as shown below:



wherein each variable in Formulae (IIa) has the same meaning as described herein, *supra* and *infra*.


In some embodiments, both A and B are $-CH_2CH_2-$.

In some embodiments, both A and B are $-CH_2CH_2-$ and E is CH .


In some embodiments, both A and B are $-CH_2CH_2-$, E is CH , and D is $N-R_2$.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein both A and B are C_2 alkylene groups wherein A and B are optionally substituted with 1 to 4 methyl groups.

In some embodiments, A is $-CH_2CH_2-$ or $-CHCH_2-$ and B is $-CH_2CH_2-$. In some embodiments, compounds of the present invention can be represented by Formulae (IIc) and (IId) as shown below:




wherein each variable in Formulae (IIc) and (IId) has the same meaning as described herein, *supra* and *infra*. In some embodiments, A and B are both $-CH_2CH_2-$, D is $N-R_2$, and E is CR_4 ; these embodiments are represented by Formula (IIf) as shown below:




wherein each variable in Formula (IIf) has the same meaning as described herein, *supra* and *infra*. In some embodiments, compounds have the Formula (IIf) and R_4 is H. In further embodiment, V_2 is a bond. In still further embodiments, V_2 is $-CH_2-$ or $-CH_2CH_2-$.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein A is a C_2 alkylene group and B is a C_3 alkylene group wherein A and B are optionally substituted with 1 to 4 methyl groups. In some embodiments, A is $-CH_2CH_2-$ or $-CHCH_2-$ and B is $-CH_2CH_2CH_2-$ and can be represented by Formulae (IIh) and (IIi) as shown below:




wherein each variable in Formulae (IIh) and (IIi) has the same meaning as described herein, *supra* and *infra*.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein A is a C_3 alkylene group and B is a C_2 alkylene group wherein A and B are optionally substituted with 1 to 4 methyl groups. In some embodiments, A is $-CHCH_2CH_2-$ and B is $-CH_2CH_2-$; these embodiments can be represented by Formula (IIk) as shown below:



wherein each variable in Formula (IIk) has the same meaning as described herein, *supra* and *infra*.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein both A and B are C₃ alkylene groups wherein A and B are optionally substituted with 1 to 4 methyl groups. In some embodiments, A is -CH₂CH₂CH₂- or -CHCH₂CH₂- and B is -CH₂CH₂CH₂- and are represented by Formula (IIm) and (IIn) respectively as shown below:




wherein each variable in Formulae (IIm) and (IIn) has the same meaning as described herein, *supra* and *infra*.


Some embodiments of the present invention pertain to compounds of Formula (I) wherein - is a single bond.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein E is a nitrogen atom (i.e., N).

Some embodiments of the present invention pertain to compounds of Formula (I) wherein E is CR₄. In some embodiments, R₄ is H and can be represented by Formula (IIp) as shown below:



wherein each variable in Formula (IIp) has the same meaning as described herein, *supra* and *infra*. In further embodiments, V₂ is a bond and represented by Formula (IIr):




wherein each variable in Formula (IIr) has the same meaning as described herein, *supra* and *infra*. In some embodiments, compounds of the present invention are of Formula (IIr) and Q is NH. In some embodiments, compounds are of Formula (IIr) and Q is O (i.e., an oxygen atom).

- 5 Some embodiments of the present invention pertain to compounds of Formula (I) wherein $\text{--}\text{--}$ is a double bond. It is understood that when $\text{--}\text{--}$ is a double bond then E is C (i.e., a carbon atom) and not N (i.e., a nitrogen atom).

Some embodiments of the present invention pertain to compounds of Formula (I) wherein V_2 is a CH_2 or CH_2CH_2 group.

- 10 Some embodiments of the present invention pertain to compounds of Formula (I) wherein V_1 is a bond and V_2 is a CH_2 or CH_2CH_2 group.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein D is CR_2R_3 and can be represented by Formula (IIIt) as shown below:





- 15 wherein each variable in Formula (IIIt) has the same meaning as described herein, *supra* and *infra*. In some embodiments, compounds of the invention are of Formula (IIIt) and R_2 is selected from the group consisting of H, amino, carboxamide, carboxy, cyano, C_{3-6} -cycloalkyl, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, halogen and hydroxyl. In some embodiments, R_2 is $-\text{NR}_{23}\text{C}(\text{O})\text{O}-(\text{C}_{1-8} \text{ alkyl})$ or $-\text{OC}(\text{O})\text{NR}_{23}-(\text{C}_{1-8} \text{ alkyl})$. In some embodiments, R_2 is selected from the group consisting of OCH_3 , OCH_2CH_3 , $\text{OCH}_2\text{CH}_2\text{CH}_3$, $\text{OCH}(\text{CH}_3)_2$, $\text{OCH}_2(\text{CH}_2)_2\text{CH}_3$, amino, carboxamide, carboxy, cyano, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, OCF_3 , OCHF_2 , CF_3 , CHF_2 and F. In some
- 20 embodiments, R_2 is C_{1-8} alkyl, aryl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-5} acyl, C_{1-5} acyloxy, C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylamino, C_{1-4} alkylcarboxamide, C_{1-4} alkylthiocarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4}
- 25 alkylsulfonyl, C_{1-4} alkylthio, C_{1-4} alkylthiourey, C_{1-4} alkylureyl, amino, carbo- C_{1-6} -alkoxy, carboxamide, carboxy, cyano, C_{3-6} -cycloalkyl, C_{3-6} -cycloalkyl- C_{1-3} -heteroalkylene, C_{2-8} dialkylamino, C_{2-6} dialkylcarboxamide, C_{2-6} dialkylthiocarboxamide, C_{2-6} dialkylsulfonamide, C_{1-4} alkylthiourey, C_{1-}

- haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino and nitro. In some embodiments, R₂ is selected from the group consisting of CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, CH(CH₃)(CH₂CH₃), CH₂(CH₂)₂CH₃, CH₂(CH₂)₃CH₃. In some embodiments, R₂ is selected from the group consisting of
- 5 CH₂OCH₃, CH₂CH₂OCH₃, CH₂OCH₂CH₃, CH₂OCH₂CH₂CH₃, CH₂CH₂OCH₂CH₃, CH₂CH₂OCH₂CH₂CH₃, CH₂OCH(CH₃)₂, CH₂OCH₂CH(CH₃)₂, CH₂CO₂H, CH₂CH₂CO₂H, CH₂OH, CH₂CH₂OH and CH₂CH₂CH₂OH. In some embodiments, R₂ is selected from the group consisting of
- 10 CH₂SCH₃, CH₂SCH₂CH₃, CH₂SCH₂CH₂CH₃, CH₂SCH(CH₃)₂, CH₂SCH₂(CH₂)₂CH₃, CH₂CH₂SCH₃, CH₂CH₂SCH₂CH₃, CH₂CH₂SCH₂CH₂CH₃, CH₂CH₂SCH(CH₃)₂, CH₂CH₂SCH₂(CH₂)₂CH₃, CH₂S(O)CH₃, CH₂S(O)CH₂CH₃, CH₂S(O)CH₂CH₂CH₃, CH₂S(O)CH(CH₃)₂, CH₂S(O)CH₂(CH₂)₂CH₃, CH₂CH₂S(O)CH₃, CH₂CH₂S(O)CH₂CH₃, CH₂CH₂S(O)CH₂CH₂CH₃, CH₂CH₂S(O)CH(CH₃)₂, CH₂CH₂S(O)CH₂(CH₂)₂CH₃, CH₂S(O)₂CH₃, CH₂S(O)₂CH₂CH₃, CH₂S(O)₂CH₂CH₂CH₃, CH₂S(O)₂CH(CH₃)₂, CH₂S(O)₂CH₂(CH₂)₂CH₃, CH₂CH₂S(O)₂CH₃, CH₂CH₂S(O)₂CH₂CH₃, CH₂CH₂S(O)₂CH₂CH₂CH₃, CH₂CH₂S(O)₂CH(CH₃)₂ and CH₂CH₂S(O)₂CH₂(CH₂)₂CH₃. In some
- 15 embodiments, R₂ is selected from the group consisting of CH₂OCH₂-cyclopropyl, CH₂OCH₂-cyclobutyl, CH₂OCH₂-cyclopentyl, CH₂OCH₂-cyclohexyl, CH₂OCH₂CH₂-cyclopropyl, CH₂OCH₂CH₂-cyclobutyl, CH₂OCH₂CH₂-cyclopentyl, CH₂OCH₂CH₂-cyclohexyl, CH₂CH₂OCH₂-cyclopropyl, CH₂CH₂OCH₂-cyclobutyl, CH₂CH₂OCH₂-cyclopentyl, CH₂CH₂OCH₂-cyclohexyl, CH₂CH₂OCH₂CH₂-cyclopropyl, CH₂CH₂OCH₂CH₂-cyclobutyl, CH₂CH₂OCH₂CH₂-cyclopentyl and
- 20 CH₂CH₂OCH₂CH₂-cyclohexyl. In some embodiments, R₂ is selected from the group consisting of 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,3,4-oxadiazol-2-yl, 1,2,4-triazol-5-yl and 1,2,4-triazol-1-yl, 3-methyl-1,2,4-oxadiazol-5-yl, 3-methyl-1,2,4-oxadiazol-5-yl, 3-ethyl-1,2,4-oxadiazol-5-yl, 3-ethyl-1,2,4-oxadiazol-5-yl, 5-methyl-1,3,4-oxadiazol-2-yl, 5-ethyl-1,3,4-oxadiazol-2-yl, 3-methyl-1,2,4-triazol-5-yl, 3-ethyl-1,2,4-triazol-5-yl, 3-methyl-1,2,4-triazol-1-yl, 3-ethyl-1,2,4-triazol-1-yl, 5-
- 25 methyl-1,2,4-triazol-1-yl and 5-ethyl-1,2,4-triazol-1-yl.

In some embodiments R₂ is a heteroaryl comprising 5-atoms in the aromatic ring and are represented by the following formulae:


TABLE 2A





wherein the 5-membered heteroaryl is bonded at any available position of the ring, for example, a imidazolyl ring can be bonded at one of the ring nitrogens (i.e., imidazol-1-yl group) or at one of the ring carbons (i.e., imidazol-2-yl, imidazol-4-yl or imidazol-5-yl group). In some embodiments R_2 is a 5-membered heteroaryl, for example but not limited to those shown in TABLE 2A, optionally substituted with 1 to 4 substituents selected from the group consisting of C_{1-5} acyl, C_{1-5} acyloxy, C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylamino, C_{1-4} alkylcarboxamide, C_{1-4} alkylthiocarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, C_{1-4} alkylthiourey, C_{1-4} alkylureyl, amino, carbo- C_{1-6} -alkoxy, carboxamide, carboxy, cyano, C_{3-6} -cycloalkyl, C_{3-6} -cycloalkyl- C_{1-3} -heteroalkylene, C_{2-8} dialkylamino, C_{2-6} dialkylcarboxamide, C_{2-6} dialkylthiocarboxamide, C_{2-6} dialkylsulfonamide, C_{1-4} alkylthiourey, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkylsulfinyl, C_{1-4} haloalkylsulfonyl, C_{1-4} haloalkyl, C_{1-4} haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino and nitro.


In some embodiments R_2 is a heteroaryl comprising 5-atoms in the aromatic ring and are represented by the following formulae:




wherein the 5-membered heteroaryl is bonded at any available position of the ring as described above. In some embodiments, R_2 is a 5-membered heteroaryl optionally substituted with 1 to 4 substituents selected from the group consisting of C_{1-5} acyl, C_{1-5} acyloxy, C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylamino, C_{1-4} alkylcarboxamide, C_{1-4} alkylthiocarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, C_{1-4} alkylthiourey, C_{1-4} alkylureyl, amino, carbo- C_{1-6} -alkoxy, carboxamide, carboxy, cyano, C_{3-6} -cycloalkyl- C_{1-3} -heteroalkylene, C_{2-8} dialkylamino, C_{2-6} dialkylcarboxamide, C_{2-6} dialkylthiocarboxamide, C_{2-6} dialkylsulfonamide, C_{1-4} alkylthiourey, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkylsulfinyl, C_{1-4} haloalkylsulfonyl, C_{1-4} haloalkyl, C_{1-4} haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino and nitro.

In some embodiments R_2 is a heterocyclic group represented, for example, by the formulae in TABLE 2B.

TABLE 2B




- 5 It is understood that any one of the heterocyclic groups shown in TABLES 2B to 2E may be bonded at any ring carbon or ring nitrogen as allowed by the respective formula unless otherwise specified. For example, a 2,5-dioxo-imidazolidinyl group may be bonded at the ring carbon or at either of the two ring nitrogens to give the following formulae respectively:




- 10 In some embodiments R_2 is a heterocyclic represented, for example, by the formulae in TABLE 2C.

TABLE 2C




- 15 In some embodiments R_2 is a heterocyclic represented, for example, by the formulae in TABLE 2D.

TABLE 2D




In some embodiments R_2 is a heterocyclic represented, for example, by the formulae in TABLE 2E.

TABLE 2E



In some embodiments R_2 is a heterocyclic represented, for example, by the formulae in TABLE 2F wherein the C_{1-6} alkyl group on the respective ring nitrogen atoms may be the same or different.


TABLE 2F



10

In some embodiments R_2 is a heterocyclic represented, for example, by the formulae in TABLE 2G wherein the C_{1-6} alkyl group on the respective ring nitrogen atoms may be the same or different.

TABLE 2G



15


Some embodiments of the present invention pertain to compounds of Formula (III) and R_2 is - Ar_2-Ar_3 wherein Ar_2 and Ar_3 are independently aryl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of H, C_{1-5} acyl, C_{1-5} acyloxy, C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylcarboxamide, C_{1-4} alkylthiocarboxamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, amino, carbo- C_{1-6} -alkoxy, carboxamide, carboxy, cyano, C_{3-6} -cycloalkyl, C_{2-6} dialkylcarboxamide, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, halogen, hydroxyl and nitro. In some embodiments, Ar_2 is a heteroaryl and Ar_3 is phenyl. In some embodiments, the heteroaryl and said phenyl are optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylcarboxamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, cyano, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, halogen, hydroxyl and nitro.

20

In some embodiments Ar_2 is a heteroaryl comprising 5-atoms in the aromatic ring and are represented by the following formulae shown in TABLE 3.

25

TABLE 3




5

10

15

wherein the 5-membered heteroaryl is bonded at any position of the ring, for example, a imidazolyl ring can be bonded at one of the ring nitrogens (i.e., imidazol-1-yl group) or at one of the ring carbons (i.e., imidazol-2-yl, imidazol-4-yl or imiadazol-5-yl group) and Ar₃ is bonded to any remaining available ring atom. In some embodiments Ar₂ is a heteroaryl and Ar₃ is phenyl. In some embodiments, Ar₂ is a phenyl and Ar₃ is heteroaryl (such as a heteroaryl selected from TABLE 2A, *supra*). In some embodiments the heteroaryl and phenyl are optionally substituted with 1 to 5 substituents selected from the group consisting of H, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, hydroxyl and nitro.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein D is CR₂R₃, or R₂ is Formula (C):




wherein:

20

R₁₇ is C₁₋₈ alkyl or C₃₋₇ cycloalkyl; and R₁₈ is F, Cl, Br or CN. In some embodiments, R₁₇ is C₁₋₈ alkyl and R₁₈ is F, Cl or CN.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein D is CR₂R₃ and R₂ is Formula (D):



wherein:

G is C(O), C(O)NR₂₃, C(O)O, OC(O), C(S), C(S)NR₂₃, C(S)O, OC(S), CR₂₃R₂₄, O, S, S(O) or S(O)₂; where R₂₃ and R₂₄ are independently H or C₁₋₈ alkyl; and R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro.


Some embodiments of the present invention pertain to compounds of Formula (I) wherein R₂ is Formula (D) and G is C(O), C(O)NR₂₃, C(O)O, OC(O), C(S), C(S)NR₂₃, C(S)O, OC(S) or CR₂₃R₂₄. In some embodiments, R₂₂ is C₁₋₈ alkyl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carboxamide, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, halogen and hydroxyl. In some embodiments, Formula (D) (i.e., -G-R₂₂) is selected from the group consisting of C(O)CH₃, C(O)CH₂CH₃, C(O)CH₂CH₂CH₃, C(O)CH(CH₃)₂, C(O)CH₂CH₂CH₂CH₃, C(O)C(CH₃)₃, C(O)CH₂C(CH₃)₃, CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, CH(CH₃)(CH₂CH₃), CH₂(CH₂)₂CH₃, C(CH₃)₃, CH₂(CH₂)₃CH₃, C(O)NHCH₃, C(O)NHCH₂CH₃, C(O)NHCH₂CH₂CH₃, C(O)NHCH(CH₃)₂, C(O)NHCH₂(CH₂)₂CH₃, C(O)N(CH₃)₂, C(O)N(CH₃)CH₂CH₃, C(O)NH(CH₂CH₃)₂, CO₂CH₃, CO₂CH₂CH₃, CO₂CH₂CH₂CH₃, CO₂CH(CH₃)₂ and CO₂CH₂(CH₂)₂CH₃.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein R₂ is Formula (D) and G is C(O), C(O)NR₂₃, C(O)O, OC(O), C(S), C(S)NR₂₃, C(S)O, OC(S) or CR₂₃R₂₄. In some embodiments, R₂₂ is phenyl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carboxamide, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, halogen and hydroxyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein R₂ is Formula (D) and G is C(O), C(O)NR₂₃, C(O)O, OC(O), C(S), C(S)NR₂₃, C(S)O, OC(S) or CR₂₃R₂₄. In some embodiments, R₂₂ is heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carboxamide, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, halogen and hydroxyl. In some embodiments, R₂₂ is a 5-membered heteroaryl, for example, as shown in TABLE 2A *supra*. In some embodiments, R₂₂ is 1-methyl-1H-imidazole-4-yl or 2,4-dimethyl-thiazole-5-yl.

In some embodiments, R_{22} is a 6-membered heteroaryl, for example, the 6-membered heteroaryls as shown in TABLE 4:

TABLE 4



wherein the heteroaryl group is bonded at any ring carbon. In some embodiments R_{22} is selected from the group consisting of pyridinyl, pyridazinyl, pyrimidinyl and pyrazinyl. In some embodiments, R_{22} is pyridinyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein R_{23} and R_{24} are independently H or C_{1-2} alkyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein R_2 is Formula (D) and G is O, S, S(O) or S(O)₂. In some embodiments, R_{22} is C_{1-8} alkyl optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-5} acyl, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylcarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, carboxamide, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkylsulfinyl, C_{1-4} haloalkylsulfonyl, C_{1-4} haloalkyl, halogen and hydroxyl. In some embodiments, Formula (D) (i.e., -G- R_{22}) is selected from the group consisting of: OCH₃, OCH₂CH₃, OCH₂CH₂CH₃, OCH(CH₃)₂, OCH₂(CH₂)₂CH₃, SCH₃, SCH₂CH₃, SCH₂CH₂CH₃, SCH(CH₃)₂, SCH₂(CH₂)₂CH₃, S(O)CH₃, S(O)CH₂CH₃, S(O)CH₂CH₂CH₃, S(O)CH(CH₃)₂, S(O)CH₂(CH₂)₂CH₃, S(O)₂CH₃, S(O)₂CH₂CH₃, S(O)₂CH₂CH₂CH₃, S(O)₂CH(CH₃)₂ and S(O)₂CH₂(CH₂)₂CH₃.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein R_2 is Formula (D) and G is O, S, S(O) or S(O)₂. In some embodiments, R_{22} is phenyl optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-5} acyl, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylcarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, carboxamide, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkylsulfinyl, C_{1-4} haloalkylsulfonyl, C_{1-4} haloalkyl, halogen and hydroxyl.


Some embodiments of the present invention pertain to compounds of Formula (I) wherein R_2 is Formula (D) and G is O, S, S(O) or S(O)₂. In some embodiments, R_{22} is heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-5} acyl, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylcarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, carboxamide, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkylsulfinyl, C_{1-4} haloalkylsulfonyl, C_{1-4} haloalkyl, halogen and hydroxyl. In some embodiments, R_{22} is a 5-membered heteroaryl, for

example, as shown in TABLE 2A *supra*. In some embodiments, R₂₂ is a 6-membered heteroaryl, as shown in TABLE 4 *supra*. In some embodiments R₂₂ is selected from the group consisting of pyridinyl, pyridazinyl, pyrimidinyl and pyrazinyl. In some embodiments, R₂₂ is pyridinyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein R₃ is H.

In some embodiments, D is N-R₂.

Some embodiments of the present invention pertain to compounds wherein D is N-R₂ and is represented by Formula (IIv):



wherein each variable in Formula (IIv) has the same meaning as described herein, *supra* and *infra*.

In some embodiments, R₂ is C₁₋₈ alkyl, aryl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₃₋₆-cycloalkyl-C₁₋₃-heteroalkylene, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino and nitro. In some embodiments, R₂ is pyridyl. In some embodiments, R₂ is 2-pyridyl. In some embodiments, R₂ is selected from the group consisting of CH₂CH₂C(CH₃)₃, CH₂CH₂CH(CH₃)₂ and CH₂(CH₂)₄CH₃. In some embodiments, R₂ is selected from the group consisting of: CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, CH(CH₃)(CH₂CH₃), CH₂(CH₂)₂CH₃ and CH₂(CH₂)₃CH₃. In some embodiments, R₂ is selected from the group consisting of CH₂OCH₃, CH₂CH₂OCH₃, CH₂OCH₂CH₃, CH₂OCH₂CH₂CH₃, CH₂CH₂OCH₂CH₃, CH₂CH₂OCH₂CH₂CH₃, CH₂OCH(CH₃)₂, CH₂OCH₂CH(CH₃)₂, CH₂CO₂H, CH₂CH₂CO₂H, CH₂OH, CH₂CH₂OH and CH₂CH₂CH₂OH. In some embodiments, R₂ is selected from the group consisting of CH₂SCH₃, CH₂SCH₂CH₃, CH₂SCH₂CH₂CH₃, CH₂SCH(CH₃)₂, CH₂SCH₂(CH₂)₂CH₃, CH₂CH₂SCH₃, CH₂CH₂SCH₂CH₃, CH₂CH₂SCH₂CH₂CH₃, CH₂CH₂SCH(CH₃)₂, CH₂CH₂SCH₂(CH₂)₂CH₃, CH₂S(O)CH₃, CH₂S(O)CH₂CH₃, CH₂S(O)CH₂CH₂CH₃, CH₂S(O)CH(CH₃)₂, CH₂S(O)CH₂(CH₂)₂CH₃, CH₂CH₂S(O)CH₃, CH₂CH₂S(O)CH₂CH₃, CH₂CH₂S(O)CH₂CH₂CH₃, CH₂CH₂S(O)CH(CH₃)₂, CH₂CH₂S(O)CH₂(CH₂)₂CH₃, CH₂S(O)₂CH₃, CH₂S(O)₂CH₂CH₃, CH₂S(O)₂CH₂CH₂CH₃, CH₂S(O)₂CH(CH₃)₂, CH₂S(O)₂CH₂(CH₂)₂CH₃,


CH₂CH₂S(O)₂CH₃, CH₂CH₂S(O)₂CH₂CH₃, CH₂CH₂S(O)₂CH₂CH₂CH₃, CH₂CH₂S(O)₂CH(CH₃)₂ and CH₂CH₂S(O)₂CH₂(CH₂)₂CH₃. In some embodiments, R₂ is CH₂-cyclopropyl. In some embodiments, R₂ is selected from the group consisting of CH₂OCH₂-cyclopropyl, CH₂OCH₂-cyclobutyl, CH₂OCH₂-cyclopentyl, CH₂OCH₂-cyclohexyl, CH₂OCH₂CH₂-cyclopropyl, CH₂OCH₂CH₂-cyclobutyl, CH₂OCH₂CH₂-cyclopentyl, CH₂OCH₂CH₂-cyclohexyl, CH₂CH₂OCH₂-cyclopropyl, CH₂CH₂OCH₂-cyclobutyl, CH₂CH₂OCH₂-cyclopentyl, CH₂CH₂OCH₂-cyclohexyl, CH₂CH₂OCH₂CH₂-cyclopropyl, CH₂CH₂OCH₂CH₂-cyclobutyl, CH₂CH₂OCH₂CH₂-cyclopentyl and CH₂CH₂OCH₂CH₂-cyclohexyl. In some embodiments, R₂ is selected from the group consisting of 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,3,4-oxadiazol-2-yl, 1,2,4-triazol-5-yl and 1,2,4-triazol-1-yl, 3-methyl-1,2,4-oxadiazol-5-yl, 3-methyl-1,2,4-oxadiazol-5-yl, 3-ethyl-1,2,4-oxadiazol-5-yl, 3-ethyl-1,2,4-oxadiazol-5-yl, 5-methyl-1,3,4-oxadiazol-2-yl, 5-ethyl-1,3,4-oxadiazol-2-yl, 3-methyl-1,2,4-triazol-5-yl, 3-ethyl-1,2,4-triazol-5-yl, 3-methyl-1,2,4-triazol-1-yl, 3-ethyl-1,2,4-triazol-1-yl, 5-methyl-1,2,4-triazol-1-yl and 5-ethyl-1,2,4-triazol-1-yl.

In some embodiments, compounds are of Formula (IIv) and R₂ is a heteroaryl comprising 5-atoms in the aromatic ring selected from the group shown in Table 2A. In some embodiments R₂ is a 5-membered heteroaryl optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₃₋₆-cycloalkyl-C₁₋₃-heteroalkylene, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino and nitro. In some embodiments R₂ is a heteroaryl selected from the group, but not limited to the group, in TABLE 2A. In some embodiments, R₂ is a 5-membered heteroaryl optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl-C₁₋₃-heteroalkylene, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino and nitro. In some embodiments R₂ is a heterocyclic group selected from the groups shown in TABLE 2B to TABLE 2G.

Some embodiments of the present invention pertain to compounds of Formula (IIv) and R₂ is -Ar₂-Ar₃ wherein Ar₂ and Ar₃ are independently aryl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of H, C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio,

amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₆ dialkylcarboxamide, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, hydroxyl and nitro. In some embodiments Ar₂ is a heteroaryl comprising 5-atoms in the aromatic ring and selected from the group shown in TABLE 3. In some embodiments Ar₂ is a heteroaryl and Ar₃ is phenyl. In some embodiments, Ar₂ is a phenyl and Ar₃ is heteroaryl (such as a heteroaryl selected from TABLE 2A or TABLE 4, *supra*). In some embodiments, the heteroaryl and phenyl are optionally substituted with 1 to 5 substituents selected from the group consisting of H, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, hydroxyl and nitro.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein D is N-R₂. In some embodiments, R₂ is Formula (C):




wherein:

R₁₇ is C₁₋₈ alkyl or C₃₋₇ cycloalkyl; and R₁₈ is F, Cl, Br or CN. In some embodiments, R₁₇ is C₁₋₈ alkyl and R₁₈ is F, Cl or CN.

In some embodiments, R₂ is selected from the group consisting of methoxycarbonyl, ethoxycarbonyl, *iso*-propoxycarbonyl, *n*-propoxycarbonyl, *n*-butoxycarbonyl, *tert*-butoxycarbonyl, *iso*-butoxycarbonyl, and *n*-pentyloxycarbonyl. In some embodiments, R₂ is *iso*-propoxycarbonyl or *tert*-butoxycarbonyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein D is N-R₂. In some embodiments, R₂ is Formula (D):



wherein:


G is C(O), C(O)NR₂₃, C(O)O, C(S), C(S)NR₂₃, C(S)O, CR₂₃R₂₄ or S(O)₂; where R₂₃ and R₂₄ are independently H or C₁₋₈ alkyl; and R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein D is N-R₂. In some embodiments, R₂ is Formula (D) and R₂₂ is C₁₋₈ alkyl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carboxamide, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, halogen and hydroxyl. In some embodiments, D is N-R₂ where R₂ is of Formula (D) (i.e., -G-R₂₂) and -G-R₂₂ is selected from the group consisting of C(O)CH₃, C(O)CH₂CH₃, C(O)CH₂CH₂CH₃, C(O)CH(CH₃)₂, C(O)CH₂CH₂CH₂CH₃, C(O)C(CH₃)₃, C(O)CH₂C(CH₃)₃, CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, CH(CH₃)(CH₂CH₃), CH₂(CH₂)₂CH₃, C(CH₃)₃, CH₂(CH₂)₃CH₃, C(O)NHCH₃, C(O)NHCH₂CH₃, C(O)NHCH₂CH₂CH₃, C(O)NHCH(CH₃)₂, C(O)NHCH₂(CH₂)₂CH₃, C(O)N(CH₃)₂, C(O)N(CH₃)CH₂CH₃, C(O)NH(CH₂CH₃)₂, CO₂CH₃, CO₂CH₂CH₃, CO₂CH₂CH₂CH₃, CO₂CH(CH₃)₂ and CO₂CH₂(CH₂)₂CH₃.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein D is N-R₂. In some embodiments, R₂ is Formula (D) and R₂₂ is phenyl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carboxamide, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, halogen and hydroxyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein D is N-R₂. In some embodiments, R₂ is Formula (D) and R₂₂ is heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carboxamide, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, halogen and hydroxyl. In some embodiments, R₂₂ is selected from the group consisting of pyridinyl, pyridazinyl, pyrimidinyl and pyrazinyl. In some embodiments, R₂₂ is pyridinyl.

In some embodiments, R₂ is a group of Formula (D):



(D)

wherein:

G is -CR₂₃R₂₄C(O)-, -C(O)-, -CR₂₃R₂₄C(O)NR₂₃-, -C(O)NR₂₃-, -C(O)O-, -C(S)-, -C(S)NR₂₃-, -C(S)O-, -CR₂₃R₂₄-, -S(O)₂-, or a bond; wherein R₂₃ and R₂₄ are each independently H or C₁₋₈ alkyl; and

R₂₂ is H, C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄

alkylureyl, amino, carbo- C_{1-6} -alkoxy, carboxamide, carboxy, cyano, C_{3-7} cycloalkyl, C_{2-8} dialkylamino, C_{2-6} dialkylcarboxamide, C_{2-6} dialkylthiocarboxamide, C_{2-6} dialkylsulfonamide, C_{1-4} alkylthiourey, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkylsulfinyl, C_{1-4} haloalkylsulfonyl, C_{1-4} haloalkyl, C_{1-4} haloalkylthio, halogen, heteroaryl, heterocyclic, hydroxyl, hydroxylamino, nitro, phenyl, phenoxy, and sulfonic acid, wherein said C_{1-7} alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-5} acyl, C_{1-5} acyloxy, C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylamino, C_{1-4} alkylcarboxamide, C_{1-4} alkylthiocarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, C_{1-4} alkylthiourey, C_{1-4} alkylureyl, amino, carbo- C_{1-6} -alkoxy, carboxamide, carboxy, cyano, C_{3-7} cycloalkyl, C_{2-8} dialkylamino, C_{2-6} dialkylcarboxamide, C_{2-6} dialkylthiocarboxamide, C_{2-6} dialkylsulfonamide, C_{1-4} alkylthiourey, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkylsulfinyl, C_{1-4} haloalkylsulfonyl, C_{1-4} haloalkyl, C_{1-4} haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino, and nitro.

In some embodiments, Formula (D) is $-C(O)OR_{22}$.

In some embodiments, Formula (D) is $-C(O)R_{22}$.

15 In some embodiments, Formula (D) is $-CR_{23}R_{24}-R_{22}$.

In some embodiments, Formula (D) is $-R_{22}$ (i.e., -G- is a bond).

In some embodiments, Formula (D) is $-S(O)_2R_{22}$.

In some embodiments, Formula (D) is $-CR_{23}R_{24}C(O)R_{22}$.

In some embodiments, Formula (D) is $-CR_{23}R_{24}C(O)NR_{25}R_{22}$.

20 In some embodiments, R_2 is $-C(O)OR_{22}$ and R_{22} is C_{1-8} alkyl, C_{3-7} cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{2-6} alkenyl, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylsulfonyl, amino, carbo- C_{1-6} -alkoxy, carboxy, cyano, C_{3-7} cycloalkyl, C_{2-8} dialkylamino, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C_{1-7} alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C_{1-4} haloalkoxy, and heterocyclic.

30 In some embodiments, R_2 is $-C(O)OR_{22}$ and R_{22} is C_{1-8} alkyl, or C_{3-7} cycloalkyl each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylsulfonyl, carboxy, cyano, C_{3-7} cycloalkyl, C_{2-8} dialkylamino, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid.

In some embodiments, R_2 is $-C(O)OR_{22}$ and R_{22} is C_{1-8} alkyl, or C_{3-7} cycloalkyl wherein said C_{3-7} cycloalkyl is optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-4} alkoxy, C_{1-7} alkyl, carboxy, C_{2-8} dialkylamino, and halogen.

In some embodiments, R_2 is $-C(O)OR_{22}$ and R_{22} is C_{1-8} alkyl, or C_{3-7} cycloalkyl.

35 In some embodiments, R_2 is $-C(O)R_{22}$ and R_{22} is C_{1-8} alkyl, C_{3-7} cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{2-6} alkenyl, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylsulfonyl, amino, carbo- C_{1-6} -alkoxy,

carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic.

5 In some embodiments, R₂ is -C(O)R₂₂ and R₂₂ is C₁₋₈ alkyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of H, C₁₋₄ alkoxy, C₁₋₇ alkyl, amino, carboxy, halogen, heteroaryl, hydroxyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl and phenoxy are optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic.

10 In some embodiments, R₂ is -CH₂R₂₂, or -R₂₂ and R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic.

In some embodiments, R₂ is -CH₂R₂₂, or -R₂₂, and R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, or heteroaryl each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₂₋₆ alkenyl, C₁₋₄ alkoxy, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, and hydroxyl.

20 R₂ is -S(O)₂R₂₂ and R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic.

In some embodiments, R₂ is -S(O)₂R₂₂ and R₂₂ is C₁₋₈ alkyl, or heteroaryl and said heteroaryl is optionally substituted with 1 to 5 C₁₋₇ alkyl.


30 In some embodiments, R₂ is -CR₂₃R₂₄C(O)R₂₂ and wherein R₂₃ and R₂₄ are each independently H or C₁₋₈ alkyl; and R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic.

In some embodiments, R_2 is $-\text{CR}_{23}\text{R}_{24}\text{C}(\text{O})\text{R}_{22}$ and wherein R_{23} and R_{24} are each independently H or C_{1-8} alkyl; and R_{22} is phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylsulfonyl, cyano, C_{2-8} dialkylamino, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, halogen, heteroaryl, and phenyl.

R_2 is $-\text{CR}_{23}\text{R}_{24}\text{C}(\text{O})\text{NR}_{25}\text{R}_{22}$ and wherein R_{23} , R_{24} and R_{25} are each independently H or C_{1-8} alkyl; and R_{22} is C_{1-8} alkyl, C_{3-7} cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{2-6} alkenyl, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylsulfonyl, amino, carbo- C_{1-6} -alkoxy, carboxy, cyano, C_{3-7} cycloalkyl, C_{2-8} dialkylamino, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C_{1-7} alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C_{1-4} haloalkoxy, and heterocyclic.


In some embodiments, R_2 is $-\text{CH}_2\text{C}(\text{O})\text{NHR}_{22}$ and wherein R_{22} is phenyl optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} haloalkyl, and halogen.

In some embodiments, A and B are both $-\text{CH}_2\text{CH}_2-$, D is NR_2 , E is CR_4 , $---$ is a single bond and V_1 and V_2 are both single bonds; these embodiments can be represented by Formula (IIx) as shown below:




wherein each variable in Formula (IIx) has the same meaning as described herein, *supra* and *infra*. In some embodiments, compounds are of Formula (IIx) and W is NR_5 . In some embodiments, R_5 is H. In some embodiments, Z is cyano. In further embodiments, Q is NR_6 , O, S, $\text{S}(\text{O})$ or $\text{S}(\text{O})_2$. In still further embodiments, Q is NH or O.


In some embodiments, compounds of the present invention are of Formula (IIx) wherein R_2 is Formula (D); these embodiments can be represented by Formula (IIy) as shown below:



wherein each variable in Formula (IIy) has the same meaning as described herein, *supra* and *infra*. In some embodiments, G is $\text{C}(\text{O})$, $\text{C}(\text{O})\text{NR}_{23}$, $\text{C}(\text{O})\text{O}$, $\text{C}(\text{S})$, $\text{C}(\text{S})\text{NR}_{23}$, $\text{C}(\text{S})\text{O}$, $\text{CR}_{23}\text{R}_{24}$ or $\text{S}(\text{O})_2$. In some embodiments, G is $\text{C}(\text{O})$ and can be represented by Formula (IIz) as shown below:



wherein each variable in Formula (IIz) has the same meaning as described herein, *supra* and *infra*. In some embodiments, G is C(O)O and can be represented by Formula (IIIa) as shown below:



5 wherein each variable in Formula (IIIa) has the same meaning as described herein, *supra* and *infra*.

In some embodiments, compounds are of either Formula (IIz) or (IIIa) and R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro.

15 In some embodiments, compounds are of either Formula (IIz) or (IIIa) and R₂₂ is C₁₋₈ alkyl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro.

20 In some embodiments, compounds are of either Formula (IIz) or (IIIa) and R₂₂ is phenyl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄

haloalkyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro. In some embodiments, the phenyl is substituted with 1 to 4 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carboxamide, carboxy, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio and halogen. In some embodiments, the phenyl is substituted with 1 to 4 substituents selected from the group consisting of C₁₋₄ alkylsulfonyl, C₁₋₄ haloalkylsulfonyl and halogen.

In some embodiments, compounds are of either Formula (IIz) or (IIIa) and R₂₂ is heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro. In some embodiments, the heteroaryl is substituted with 1 to 4 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carboxamide, carboxy, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio and halogen. In some embodiments, the heteroaryl is substituted with 1 to 4 substituents selected from the group consisting of C₁₋₄ alkylsulfonyl, C₁₋₄ haloalkylsulfonyl and halogen. In some embodiments, the heteroaryl is a 5-membered heteroaryl, for example, as shown in TABLE 2A, *supra*. In some embodiments, the heteroaryl is a 6-membered heteroaryl, for example, as shown in TABLE 4, *supra*. In some embodiments, the heteroaryl is selected from the group consisting of pyridinyl, pyridazinyl, pyrimidinyl and pyrazinyl. In some embodiments, the heteroaryl is pyridinyl.

In some embodiments, R₂₂ is 1-methyl-1H-imidazole-4-yl, or 2,4-dimethyl-thiazole-5-yl.

In some embodiments, compounds are of Formula (IIy), (IIx) or (IIIa) and W is NR₅. In some embodiments, R₅ is H. In some embodiments, Z is cyano. In further embodiments, Q is NR₆, O, S, S(O) or S(O)₂. In still further embodiments, Q is NH or O.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein D is O, S, S(O) or S(O)₂.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein R₂₃ and R₂₄ are independently H or C₁₋₂ alkyl. In some embodiments, R₂₃ and R₂₄ are H.

In some embodiments, Z is selected from the group consisting of C₁₋₅ acyl, C₁₋₈ alkyl, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, amino, carbamimidoyl, cyano, C₃₋₇ cycloalkyl, heterocyclic, and hydroxycarbamimidoyl, wherein C₁₋₈ alkyl, C₃₋₇ cycloalkyl, and heterocyclic are each optionally substituted with 1, 2, 3 or 4 groups selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄

alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, amino, C₁₋₂ alkylamino, C₂₋₄ dialkylamino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, formyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro, and wherein said C₁₋₇ alkyl is
 5 optionally substituted with amino.


In some embodiments, Z is selected from the group consisting of C₁₋₅ acyl, C₁₋₈ alkyl, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, amino, carbamimidoyl, cyano, C₃₋₇ cycloalkyl, heterocyclic, and hydroxycarbamimidoyl, wherein said heterocyclic is optionally substituted with a -CH₂NH₂ group.

In some embodiments, Z is selected from the group consisting of C(O)CH₃, C(O)CH₂CH₃,
 10 CH₃, CH₂CH₃, C≡CH, NHS(O)₂CH₃, amino, carbamimidoyl, cyano, cyclopropyl, 4,5-dihydro-1H-imidazol-2-yl, 5-aminomethyl-4,5-dihydro-oxazol-2-yl, and hydroxycarbamimidoyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Z is selected from the group consisting of C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, carboxamide, carboxy, cyano, formyl, aryl,
 15 C₁₋₄ haloalkyl, C₁₋₄ haloalkylcarboxamide, heteroaryl, hydroxyl, hydroxylamino, nitro and tetrazolyl.

In some embodiments, Z is selected from the group consisting of formyl, NHC(O)CH₃, NHC(O)CH₂CH₃, NHC(O)CH(CH₃)₂, CH₃, CH₂CH₃, CH(CH₃)₂, CH₂CH₂CH₂CH₃, NHC(O)CF₃, carboxy, cyano, CF₃, CF₂CF₃, nitro and 1H-tetrazol-5-yl. In some embodiments, Z is selected from the group consisting of carboxy, CF₃, nitro and 1H-tetrazol-5-yl. In some embodiments, Z is cyano.
 20 In still further embodiments, Z is formyl [i.e. -C(O)H].

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Z is Formula (A):



wherein:

25 R₇ is H, C₁₋₈ alkyl or C₃₋₇ cycloalkyl; and R₈ is H, nitro or nitrile. In some embodiments, R₇ is H or C₁₋₈ alkyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein R₁, R₇ and R₈ are independently selected from the group consisting of H, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₂₋₆ alkynyl, amino, C₃₋₇ cycloalkyl and C₁₋₄ haloalkyl. In some embodiments, R₁, R₇ and R₈ are
 30 independently H, halogen or amino. In still further embodiments, R₁, R₇ and R₈ are H.

In some embodiments, Ar₁ is aryl or heteroaryl each optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅; wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen,

heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonamide, alkylsulfonamide, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonooxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

In some embodiments, Ar₁ is aryl.

In some embodiments, Ar₁ is phenyl optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅; wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, carboxy, C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonooxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

In some embodiments, Ar₁ is phenyl optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅; wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylthio, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are

each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonooxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with
 5 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

In some embodiments, Ar₁ is phenyl optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅;
 10 wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄
 15 alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonooxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with
 20 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₈ alkyl, and halogen.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Ar₁ is phenyl. In some embodiments, the phenyl is optionally substituted with R₁₁. In some
 embodiments, R₁₁ is selected from the group consisting of H, C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄
 25 alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carboxamide, C₃₋₇ cycloalkyl, halogen and sulfonamide. In some embodiments, R₁₁ is selected from the group consisting of C(O)CH₃, C(O)CH₂CH₃, C(O)CH₂CH₂CH₃, C(O)CH(CH₃)₂, C(O)CH₂CH₂CH₂CH₃, OCH₃, OCH₂CH₃, OCH₂CH₂CH₃, OCH(CH₃)₂, OCH₂CH₂CH₂CH₃, CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, CH(CH₃)(CH₂CH₃), CH₂(CH₂)₂CH₃, CH₂(CH₂)₃CH₃,
 30 CH₂(CH₂)₄CH₃, CH₂(CH₂)₅CH₃, C(O)NHCH₃, C(O)NHCH₂CH₃, C(O)NHCH₂CH₂CH₃, C(O)NHCH(CH₃)₂, C(O)NHCH₂(CH₂)₂CH₃, CCH, S(O)₂NHCH₃, S(O)₂NHCH₂CH₃, S(O)₂NHCH₂CH₂CH₃, S(O)₂NHCH(CH₃)₂, S(O)₂NHCH₂(CH₂)₂CH₃, S(O)₂NHCH(CH₃)CH₂CH₃, S(O)CH₃, S(O)CH₂CH₃, S(O)CH₂CH₂CH₃, S(O)CH(CH₃)₂, S(O)CH₂(CH₂)₂CH₃, S(O)CH(CH₃)CH₂CH₃, S(O)₂CH₃, S(O)₂CH₂CH₃, S(O)₂CH₂CH₂CH₃, S(O)₂CH(CH₃)₂,
 35 S(O)₂CH₂(CH₂)₂CH₃, S(O)₂CH(CH₃)CH₂CH₃, SCH₃, SCH₂CH₃, SCH₂CH₂CH₃, SCH(CH₃)₂ and SCH₂(CH₂)₂CH₃. In some embodiments, R₁₁ is selected from the group consisting of amino, arylsulfonyl, carboxy, cyano, C₃₋₇ cycloalkyl, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl and C₁₋₄

haloalkylthio. In some embodiments, R_{11} is selected from the group consisting of phenylsulfonyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Cl, F, Br, OCF_3 , $OCHF_2$, OCH_2CF_3 , CF_3 , CHF_2 , CH_2CF_3 , SCF_3 , $SCHF_2$ and SCH_2CF_3 . In some embodiments, R_{11} is selected from the group consisting of heterocyclic, heteroaryl, C_{4-7} oxo-cycloalkyl, phenoxy and phenyl. In some

5 embodiments, R_{11} is selected from the group consisting of morpholin-4-yl, thiomorpholin-4-yl, 1-oxo-1 λ^4 -thiomorpholin-4-yl, 1,1-Dioxo-1 λ^6 -thiomorpholin-4-yl, piperazin-1-yl, 4-methyl-piperazin-1-yl, 4-ethyl-piperazin-1-yl, 4-propyl-piperazin-1-yl, piperidin-1-yl, pyrrolidin-1-yl, 2,5-dioxo-imidazolidin-4-yl, 2,4-dioxo-thiazolidin-5-yl, 4-oxo-2-thioxo-thiazolidin-5-yl, 3-methyl-2,5-dioxo-imidazolidin-4-yl, 3-methyl-2,4-dioxo-thiazolidin-5-yl, 3-methyl-4-oxo-2-thioxo-thiazolidin-5-yl, 3-

10 ethyl-2,5-dioxo-imidazolidin-4-yl, 3-ethyl-2,4-dioxo-thiazolidin-5-yl, and 3-ethyl-4-oxo-2-thioxo-thiazolidin-5-yl. In some embodiments, R_{11} is selected from the group consisting of 1H-imidazol-4-yl, [1,2,4]triazol-1-yl, [1,2,3]triazol-1-yl, [1,2,4]triazol-4-yl, pyrrol-1-yl, pyrazol-1-yl, 1H-pyrazol-3-yl, imidazol-1-yl, oxazol-5-yl, oxazol-2-yl, [1,3,4]oxadiazol-2-yl, [1,3,4]thiadiazol-2-yl, [1,2,4]oxadiazol-3-yl, [1,2,4]thiadiazol-3-yl, tetrazol-1-yl, pyrimidin-5-yl, pyrimidin-2-yl, pyrimidin-

15 4-yl, pyridazin-3-yl, pyridazin-4-yl, pyrazin-2-yl, 1,3-dioxo-1,3-dihydro-isoindol-2-yl and [1,2,3]thiadiazol-4-yl. In some embodiments, R_{11} is C_{1-8} alkyl or C_{1-4} alkoxy optionally substituted with 1 to 5 substituents selected independently from the group consisting of C_{1-4} alkoxy, C_{1-4} alkylcarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, carbo-

20 C_{1-6} -alkoxy, carboxamide, carboxy, cyano, heterocyclic, hydroxyl and phenyl. In some embodiments, R_{11} is C_{1-4} alkylsulfonyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C_{1-4} alkoxy, carboxamide, heteroaryl, heterocyclic and phenyl. In some


embodiments, the C_{1-4} alkylsulfonyl is substituted with the heteroaryl group. In some embodiments, the heteroaryl group is selected from the group consisting of 1H-imidazol-4-yl, [1,2,4]triazol-1-yl, [1,2,3]triazol-1-yl, [1,2,4]triazol-4-yl, pyrrol-1-yl, pyrazol-1-yl, 1H-pyrazol-3-yl, imidazol-1-yl,

25 oxazol-5-yl, oxazol-2-yl, [1,3,4]oxadiazol-2-yl, [1,3,4]thiadiazol-2-yl, [1,2,4]oxadiazol-3-yl, [1,2,4]thiadiazol-3-yl, tetrazol-1-yl, pyrimidin-5-yl, pyrimidin-2-yl, pyrimidin-4-yl, pyridazin-3-yl, pyridazin-4-yl, pyrazin-2-yl, 1,3-dioxo-1,3-dihydro-isoindol-2-yl and [1,2,3]thiadiazol-4-yl. In some

embodiments, R_{11} is arylsulfonyl, heteroaryl, phenoxy or phenyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C_{1-5} acyl, C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, carboxamide, carboxy, cyano,

30 halogen, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkylthio and hydroxyl. In some embodiments, R_{11} is arylsulfonyl, heteroaryl, phenoxy or phenyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C_{1-4} alkoxy, C_{1-8} alkyl, cyano, halogen, C_{1-4} haloalkoxy, C_{1-4} haloalkyl and hydroxyl.

35 Some embodiments of the present invention pertain to compounds of Formula (I) wherein Ar_1 is phenyl. In some embodiments, the phenyl is optionally substituted with R_{11} . In some embodiments, R_{11} is a group of Formula (B):




wherein:

“p” and “r” are independently 0, 1, 2 or 3; and R₁₆ is H, C₁₋₅ acyl, C₂₋₆ alkenyl, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, halogen, heteroaryl or phenyl, and wherein the heteroaryl or phenyl may be optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₄ alkoxy, amino, C₁₋₄ alkylamino, C₂₋₆ alkynyl, C₂₋₈ dialkylamino, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl and hydroxyl. In some embodiments, p = 0 and r = 0. In some embodiments, R₁₆ is heteroaryl or phenyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₄ alkoxy, amino, C₁₋₄ alkylamino, C₂₋₆ alkynyl, C₂₋₈ dialkylamino, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl and hydroxyl. In some embodiments, the heteroaryl is selected from the group consisting of 1H-imidazol-4-yl, [1,2,4]triazol-1-yl, [1,2,3]triazol-1-yl, [1,2,4]triazol-4-yl, pyrrol-1-yl, pyrazol-1-yl, 1H-pyrazol-3-yl, imidazol-1-yl, oxazol-5-yl, oxazol-2-yl, [1,3,4]oxadiazol-2-yl, [1,3,4]thiadiazol-2-yl, [1,2,4]oxadiazol-3-yl, [1,2,4]thiadiazol-3-yl, tetrazol-1-yl, pyrimidin-5-yl, pyrimidin-2-yl, pyrimidin-4-yl, pyridazin-3-yl, pyridazin-4-yl, pyrazin-2-yl, 1,3-dioxo-1,3-dihydro-isindol-2-yl and [1,2,3]thiadiazol-4-yl. In some embodiments, p = 0 and r = 1. In some embodiments, R₁₆ is carbo-C₁₋₆-alkoxy or carboxy. In some embodiments, p = 2 and r = 1. In some embodiments, R₁₆ is H, C₁₋₅ acyl or C₁₋₈ alkyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Ar₁ is phenyl. In some embodiments, the phenyl is optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅. In some embodiments, R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅ are independently selected from the group consisting of H, C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylureyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, halogen, C₁₋₄ haloalkoxy and C₁₋₄ haloalkyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Ar₁ is phenyl and R₁₁ is substituted at the para position on the phenyl; these embodiments can be represented by Formula (IIIc) as shown below:




wherein each variable in Formula (IIIc) has the same meaning as described herein, *supra* and *infra*.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Ar₁ is phenyl and two adjacent R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅ groups together with the atoms to which they are attached form a 5-, 6- or 7-membered cycloalkyl, cycloalkenyl or heterocyclic group fused with Ar₁, wherein the 5-, 6- or 7-membered group is optionally substituted with halogen. In some

embodiments, the phenyl and two adjacent R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅ groups form a 5-, 6- or 7-membered cycloalkyl as represented in TABLE 5:

TABLE 5



wherein "a" is 1, 2 or 3 to give a 5-, 6- or 7-membered cycloalkyl fused together with the phenyl group where two of the ring carbons are shared between the cycloalkyl and phenyl group. In some embodiments, 1, 2, or 3 ring carbons are replaced by a heteroatom selected from, but not limited to, O, S, and N, wherein N is substituted with H or C₁₋₄ alkyl. In some embodiments, the two adjacent groups form a 5 membered heterocyclic group with the phenyl group. In some embodiments, the 5 membered heterocyclic group with the phenyl group together is a 2,3-dihydro-benzofuran-5-yl or benzo[1,3]dioxol-5-yl group. In some embodiments, the two adjacent groups form a 6 membered heterocyclic group with the phenyl group. In some embodiments, the 6 membered heterocyclic group with the phenyl group together is a 2,3-dihydro-benzo[1,4]dioxin-6-yl or 2,3-dihydro-benzo[1,4]dioxin-2-yl group. In some embodiments, the two adjacent groups form a 7 membered heterocyclic group with the phenyl group. In some embodiments, the 7 membered heterocyclic group with the phenyl group together is a 3,4-dihydro-2H-benzo[b][1,4]dioxepin-7-yl group.

In some embodiments, Ar₁ is heteroaryl.

In some embodiments, Ar₁ is pyridyl optionally substituted with R₁₁, R₁₂, R₁₃, and R₁₄; wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, and sulfonamide, and wherein C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonamide, alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonooxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and R₁₂, R₁₃, and R₁₄ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄

alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

In some embodiments, Ar₁ is pyridyl optionally substituted with R₁₁, R₁₂, R₁₃, and R₁₄; wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, C₂₋₆ dialkylamino, halogen, heterocyclic, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, alkylsulfonyl, C₁₋₄ alkylthio, C₂₋₆ dialkylamino, and heterocyclic are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₄ alkylsulfonyl, C₃₋₇ cycloalkyloxy, heteroaryl, hydroxyl, phenyl, and phosphonooxy; and R₁₂, R₁₃, and R₁₄ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

In some embodiments, Ar₁ is pyridyl optionally substituted with R₁₁, R₁₂, R₁₃, and R₁₄; wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, C₂₋₆ dialkylamino, halogen, heterocyclic, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, alkylsulfonyl, C₁₋₄ alkylthio, C₂₋₆ dialkylamino, and heterocyclic are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₄ alkylsulfonyl, C₃₋₇ cycloalkyloxy, heteroaryl, hydroxyl, phenyl, and phosphonooxy; and R₁₂, R₁₃, and R₁₄ are each independently selected from the group consisting of C₁₋₈ alkyl, and halogen.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Ar₁ is heteroaryl. In some embodiments, the heteroaryl is optionally substituted with R₁₁. In some embodiments, R₁₁ is selected from the group consisting of C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carboxamide, C₃₋₇ cycloalkyl, halogen and sulfonamide. In some embodiments, R₁₁ is selected from the group consisting of C(O)CH₃, C(O)CH₂CH₃, C(O)CH₂CH₂CH₃, C(O)CH(CH₃)₂, C(O)CH₂CH₂CH₂CH₃, OCH₃, OCH₂CH₃, OCH₂CH₂CH₃, OCH(CH₃)₂, OCH₂CH₂CH₂CH₃, CH₃, CH₂CH₃, CH₂CH₂CH₃, CH(CH₃)₂, CH(CH₃)(CH₂CH₃), CH₂(CH₂)₂CH₃, CH₂(CH₂)₃CH₃, CH₂(CH₂)₄CH₃, CH₂(CH₂)₅CH₃, C(O)NHCH₃, C(O)NHCH₂CH₃, C(O)NHCH₂CH₂CH₃, C(O)NHCH(CH₃)₂, C(O)NHCH₂(CH₂)₂CH₃, CCH, S(O)₂NHCH₃, S(O)₂NHCH₂CH₃, S(O)₂NHCH₂CH₂CH₃, S(O)₂NHCH(CH₃)₂, S(O)₂NHCH₂(CH₂)₂CH₃, S(O)₂NHCH(CH₃)CH₂CH₃, S(O)CH₃, S(O)CH₂CH₃, S(O)CH₂CH₂CH₃, S(O)CH(CH₃)₂, S(O)CH₂(CH₂)₂CH₃, S(O)CH(CH₃)CH₂CH₃, S(O)₂CH₃, S(O)₂CH₂CH₃, S(O)₂CH₂CH₂CH₃, S(O)₂CH(CH₃)₂, S(O)₂CH₂(CH₂)₂CH₃, S(O)₂CH(CH₃)CH₂CH₃, SCH₃, SCH₂CH₃, SCH₂CH₂CH₃, SCH(CH₃)₂ and SCH₂(CH₂)₂CH₃. In some embodiments, R₁₁ is selected from the group consisting of amino, arylsulfonyl, carboxy, cyano, C₃₋₇ cycloalkyl, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl and C₁₋₄

haloalkylthio. In some embodiments, R_{11} is selected from the group consisting of phenylsulfonyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Cl, F, Br, OCF_3 , $OCHF_2$, OCH_2CF_3 , CF_3 , CHF_2 , CH_2CF_3 , SCF_3 , $SCHF_2$ and SCH_2CF_3 . In some embodiments, R_{11} is selected from the group consisting of heterocyclic, heteroaryl, C_{4-7} oxo-cycloalkyl, phenoxy and phenyl. In some

5 embodiments, R_{11} is selected from the group consisting of morpholin-4-yl, thiomorpholin-4-yl, 1-oxo-1 λ^4 -thiomorpholin-4-yl, 1,1-Dioxo-1 λ^6 -thiomorpholin-4-yl, piperazin-1-yl, 4-methyl-piperazin-1-yl, 4-ethyl-piperazin-1-yl, 4-propyl-piperazin-1-yl, piperidin-1-yl, pyrrolidin-1-yl, 2,5-dioxo-imidazolidin-4-yl, 2,4-dioxo-thiazolidin-5-yl, 4-oxo-2-thioxo-thiazolidin-5-yl, 3-methyl-2,5-dioxo-imidazolidin-4-yl, 3-methyl-2,4-dioxo-thiazolidin-5-yl, 3-methyl-4-oxo-2-thioxo-thiazolidin-5-yl, 3-ethyl-2,5-dioxo-imidazolidin-4-yl, 3-ethyl-2,4-dioxo-thiazolidin-5-yl, and 3-ethyl-4-oxo-2-thioxo-thiazolidin-5-yl. In some embodiments, R_{11} is selected from the group consisting of 1H-imidazol-4-yl, [1,2,4]triazol-1-yl, [1,2,3]triazol-1-yl, [1,2,4]triazol-4-yl, pyrrol-1-yl, pyrazol-1-yl, 1H-pyrazol-3-yl, imidazol-1-yl, oxazol-5-yl, oxazol-2-yl, [1,3,4]oxadiazol-2-yl, [1,3,4]thiadiazol-2-yl, [1,2,4]oxadiazol-3-yl, [1,2,4]thiadiazol-3-yl, tetrazol-1-yl, pyrimidin-5-yl, pyrimidin-2-yl, pyrimidin-4-yl, pyridazin-3-yl, pyridazin-4-yl, pyrazin-2-yl, 1,3-dioxo-1,3-dihydro-isoindol-2-yl and [1,2,3]thiadiazol-4-yl. In some embodiments, R_{11} is C_{1-8} alkyl or C_{1-4} alkoxy, optionally substituted with 1 to 5 substituents selected independently from the group consisting of C_{1-4} alkoxy, C_{1-4} alkylcarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, carbo- C_{1-6} -alkoxy, carboxamide, carboxy, cyano, heterocyclic, hydroxyl and phenyl. In some


15 R_{11} is C_{1-4} alkylsulfonyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C_{1-4} alkoxy, carboxamide, heteroaryl, heterocyclic and phenyl. In some embodiments, the C_{1-4} alkylsulfonyl is substituted with the heteroaryl group. In some embodiments, the heteroaryl is selected from the group consisting of 1H-imidazol-4-yl, [1,2,4]triazol-1-yl, [1,2,3]triazol-1-yl, [1,2,4]triazol-4-yl, pyrrol-1-yl, pyrazol-1-yl, 1H-pyrazol-3-yl, imidazol-1-yl, oxazol-5-yl, oxazol-2-yl, [1,3,4]oxadiazol-2-yl, [1,3,4]thiadiazol-2-yl, [1,2,4]oxadiazol-3-yl, [1,2,4]thiadiazol-3-yl, tetrazol-1-yl, pyrimidin-5-yl, pyrimidin-2-yl, pyrimidin-4-yl, pyridazin-3-yl, pyridazin-4-yl, pyrazin-2-yl, 1,3-dioxo-1,3-dihydro-isoindol-2-yl and [1,2,3]thiadiazol-4-yl. In some

20 R_{11} is arylsulfonyl, heteroaryl, phenoxy or phenyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C_{1-5} acyl, C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, carboxamide, carboxy, cyano, halogen, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkylthio and hydroxyl. In some embodiments, R_{11} is arylsulfonyl, heteroaryl, phenoxy or phenyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C_{1-4} alkoxy, C_{1-8} alkyl, cyano, halogen, C_{1-4} haloalkoxy, C_{1-4} haloalkyl and hydroxyl.

25

30

35 Some embodiments of the present invention pertain to compounds of Formula (I) wherein Ar_1 is heteroaryl. In some embodiments, the heteroaryl is optionally substituted with R_{11} . In some embodiments, R_{11} is of Formula (B):



wherein:

“p” and “r” are independently 0, 1, 2 or 3; and R₁₆ is H, C₁₋₅ acyl, C₂₋₆ alkenyl, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, halogen, heteroaryl or phenyl, and wherein the heteroaryl or phenyl may be optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₄ alkoxy, amino, C₁₋₄ alkylamino, C₂₋₆ alkynyl, C₂₋₈ dialkylamino, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl and hydroxyl. In some embodiments, p = 0 and r = 0. In some embodiments, R₁₆ is heteroaryl or phenyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₄ alkoxy, amino, C₁₋₄ alkylamino, C₂₋₆ alkynyl, C₂₋₈ dialkylamino, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl and hydroxyl. In some embodiments, the heteroaryl is selected from the group consisting of 1H-imidazol-4-yl, [1,2,4]triazol-1-yl, [1,2,3]triazol-1-yl, [1,2,4]triazol-4-yl, pyrrol-1-yl, pyrazol-1-yl, 1H-pyrazol-3-yl, imidazol-1-yl, oxazol-5-yl, oxazol-2-yl, [1,3,4]oxadiazol-2-yl, [1,3,4]thiadiazol-2-yl, [1,2,4]oxadiazol-3-yl, [1,2,4]thiadiazol-3-yl, tetrazol-1-yl, pyrimidin-5-yl, pyrimidin-2-yl, pyrimidin-4-yl, pyridazin-3-yl, pyridazin-4-yl, pyrazin-2-yl, 1,3-dioxo-1,3-dihydro-isindol-2-yl and [1,2,3]thiadiazol-4-yl. In some embodiments, p = 0 and r = 1. In some embodiments, R₁₆ is carbo-C₁₋₆-alkoxy or carboxy. In some embodiments, p = 2 and r = 1. In some embodiments, R₁₆ is H, C₁₋₅ acyl or C₁₋₈ alkyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Ar₁ is heteroaryl. In some embodiments, the heteroaryl is optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅. In some embodiments, R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅ are independently selected from the group consisting of H, C₁₋₅ acyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylureyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, halogen, C₁₋₄ haloalkoxy and C₁₋₄ haloalkyl.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Ar₁ is heteroaryl. In some embodiments, the heteroaryl is optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅ wherein two adjacent R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅ groups together with the atoms to which they are attached form a 5-, 6- or 7-membered cycloalkyl, cycloalkenyl or heterocyclic group fused with Ar₁, wherein the 5-, 6- or 7-membered group is optionally substituted with halogen. In some embodiments, the two adjacent groups form a 5-membered heterocyclic group with the heteroaryl group. In some embodiments, the two adjacent groups form a 6-membered heterocyclic group with the heteroaryl group. In some embodiments, the two adjacent groups form a 7-membered heterocyclic group with the heteroaryl group.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein R₄, R₅ and R₆ are independently H or CH₃.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein X is N.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein Y is N.

5 Some embodiments of the present invention pertain to compounds of Formula (I) wherein X is N and Y is CH.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein X is CH and Y is N.

10 Some embodiments of the present invention pertain to compounds of Formula (I) wherein X and Y are N.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein X and Y are CH.

15 Some embodiments of the present invention pertain to compounds of Formula (I) wherein:

A and B are each independently -CH₂CH₂- or -CH₂-;

D is N-R₂;

V₁ is a bond;

V₂ is -CH₂-, -CH₂CH₂-, or a bond;

W and Q are each independently NH or O;

20 X and Y are each independently N or CH, provided that if either X or Y is CH then the other is N;

Z is selected from the group consisting of nitro, C₁₋₅ acyl, C₁₋₈ alkyl, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, amino, carbamimidoyl, cyano, C₃₋₇ cycloalkyl, heterocyclic, and hydroxycarbamimidoyl, wherein said heterocyclic is optionally substituted with a -CH₂NH₂ group;

25 R₂ is -C(O)OR₂₂, -C(O)R₂₂, -CH₂R₂₂, -R₂₂, -S(O)₂R₂₂, -CR₂₃R₂₄C(O)R₂₂, or -CR₂₃R₂₄C(O)NR₂₅R₂₂, wherein R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, 30 hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic; and R₂₃ and R₂₄ are each independently H or C₁₋₈ alkyl;

Ar₁ is aryl or heteroaryl each optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅; wherein R₁₁ is selected from the group consisting of C₁₋₆ alkylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₂₋₆ alkynyl, 35 C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, and sulfonamide, and wherein C₁₋₆

acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonamide, alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonoxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein:

A and B are both -CH₂CH₂-;

D is N-R₂;

V₁ and V₂ are both a bond;

W and Q are each independently NH or O;

X and Y are both N;

Z is selected from the group consisting of nitro, C(O)CH₃, C(O)CH₂CH₃, CH₃, CH₂CH₃, C≡CH, NHS(O)₂CH₃, amino, carbamimidoyl, cyano, cyclopropyl, 4,5-dihydro-1H-imidazol-2-yl, 5-aminomethyl-4,5-dihydro-oxazol-2-yl, and hydroxycarbamimidoyl;

R₂ is -C(O)OR₂₂, wherein R₂₂ is C₁₋₈ alkyl, or C₃₋₇ cycloalkyl each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylsulfonyl, amino, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, and hydroxyl;

Ar₁ is phenyl optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅;

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonoxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R_{12} , R_{13} , R_{14} , and R_{15} are each independently selected from the group consisting of C_{1-8} alkyl, and halogen.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein:

A and B are both $-CH_2CH_2-$;

5 D is $N-R_2$;

V_1 and V_2 are both a bond;

W is NH;

Q is O;

X and Y are both N;

10 Z is nitro, cyano, $C(O)CH_3$, amino, CH_3 , CH_2CH_3 , or $C\equiv CH$;

R_2 is $-C(O)OR_{22}$, $-C(O)R_{22}$, $-R_{22}$, or $-S(O)_2R_{22}$ wherein R_{22} is C_{1-8} alkyl, C_{3-7} cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-5} acyl, C_{2-6} alkenyl, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylsulfonyl, amino, carbo-
 15 C_{1-6} -alkoxy, carboxy, cyano, C_{3-7} cycloalkyl, C_{2-8} dialkylamino, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C_{1-7} alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C_{1-4} haloalkoxy, and heterocyclic;

Ar_1 is phenyl, 3-pyridyl, or 2-pyridyl each optionally substituted with R_{11} , R_{12} , R_{13} , R_{14} , and R_{15} ,

20 wherein R_{11} is selected from the group consisting of C_{1-6} acylsulfonamide, C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylamino, C_{1-6} alkylcarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, carbamimidoyl, carboxamide, carboxy, cyano, C_{2-6} dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylamino, C_{1-6} alkylcarboxamide, C_{1-4}
 25 alkylsulfonamide, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, carbamimidoyl, C_{2-6} dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C_{1-6} acylsulfonamide, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylcarboxamide, C_{1-4} alkylsulfonyl, carboxy, C_{2-6} dialkylamino, C_{2-6} dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonooxy wherein said C_{1-7} alkyl and C_{1-4} alkylcarboxamide
 30 are each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-4} alkoxy and hydroxy; and

R_{12} , R_{13} , R_{14} , and R_{15} are each independently CH_3 , or F.

Some embodiments of the present invention pertain to compounds of Formula (I) wherein:

A and B are both $-CH_2CH_2-$;

35 D is $N-R_2$;

V_1 and V_2 are both a bond;

W and Q are both O;

X and Y are both N;

Z is selected from the group consisting of CH₃, CH₂CH₃, cyclopropyl, or C≡CH;

R₂ is -C(O)OR₂₂, -C(O)R₂₂, -R₂₂, -CH₂C(O)R₂₂, or -CH₂C(O)NHR₂₂, wherein R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carboxy, cyano, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, hydroxyl, phenyl, and phenoxy, wherein said C₁₋₇ alkyl, is optionally substituted with 1 or 2 substituents selected from the group consisting of C₁₋₄ haloalkoxy, and heterocyclic;

Ar₁ is phenyl, 2-pyridyl, or 3-pyridyl each optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅,

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₂₋₆ alkynyl C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₂₋₆ dialkylamino, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, heteroaryl, hydroxyl, and phosphonooxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₈ alkyl, and halogen.

In some embodiments, compounds of the present invention are when R₁₁ is selected from the group consisting of:

sulfamoyl [-S(O)₂NH₂],
 25 acetylsulfamoyl [-S(O)₂NHC(O)CH₃],
 propionylsulfamoyl [-S(O)₂NHC(O)CH₂CH₃],
 butyrylsulfamoyl [-S(O)₂NHC(O)CH₂CH₂CH₃],
 pentanoylsulfamoyl [-S(O)₂NHC(O)CH₂CH₂CH₂CH₃],
 methanesulfonyl [-S(O)₂CH₃],
 30 ethanesulfonyl [-S(O)₂CH₂CH₃],
 propane-1-sulfonyl [-S(O)₂CH₂CH₂CH₃],
 hydroxymethyl (-CH₂OH);
 2-hydroxyethyl (-CH₂CH₂OH);
 3-hydroxypropyl (-CH₂CH₂CH₂OH);
 35 4-hydroxy-butyl (-CH₂CH₂CH₂CH₂OH);
 phosphonooxymethyl [-CH₂OP(O)(OH)₂];
 2-phosphonooxy-ethyl [-CH₂CH₂OP(O)(OH)₂];

3-phosphonooxy-propyl $[-\text{CH}_2\text{CH}_2\text{CH}_2\text{OP}(\text{O})(\text{OH})_2]$; and

4-phosphonooxy-butyl $[-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OP}(\text{O})(\text{OH})_2]$.

In some embodiments, R_{11} is methoxy, ethoxy, isobutoxy or 3-methyl-butoxy.

In some embodiments, R_{11} is pyridyl optionally substituted with C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylamino, halogen or hydroxyl.

In some embodiments, R_{11} is 2-pyridyl optionally substituted with C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylamino, halogen or hydroxyl.

In some embodiments, R_{11} is 3-pyridyl optionally substituted with C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylamino, halogen or hydroxyl.

Some embodiments of the present invention include one or more of the compounds illustrated in TABLES A, B, C, D and E; these TABLES are shown below.

TABLE A

Cmpd#	Structure	Chemical Name
A1		4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester
A2		(4-Methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine
A3		1-{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-3,3-dimethyl-butan-1-one
A4		(4-Methanesulfonyl-phenyl)-[5-nitro-6-(1-thiophen-3-ylmethyl-piperidin-4-yloxy)-pyrimidin-4-yl]-thiophen-3-ylmethyl-amine
A5		(4-Methanesulfonyl-phenyl)-[5-nitro-6-(1-pyridin-2-ylmethyl-piperidin-4-yloxy)-pyrimidin-4-yl]-amine

Cmpd#	Structure	Chemical Name
A6		(4-Methanesulfonyl-phenyl)-[5-nitro-6-(1-pyridin-3-ylmethyl)-piperidin-4-yloxy]-pyrimidin-4-yl]-amine
A7		{6-[1-(3,3-Dimethyl-butyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine
A8		(4-Methanesulfonyl-phenyl)-{6-[1-(3-methyl-butyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl]-amine
A9		(4-Methanesulfonyl-phenyl)-[5-nitro-6-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yloxy)-pyrimidin-4-yl]-amine
A10		4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid ethyl ester
A11		1-{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-3,3-dimethyl-butan-2-one
A12		{6-[1-(2-Ethoxy-ethyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine
A13		4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxymethyl]-piperidine-1-carboxylic acid tert-butyl ester

Cmpd#	Structure	Chemical Name
A14		4-{2-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-ethyl}-piperidine-1-carboxylic acid tert-butyl ester
A15		3-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester
A16		3-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxymethyl]-pyrrolidine-1-carboxylic acid tert-butyl ester
A17		3-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxymethyl]-pyrrolidine-1-carboxylic acid tert-butyl ester
A18		4-[5-Cyano-6-(6-methylsulfonyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester
A19		4-[5-Cyano-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester
A20		[6-(1-Hexyl-piperidin-4-yloxy)-5-nitro-pyrimidin-4-yl]-(4-methanesulfonyl-phenyl)-amine
A21		[6-(1-Cyclopropylmethyl-piperidin-4-yloxy)-5-nitro-pyrimidin-4-yl]-(4-methanesulfonyl-phenyl)-amine
A22		4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
A23		4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid 2-isopropyl-5-methyl-cyclohexyl ester
A24		{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-pyridin-3-yl-methanone
A25		(2-Chloro-pyridin-3-yl)-{4-[6-(4-methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-methanone
A26		{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-pyridin-2-yl-methanone
A27		(4-Methanesulfonyl-phenyl)-[6-(1-methanesulfonyl-piperidin-4-yloxy)-5-nitro-pyrimidin-4-yl]-amine
A28		(4-Methanesulfonyl-phenyl)-{5-nitro-6-[1-(propane-1-sulfonyl)-piperidin-4-yloxy]-pyrimidin-4-yl}-amine
A29		{6-[1-(Butane-1-sulfonyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine
A30		(4-Methanesulfonyl-phenyl)-{5-nitro-6-[1-(thiophene-2-sulfonyl)-piperidin-4-yloxy]-pyrimidin-4-yl}-amine
A31		(4-Methanesulfonyl-phenyl)-{6-[1-(1-methyl-1H-imidazole-4-sulfonyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-amine









Cmpd#	Structure	Chemical Name
A32		{6-[1-(2,4-Dimethyl-thiazole-5-sulfonyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine
A33		4-[5-Cyano-6-(3-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester
A34		4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester
A35		4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester
A36		4-[6-(6-Methanesulfonyl-pyridin-3-ylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester
A37		4-[5-Acetyl-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester
A38		4-[5-Amino-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester
A39		4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A40		4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid ethyl ester

Cmpd#	Structure	Chemical Name
A41		4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isobutyl ester
A42		4-(4-Methanesulfonyl-phenylamino)-6-[1-(tetrahydrofuran-2-carbonyl)-piperidin-4-yloxy]-pyrimidine-5-carbonitrile
A43		4-[1-(3,3-Dimethyl-2-oxo-butyl)-piperidin-4-yloxy]-6-(4-methanesulfonyl-phenylamino)-pyrimidine-5-carbonitrile
A44		4-(4-Methanesulfonyl-phenylamino)-6-[1-(pyridine-3-carbonyl)-piperidin-4-yloxy]-pyrimidine-5-carbonitrile
A45		4-(1-Formyl-piperidin-4-yloxy)-6-(4-methanesulfonyl-phenylamino)-pyrimidine-5-carbonitrile
A46		4-(4-Methanesulfonyl-phenylamino)-6-[1-(pyridine-2-carbonyl)-piperidin-4-yloxy]-pyrimidine-5-carbonitrile
A47		4-[6-(4-Cyano-2-fluoro-phenylamino)-5-ethynyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A48		4-[5-Ethynyl-6-(2-fluoro-4-[1,2,4]triazol-1-yl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
A49		4-{5-Ethynyl-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-pyrimidin-4-ylamino}-3-fluoro-benzonitrile
A50		{5-Ethynyl-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-pyrimidin-4-yl}-(2-fluoro-4-methanesulfonyl-phenyl)-amine
A51		4-{6-[2,5-Difluoro-4-(2-methanesulfonyl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A52		4-{6-[2-Fluoro-4-(2-sulfamoyl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A53		4-{6-[6-(2-Fluoro-ethyl)-2-methyl-pyridin-3-ylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A54		4-{2-[4-Fluoro-6-(2-isopropoxy-ethyl)-pyridin-3-ylamino]-3-methyl-pyridin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A55		4-{6-[2,5-Difluoro-4-(2-[1,2,4]triazol-1-yl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A56		4-{5-Ethynyl-6-[2-fluoro-4-(4-methoxy-pyridin-2-yl)-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester









Cmpd#	Structure	Chemical Name
A57		4-{6-[2-Fluoro-4-(2-propionylsulfamoyl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A58		4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A59		4-{6-[2,3-Difluoro-4-(2-methanesulfonyl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A60		4-[5-Acetyl-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isobutyl ester
A61		1-[4-(1-Benzyl-azetidin-3-yloxy)-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-5-yl]-ethanone
A62		4-[5-Cyano-6-(6-propylamino-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A63		4-[5-Cyano-6-(2-fluoro-4-isopropylamino-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A64		4-[5-Cyano-6-(2-fluoro-4-propylamino-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
A73		4-[5-Cyano-6-(4-dimethylamino-2-fluoro-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A74		4-{5-Cyano-6-[2-fluoro-4-(2-pyrrolidin-1-yl-ethylamino)-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A75		4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A76		4-{5-Cyano-6-[2-fluoro-4-(2-morpholin-4-yl-ethylamino)-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A77		4-[6-(2-Fluoro-4-iodo-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A78		4-[5-Cyano-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A79		4-[6-(2-Fluoro-4-morpholin-4-yl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A80		4-[6-(2,5-Difluoro-4-propoxy-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A81		4-[6-(2-Fluoro-4-propylamino-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
A82		4-{6-[2-Fluoro-4-(2-methoxyethylamino)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A83		4-(6-{2-Fluoro-4-[(tetrahydrofuran-2-ylmethyl)-amino]-phenylamino}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
A84		4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethylamino)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A85		4-(6-{2-Fluoro-4-[(2-methanesulfonyl-ethyl)-methyl-amino]-phenylamino}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
A86		4-[6-(4-Bromo-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A87		4-[6-(4-Cyano-2-fluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A88		4-[6-(4-Cyano-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A89		4-[6-(2,5-Difluoro-4-morpholin-4-yl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
A90		4-[6-(6-Chloro-2-methyl-pyridin-3-ylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A91		4-[5-Methyl-6-(2-methyl-6-morpholin-4-yl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A92		4-[5-(4,5-Dihydro-1H-imidazol-2-yl)-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A93		(2-Fluoro-4-methanesulfonyl-phenyl)-{6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidin-4-yl}-amine
A94		4-[6-(2-Fluoro-4-propoxy-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A95		4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A96		4-{6-[2-Fluoro-4-(2-methoxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A97		4-{6-[2-Fluoro-4-(2-isopropoxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
A98		4-[6-(6-Chloro-4-methyl-pyridin-3-ylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A99		4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-(N-hydroxycarbamimidoyl)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A100		4-[5-Carbamimidoyl-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A101		4-{6-[2-Fluoro-4-(tetrahydro-furan-2-ylmethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A102		4-[5-Methyl-6-(4-methyl-6-morpholin-4-yl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A103		4-{6-[6-(2-Methoxy-ethoxy)-2-methyl-pyridin-3-ylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A104		4-{6-[6-(2-Methoxy-ethoxy)-4-methyl-pyridin-3-ylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A105		4-{6-[2,5-Difluoro-4-(2-methoxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
A106		4-{6-[2-Fluoro-4-(2-isopropoxyethylsulfamoyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A107		4-{6-[2,5-Difluoro-4-(N-hydroxycarbamimidoyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A108		4-[6-(4-Carbamoyl-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A109		4-{6-[(2-Fluoro-4-methanesulfonyl-phenyl)-(2-methoxy-ethyl)-amino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A110		4-[6-(4-Carbamimidoyl-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A111		4-{6-[4-(2-Ethoxy-ethoxy)-2-fluoro-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A112		4-{6-[2-Fluoro-4-(tetrahydro-pyran-4-yloxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A113		4-{6-[2-Fluoro-4-(2-hydroxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
A114		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butan-1-one
A115		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-pentan-1-one
A116		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-butan-1-one
A117		4-{6-[2-Fluoro-4-(pyridin-2-ylmethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
A118		4-[2-(2-Fluoro-4-methanesulfonyl-phenylamino)-3-methyl-pyridin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A119		4-[6-(6-Chloro-4-fluoro-pyridin-3-ylamino)-5-cyano-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
A120		4-[5-Amino-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester

TABLE B

Cmpd#	Structure	Chemical Name
B1		4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-ylamino]-piperidine-1-carboxylic acid tert-butyl ester
B2		N-(4-Methanesulfonyl-phenyl)-5-nitro-N'-piperidin-4-yl-pyrimidine-4,6-diamine
B3		1-[4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-ylamino]-piperidin-1-yl]-ethanone
B4		1-[4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-ylamino]-piperidin-1-yl]-2,2-dimethyl-propan-1-one
B5		4-([6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yl]-isopropyl-amino)-methyl-piperidine-1-carboxylic acid tert-butyl ester

TABLE C

Cmpd#	Structure	Chemical Name
C1		4-(2-Fluoro-4-methanesulfonyl-phenoxy)-6-[1-(3-methoxy-propyl)-piperidin-4-yloxy]-5-methyl-pyrimidine
C2		1-[4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl]-3-methoxy-propan-2-ol

Cmpd#	Structure	Chemical Name
C3		4-[6-(2-Fluoro-4-morpholin-4-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C4		{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-[6-(2-pyrrolidin-1-yl-ethyl)-pyridin-3-yl]-methanone
C5		(6-Amino-pyridin-3-yl)-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-methanone
C6		4-[5-Ethyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C7		4-{6-[2-Fluoro-4-(5-isopropoxymethyl-[1,2,4]oxadiazol-3-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C8		4-{6-[2-Fluoro-4-(5-methoxy-pyridin-2-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C9		4-[6-(2-Fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C10		4-{6-[6-(2-Isopropoxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C11		4-{6-[6-(2-Cyclopropoxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C12		4-{6-[6-(2-Hydroxy-ethylsulfanyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C13		4-{6-[2-Fluoro-4-(pyridine-2-carbonyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C14		4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methanesulfonylamino-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C15		4-[5-Methyl-6-(2-methyl-6-pentyl-pyridin-3-yloxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C16		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(3-fluoro-phenyl)-ethanone
C17		4-[6-(4-Methoxy-6'-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C18		4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(2-pyridin-3-yl-ethyl)-piperidin-4-yloxy]-pyrimidine

Cmpd#	Structure	Chemical Name
C19		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-2-(4-trifluoromethoxy-phenoxy)-propan-1-one
C20		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-2-(4-trifluoromethoxy-phenoxy)-ethanone
C21		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-trifluoromethoxy-phenyl)-ethanone
C22		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-pyridin-2-yl-ethanone
C23		N-(4-Chloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetamide
C24		4-{6-[6-(2-Methoxy-ethanesulfonyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C25		N-(3-Chloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetamide
C26		N-(3,5-Dichloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetamide

Cmpd#	Structure	Chemical Name
C27		4-(2-Fluoro-4-methanesulfonyl-phenoxy)-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine
C28		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-N-(4-trifluoromethyl-phenyl)-acetamide
C29		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-N-phenyl-acetamide
C30		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-N-(4-isopropyl-phenyl)-acetamide
C31		4-(6-{2-Fluoro-4-[(2-hydroxy-ethylcarbamoyl)-methyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C32		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-N-(4-methoxy-phenyl)-acetamide
C33		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-N-(3-trifluoromethyl-phenyl)-acetamide
C34		4-[6-(5-Iodo-pyridin-2-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C35		4-{6-[2-Fluoro-4-(3-methoxypropane-1-sulfonyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C36		4-(6-{2-Fluoro-4-[N-(2-isopropoxyethyl)-carbamimidoyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C37		4-{6-[6-(2-Isopropoxy-ethyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C38		4-[6-(4-Carboxy-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C39		4-{5-Methyl-6-[2-methyl-6-(2-pyridin-2-yl-ethoxy)-pyridin-3-yloxy]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C40		4-(4-Bromo-2-fluoro-phenoxy)-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine
C41		4-{6-[2-Fluoro-4-(thiophene-2-carbonyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C42		4-[6-(5-Methanesulfonyl-pyridin-2-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C43		4-{6-[6-(2-Hydroxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester









Cmpd#	Structure	Chemical Name
C44		4-[5-Cyclopropyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C45		4-(6-{6-[(2-Isopropoxy-ethyl)-methyl-amino]-2-methyl-pyridin-3-yloxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C46		4-{6-[6-(2-Methanesulfonyl-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C47		4-{6-[6-(2-Isopropoxy-ethanesulfonyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C48		4-{6-[6-(2-Hydroxy-ethanesulfonyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C49		4-[6-(6-Amino-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C50		4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(3-methyl-butyl)-piperidin-4-yloxy]-pyrimidine
C51		4-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-oxo-butyric acid

Cmpd#	Structure	Chemical Name
C52		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-morpholin-4-yl-ethanone
C53		1-(3,4-Dichloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C54		1-(3-Chloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C55		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(3-trifluoromethyl-phenyl)-ethanone
C56		4-{6-[6-(2-Methoxy-ethylsulfanyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C57		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-thiophen-3-yl-ethanone
C58		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-phenyl-ethanone
C59		1-(2,4-Dimethoxy-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone

Cmpd#	Structure	Chemical Name
C60		1-(2,5-Dimethoxy-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C61		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-pyridin-2-yl-ethanone
C62		4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(4-methyl-pentyl)-piperidin-4-yloxy]-pyrimidine
C63		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-isopropoxy-propan-1-one
C64		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-isopropoxy-butan-1-one
C65		4-[6-(6-Chloro-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C66		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-hydroxy-propan-1-one
C67		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-fluoro-phenyl)-ethanone

Cmpd#	Structure	Chemical Name
C68		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-trifluoromethyl-phenyl)-ethanone
C69		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(5-pyridin-2-yl-thiophen-2-yl)-ethanone
C70		4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(5-methyl-hexyl)-piperidin-4-yloxy]-pyrimidine
C71		3-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-oxo-propane-1-sulfonic acid
C72		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-thiophen-2-yl-ethanone
C73		4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-(1-pentyl-piperidin-4-yloxy)-pyrimidine
C74		4-(1-Butyl-piperidin-4-yloxy)-6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidine
C75		4-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-cyclohexanecarboxylic acid

Cmpd#	Structure	Chemical Name
C76		1-(4-Diethylamino-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C77		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(2-methyl-4-phenyl-furan-3-yl)-ethanone
C78		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3,3-dimethyl-butan-2-one
C79		4-(2-Fluoro-4-methanesulfonyl-phenoxy)-6-(1-hexyl-piperidin-4-yloxy)-5-methyl-pyrimidine
C80		4-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butyric acid
C81		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-pentan-2-one
C82		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-hexan-2-one
C83		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-hexan-2-one

Cmpd#	Structure	Chemical Name
C84		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-methyl-pentan-2-one
C85		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-5-methyl-hexan-2-one
C86		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-6-methyl-heptan-2-one
C87		5-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-oxo-pentanoic acid
C88		5-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-oxo-pentanenitrile
C89		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-2-pyridin-2-yl-ethanone
C90		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-pyridin-4-yl-ethanone
C91		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-pyridin-3-yl-ethanone

Cmpd#	Structure	Chemical Name
C92		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-ylmethyl}-acrylic acid
C93		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butan-2-one
C94		1-[1,4]Dioxan-2-yl-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C95		1-(2,3-Dihydro-[1,4]dioxin-2-yl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C96		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-p-tolyl-ethanone
C97		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-methoxy-phenyl)-ethanone
C98		1-(2-Chloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C99		3-(2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetyl)-benzonitrile

Cmpd#	Structure	Chemical Name
C100		1-(2,4-Dimethyl-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C101		4-(6-{2-Fluoro-4-[(2-isopropoxyethylcarbamoyl)-methyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C102		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-methanesulfonyl-phenyl)-ethanone
C103		1-(4-Chloro-3-methyl-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C104		1-(4-Difluoromethoxy-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C105		1-(4-Chloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C106		4-(2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetyl)-benzonitrile
C107		1-(3,4-Difluoro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone

Cmpd#	Structure	Chemical Name
C108		1-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C109		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(5-phenyl-thiophen-2-yl)-ethanone
C110		2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-thiophen-2-yl-ethanone
C111		{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetic acid ethyl ester
C112		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methoxy-propan-2-ol
C113		4-{6-[2-Fluoro-4-(2-isopropoxy-ethylcarbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C114		4-(2-Fluoro-4-methanesulfonyl-phenoxy)-6-[1-(4-methoxy-cyclohexyl)-piperidin-4-yloxy]-5-methyl-pyrimidine
C115		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butan-1-one

Cmpd#	Structure	Chemical Name
C116		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-pentan-1-one
C117		4-[6-(2,4-Difluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C118		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-hexan-1-one
C119		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-butan-1-one
C120		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-methyl-pentan-1-one
C121		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-5-methyl-hexan-1-one
C122		4-{6-[2-Fluoro-4-(2-methoxyethylcarbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C123		4-{6-[2-Fluoro-4-(2-isobutoxyethoxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C124		4-{6-[4-(2-Cyclopropoxy-ethoxy)-2-fluoro-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C125		4-{6-[4-(2-Ethoxy-ethoxy)-2-fluoro-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C126		4-{6-[2-Fluoro-4-(3-methoxy-propoxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C127		4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C128		4-{6-[2-Fluoro-4-(2-pyridin-2-yl-ethoxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C129		4-{6-[2-Fluoro-4-(tetrahydro-pyran-4-yloxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C130		4-[6-(4-Bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C131		4-{6-[4-(2-tert-Butoxy-ethoxy)-2-fluoro-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C132		4-{6-[2-Fluoro-4-(methoxy-methyl-carbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C133		1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methoxy-propan-1-one
C134		4-[6-(4-Cyano-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C135		4-[5-(5-Aminomethyl-4,5-dihydro-oxazol-2-yl)-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C136		4-{6-[6-(2-Methoxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C137		4-{6-[6-(3-Methanesulfonyl-pyrrolidin-1-yl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C138		4-[6-(6-Benzylamino-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C139		4-[6-(4-Carbamoyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C140		4-{6-[2-Fluoro-4-(2-isopropoxy-ethylamino)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C141		4-(6-{2-Fluoro-4-[(tetrahydrofuran-2-ylmethyl)-amino]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C142		4-(6-{6-[(2-Methanesulfonyl-ethyl)-methyl-amino]-2-methyl-pyridin-3-yloxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C143		4-[6-(2-Fluoro-4-hydroxycarbonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C144		4-{6-[2-Fluoro-4-(2-pyrrolidin-1-ylethylcarbonyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C145		4-{6-[2-Fluoro-4-(4-isopropylpiperazine-1-carbonyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C146		4-{6-[2-Fluoro-4-(2-morpholin-4-ylethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C147		4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C148		4-{6-[2-Fluoro-4-(2-hydroxy-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C149		4-[6-(4-Carboxymethyl-2-fluorophenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C150		4-[6-(4-Dimethylcarbamoylmethyl-2-fluorophenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C151		4-[6-(2-Fluoro-4-sulfamoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C152		4-[6-(2-Fluoro-4-propionylsulfamoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C153		4-[5-Ethynyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C154		4-[6-[2-Fluoro-4-(2-phosphonoxyethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C155		4-[5-Bromo-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C156		4-(6-{2-Fluoro-4-[2-(2-methanesulfonyl-pyrrolidin-1-yl)-2-oxo-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester









Cmpd#	Structure	Chemical Name
C157		4-[6-(4-Carbamoylmethyl-2-fluorophenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C158		4-[6-(2-Fluoro-4-[(tetrahydrofuran-2-ylmethyl)-carbamoyl]-methyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C159		4-[6-(2-Fluoro-3-sulfamoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C160		C-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-C-(4-fluorophenyl)-methyleamine
C161		3-tert-Butoxy-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-propan-1-one
C162		4-[6-(2-Fluoro-4-sulfo-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C163		2-Ethoxy-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone
C164		{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-(tetrahydrofuran-2-yl)-methanone

Cmpd#	Structure	Chemical Name
C165		(S)-1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-2-methylamino-butan-1-one
C166		4-(6-{2-Fluoro-4-[2-(3-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C167		4-{6-[2-Fluoro-4-(2-morpholin-4-yl-2-oxo-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C168		4-{6-[2-Fluoro-4-(2-imidazol-1-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C169		4-{6-[2-Fluoro-4-(2-[1,2,3]triazol-1-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C170		(R)-1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-2-methylamino-butan-1-one
C171		(S)-1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-hydroxy-butan-1-one
C172		(R)-N-(1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carbonyl}-2-methyl-propyl)-acetamide

Cmpd#	Structure	Chemical Name
C173		(S)-N-(1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carbonyl}-2-methyl-propyl)-acetamide
C174		(R)-N-(2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-methyl-2-oxo-ethyl)-acetamide
C175		(S)-N-(2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-methyl-2-oxo-ethyl)-acetamide
C176		4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid (S)-tetrahydro-furan-3-yl ester
C177		4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid (R)-tetrahydro-furan-3-yl ester
C178		4-[6-(2-Amino-4-ethanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C179		4-[6-(4-Methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C180		(1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carbonyl}-2-methyl-propyl)-carbamic acid tert-butyl ester

Cmpd#	Structure	Chemical Name
C181		4-{6-[2-Fluoro-4-(6-methoxy-pyridin-3-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C182		3-Amino-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-methyl-pentan-1-one
C183		2-Amino-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-butan-1-one
C184		4-{6-[2-Fluoro-4-(2-isopropoxy-ethoxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C185		4-[5-Methyl-6-(4-sulfo-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C186		4-[6-(2,5-Difluoro-4-trifluoromethoxy-phenoxy)-5-ethynyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C187		4-[6-(2,5-Difluoro-4-trifluoromethoxy-phenoxy)-5-prop-1-ynyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C188		4-[5-Ethynyl-6-(2-fluoro-4-methoxy-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C189		4-[5-Ethynyl-6-(6-methoxy-4-methyl-pyridin-3-yloxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C190		4-{5-Ethynyl-6-[6-(2-isopropoxy-ethyl)-2-methyl-pyridin-3-yloxy]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C191		4-[6-(4-Cyano-2-fluoro-phenoxy)-5-ethynyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C192		4-[5-Ethynyl-6-(2-fluoro-4-[1,2,4]triazol-4-yl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C193		4-[5-Ethynyl-6-(2-fluoro-4-[1,2,4]triazol-1-yl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C194		1-{4-[5-Ethynyl-6-(2-fluoro-4-[1,2,4]triazol-1-yl-phenoxy)-pyrimidin-4-yloxy]-piperidin-1-yl}-3-pyridin-2-yl-propan-1-one
C195		4-{5-Ethynyl-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-pyrimidin-4-yloxy}-3-fluoro-benzonitrile
C196		5-Ethynyl-4-(2-fluoro-4-methanesulfonyl-phenoxy)-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-pyrimidine

Cmpd#	Structure	Chemical Name
C197		4-[1-(3-Ethyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-ethynyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidine
C198		4-[1-(3-Ethyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidine
C199		4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(3-methyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-pyrimidine
C200		4-[6-(2-Fluoro-4-methanesulfonylamino-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C201		<i>cis</i> -{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-cyclohexyl}-carbamic acid isopropyl ester
C202		<i>trans</i> -{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-cyclohexyl}-carbamic acid isopropyl ester
C203		N-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-cyclohexyl}-3-methyl-butyramide
C204		N-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-cyclohexyl}-isobutyramide

Cmpd#	Structure	Chemical Name
C205		4-{6-[2,5-Difluoro-4-(2-methanesulfonyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C206		4-{6-[4-Fluoro-6-(2-methanesulfonyl-ethyl)-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C207		4-{5-Cyclopropyl-6-[2,5-difluoro-4-(2-hydroxy-ethyl)-phenoxy]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C208		4-(5-Cyclopropyl-6-{2,5-difluoro-4-[2-(4-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C209		4-{6-[2,5-Difluoro-4-(2-morpholin-4-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C210		4-(6-{2-Fluoro-4-[2-(4-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C211		4-{6-[6-(2-Fluoro-ethyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C212		4-{6-[2-Fluoro-4-(1-hydroxy-cyclopropylmethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C213		4-{2-[2,5-Difluoro-4-(2-methanesulfonyl-ethyl)-phenoxy]-3-methyl-pyridin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C214		(R)-4-(6-{2-Fluoro-4-[2-(3-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C215		(S)-4-(6-{2-Fluoro-4-[2-(3-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C216		(R)-4-(5-Ethynyl-6-{2-fluoro-4-[2-(2-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C217		(S)-4-(2-{2-Fluoro-4-[2-(2-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-3-methyl-pyridin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C218		4-{6-[4-Fluoro-6-(2-morpholin-4-yl-ethyl)-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C219		4-{5-Ethynyl-6-[4-fluoro-6-(2-methanesulfonyl-ethyl)-pyridin-3-yloxy]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C220		4-{2-[2,5-Difluoro-4-(2-isopropoxy-ethyl)-phenoxy]-3-methyl-pyridin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C221		4-{6-[2-Fluoro-4-(2-propionylsulfamoyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C222		4-{6-[2-Fluoro-4-(2-sulfamoyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C223		4-{6-[2,5-Difluoro-4-(2-sulfamoyl-ethyl)-phenoxy]-5-ethynyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C224		4-{6-[2,5-Difluoro-4-(2-[1,2,4]triazol-1-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C225		4-{6-[2,3-Difluoro-4-(2-methanesulfonyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C226		4-(2-{2-Fluoro-4-[2-(6-methoxy-pyridin-2-yl)-ethyl]-phenoxy}-3-methyl-pyridin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C227		4-(6-{2-Fluoro-4-[2-(3-methoxy-pyridin-2-yl)-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C228		4-[6-(3-Fluoro-1-oxy-pyridin-4-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C229		4-[6-(5'-Methoxy-6-methyl-[2,2']bipyridinyl-5-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C230		4-{5-Ethynyl-6-[2-fluoro-4-(4-methoxy-pyridin-2-yl)-phenoxy]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C231		4-{6-[2-Fluoro-4-(3-methoxy-pyridin-2-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C232		4-(6-{2,5-Difluoro-4-[2-(3-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C233		4-(6-{2,5-Difluoro-4-[2-(3-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-5-ethynyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester
C234		4-{6-[2-Fluoro-4-(5-methoxy-pyridin-3-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

Cmpd#	Structure	Chemical Name
C235		4-[6-(2-Fluoro-4-pyridin-4-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C236		4-[6-(3-Fluoro-biphenyl-4-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C237		4-[6-(2-Fluoro-4-pyridin-3-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C238		4-[6-(2-Fluoro-4-pyrimidin-5-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C239		4-[6-(2-Fluoro-4-thiophen-3-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C240		4-[6-(4-Ethynyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester
C241		(R)-4-{6-[2-Fluoro-4-(2-oxo-oxazolidin-4-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester
C242		(S)-4-{6-[2-Fluoro-4-(2-oxo-oxazolidin-4-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester

TABLE D

Cmpd#	Structure	Chemical Name
D1		4-((Cyclopropyl-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-amino)-methyl)-piperidine-1-carboxylic acid tert-butyl ester
D2		4-((Cyclopropyl-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-amino)-methyl)-piperidine-1-carboxylic acid isopropyl ester
D3		4-([6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-isopropyl-amino)-methyl)-piperidine-1-carboxylic acid isopropyl ester
D4		4-((Cyclopropylmethyl-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-amino)-methyl)-piperidine-1-carboxylic acid isopropyl ester

TABLE E


Cmpd#	Structure	Chemical Name
E1		4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-ylsulfanyl]-piperidine-1-carboxylic acid isopropyl ester

5 Additionally, compounds of the present invention encompass all pharmaceutically acceptable salts, solvates, and particularly hydrates, thereof.


General Synthetic Methods

10 As a result of their profound biological significance in higher eukaryotes and utilization of the pyrimidine core in a number of marketed drugs (**Scheme 1**) and other medicinally relevant compounds, pyrimidines and pyridines play pivotal roles as chemotypes in drug discovery campaigns. As a direct consequence of this there is a wealth of scientific literature describing synthetic construction, as well as chemical modification and elaboration of these classes of heterocycles.

Scheme 1




Scheme 1



The novel substituted pyrimidines and pyridines of the present invention can be prepared according to a variety of synthetic manipulations, all of which would be familiar to one skilled in the art of synthetic organic chemistry. Certain methods for the preparation of compounds of the present invention include, but are not limited to, those described in **Schemes 2-13** set forth in this section of the specification and in the Examples, *infra*.

Common dihalo-substituted intermediates **9.1** and **9.2**, used as a starting point for the synthesis of compounds of the present invention are commercially available or can be prepared by methods know in the art, for example as depicted in **Scheme 2a**.

Scheme 2a




This is accomplished in two steps from a di- C_{1-6} -alkylmalonate, one particularly useful di- C_{1-6} -alkylmalonate is diethyl malonate **5**. Cyclization to the C5-Z-substituted-4,6-dihydroxypyrimidine **8** is achieved via an initial alkali metal base catalysed deprotonation, alkylation strategy or via generation of the monoanion using sodium / EtOH followed by alkylation using Z-Hal, subsequent reaction of monoalkyl species **6** with formamidine in the presence of an alkali metal alkoxide, by

mixing the malonate and all or part of the formamidine with the alkoxide or with the alkoxide and the rest of the formamide. Alternative reagents such as dimethylmalonate, sodium methoxide, formamide, in low molecular weight alcoholic solvents, including methanol, ethanol, 2-propanol and the like, may be utilized in the synthesis by heating at a temperature range between about 80°C to about 100°C for about 30 mins to about 90 mins followed by a mineral acid work up. In a preferred variation chlorinated intermediate **6.1** may be used as a starting point to obtain pyrimidines wherein alternate C5 substituents such as R_5R_6N are introduced by performing thermal nucleophilic displacements. Preparation of dihydroxypyrimidines can also be achieved using microorganisms such as *Rhodococcus* (see for reference WO97008152 A1). An ortho metallation strategy can be utilized to facilitate C3-alkylation of the corresponding 2,4-dichloropyridyl core **15**. Using *n*-BuLi at -78 °C under anhydrous / inert conditions followed by trapping of the resulting monoanion with an appropriate alkyl bromide or iodide (**Scheme 2c**) [for references see Mongin, F.; Queguiner, G. Advances in the directed metallation of azines and diazines (pyridines, pyrimidines, pyrazines, pyridazines, quinolines, benzodiazines and carbolines). Part 1: Metallation of pyridines, quinolines and carbolines. Tetrahedron (2001), 57(19), 4059-4090. Turck, A.; Ple, N.; Mongin, F.; Queguiner, G. Advances in the directed metalation of azines and diazines (pyridines, pyrimidines, pyrazines, pyridazines, quinolines, benzodiazines and carbolines). Part 2. Metalation of pyrimidines, pyrazines, pyridazines and benzodiazines. Tetrahedron (2001), 57(21), 4489-4505]


Chlorination of the 4 and 6 ring positions to produce intermediate **8** maybe carried out by reacting **8** with a chlorinating reagent, such as, phosgene, $POCl_3$ (for reference see A. Gomtsyan *et al.*, J. Med. Chem. 2002, 45, 3639-3648), thionyl chloride, oxalyl chloride and by mixtures of the above reagents including PCl_3 / $POCl_3$ at elevated reaction temperatures.

Scheme 2b



In some embodiments of the current invention alternate functionalities are required at the C5 pyrimidinyl position to achieve the desired biological outcome. Such functionality may be introduced via a broad range of organic synthetic procedures. Some examples are depicted in **Scheme 2b**,


wherein common intermediate **10** can be converted to intermediates such as **11**, **12**, **13** by synthetic chemistries familiar to those in the art. **Schemes 2bii** and **2biii** are initially reliant upon the one pot chlorinating formylation variant of the Villsmeier-Haack reaction which introduces “synthetic handles” at ring positions 3, 4 and 5 of the core simultaneously (for references see; Chlorinating formylation reactions with pyrimidines. Kloetzer, W.; Herberz, M., Monatshefte fuer Chemie (1965), 96(5), 1567-72. Also see Gontsyan *et al* Journal of Medicinal Chemistry, 2002, 45, 3639 – 3648 and references therein). Wherein Z = nitro, commercially available 2,6-dichlor-5-nitropyrimidine was utilized. Where necessary all dichloro intermediate pyrimidines (**9.1**, **11**, **12**, **13** etc) that are used as core building blocks in the present invention may be optionally converted to 4,6-diiodopyrimidines by halo exchange using sodium iodide and 45 % hydroiodic acid as depicted in **Scheme 2a**.



Conventional thermal aromatic substitution reactions of amines and alcohols with halogenated pyrimidines have been well documented (see for example A. G. Arvanitis *et al.*, J. Medicinal Chemistry, 1999, 42, 805–818 and references therein). Nucleophilic aromatic ($\text{S}_{\text{N}}\text{Ar}$) substitution reactions of electron deficient halogenated pyrimidines are usually rapid and high yielding. However, in certain cases, such as electron rich or neutral halogenated heterocycles, successful substitution is afforded by prolonged heating.

To facilitate rapid entry into many of the compounds of the invention microwave synthesis was utilized (**Schemes 3 and 4**). The Smith synthesizer from Personal Chemistry is a commercially available focused field heating instrument that provides safer and more uniform conditions for performing the base catalyzed substitution reactions depicted in **Schemes 3a**, **3b** and **3c**. Bases employed for such conversions (whereby Q = N) include tertiary amines such as triethylamine, Hunig's base (i.e. diisopropyl-ethylamine), N-methylmorpholine and the like. Alternatively, one skilled in the art can employ alkali metal hydrides, alkali metal carbonates (such as, Li_2CO_3 , Na_2CO_3 , K_2CO_3 and the like), an alkali metal hydrogencarbonate (such as, LiHCO_3 , NaHCO_3 , KHCO_3 and the like). Wherein Q = N, inert lower alkyl alcoholic solvent can be employed (such as, MeOH, EtOH, i-PrOH, n-BuOH and the like) or wherein Q = O, an ethereal solvent such as tetrahydrofuran, 1,4-dioxane, and the like can be used. Reaction times to access typical monosubstituted intermediates such as, **15** and **16**, can range from about 300 s to about 3000 s and when conventional thermal methods are employed (wherein Q = O) about 20 mins to about 120 mins.


Scheme 3



Methods for conversion of intermediate monosubstituted pyrimidines and pyridines 15 and 16 are illustrated in Scheme 4. Examples wherein Q = NR₆ (Schemes 4a, 4b and 4d) were obtained using palladium catalysed aminations. This synthetic strategy has emerged as a powerful tool for synthesis of substituted aryl and heteroaryl anilines in recent times (for reference see S. L. Buchwald., Top. Curr. Chem., 2002, 219, 131 and references therein). Reaction of a suitably substituted amine (such as, intermediate 17) in the presence of a palladium or alternative transition metal catalyst selected from but not limited to Pd₂(dba)₃, Pd(OAc)₂, CuI, Cu(OTf)₂, Ni(COD)₂, Ni(acac)₂ in a suitable anhydrous solvent (such as, THF, 1,4-dioxane, and the like) with as strong alkali metal alkoxide base (such as, NaO^tBu, KO^tBu and the like). A suitable ligand employed in this step can be selected from BINAP, P(o-tolyl)₃, tBu₃P, DPPF, P[N(^tBu)CH₂CH₃]₃N and the like when the catalyst is a palladium derived complex.

Alternatively, for "Ullman-type" aryl aminations catalysed by copper derived complexes the base employed maybe selected from an alkali metal carbonate in an aprotic polar solvent (such as N,N-dimethylacetamide, DMF, DMSO, and the like) with L-proline, N-methylglycine or diethylsalicyclamide as the ligand (for reference see D. Ma, Organic Lett., 2003, 5, 14, 2453 – 2455).

Scheme 4




Compounds of general formula 19 to 22 may also be obtained by reversing the order of the reaction steps (i.e. introduction of W followed by Q), wherein the initial step comprises of introduction of either Intermediate 17 or 18 by using base in $^i\text{PrOH}$ followed by addition of 4N HCl in dioxane.

As illustrated in Scheme 5, a similar transition metal catalysed couplings were utilized to obtain molecules of general formula 24 and 27 (Scheme 5.1) wherein the Ar_1 substituent ($\text{Hal} = \text{Br}, \text{I}$) of intermediate 23 is modified to give analogs with alkyl amino substituents (i.e., NR_aR_b , wherein R_a and R_b are each independently H, C_{1-6} alkyl or a substituted C_{1-6} alkyl, or R_a and R_b together with the nitrogen form a heterocyclic ring, as described herein). Alternatively, the linker atom can be oxygen by utilizing the CuI catalysed method for aromatic C-O formation described by Buchwald (see for reference S. L. Buchwald; Organic Lett., 2002, 4, 6, 973-976) by utilizing, for example, 10 mol % CuI, 20 mol% 1,10-phenanthroline, 2 equivalents of Cs_2CO_3 , at 110°C for 18 h (Scheme 5d), with an


Ar₁ iodo substitution in the substrate. Additional important organometallic transformations from halo intermediates **23** to active analogues of the current invention include the well know palladium catalyzed couplings of appropriately substituted aryl boronic acids via the "Suzuki coupling reaction" (Scheme 5e).

5

Scheme 5.1




10




The Suzuki coupling represents a widely used method for the synthesis of biaryl compounds and is already applied on industrial scale. For a long time this reaction was limited to the use of aryl bromides, aryl iodides or electron-deficient aryl chlorides as starting materials. Thus, a general approach to the desired biaryl compounds using the cheap and easy available aryl chlorides was not available. In the last two years, however, several new protocols for the Suzuki coupling with aryl chlorides were developed. These methods allow an efficient synthesis of biaryls, independently of the substitution pattern and electronic properties of the starting materials. These concepts which were developed by the research groups of Fu, Buchwald, Guram, Beller as well as Trudell and Nolan are highlighted in "Modern methods of the Suzuki cross coupling: the long expected general synthetic routes using aryl chlorides. Groger, Harald, Journal fuer Praktische Chemie (Weinheim, Germany) (2000), 342(4), 334-339. Alternatively additional functionality maybe introduced using other metal catalyzed transformations such as cyanation using zinc(II)cyanide under microwave irradiation conditions to obtain compounds of general formula 25 or the well documented Pd catalyzed "Sonogashira reaction" (Scheme 5c) for introduction of terminal alkynes. Most recently the Sonogashira Coupling has been described to produce almost quantitative yields of desired product using appropriate reaction conditions in the complete absence of palladium catalysts (for ref see "First Examples of Transition-Metal Free Sonogashira-Type Couplings" Leadbeater, Nicholas E.; Marco, Maria; Tominack, Bonnie J, Organic Letters (2003), 5(21), 3919-3922, and also also Transition-metal-free Sonogashira-type coupling reactions in water, Appukkuttan, Prasad; Dehaen, Wim; Van der Eycken, Erik, European Journal of Organic Chemistry (2003), (24), 4713-4716. In other preferred embodiments of the present invention, such organotransition metal chemistries may be used to introduce similar functional groups to the C5 position or the C3 position of the respective pyrimidine and pyridyl cores. For example C5 bromo or iodo intermediates may be cyanated or alkynylated as depicted in Schemes 5.2 and 5.3. Indeed, advanced nitrile derivatives of the present invention may be optionally modified via synthetic manipulations outlined in Schemes 5.1f and Schemes 5.2a-c.

Scheme 5.2



One particular embodiment is when the Hal group on Ar is located at the para position of a phenyl ring (Ar). In another particular embodiment of the invention, the Hal group is chloro at the 2 position of a trisubstituted pyridyl moiety (intermediate 28). Organotransition metal catalysed methods for substitution of this halogen are depicted in Scheme 6.


Scheme 5.3



A particular substitution for compounds 19-29 is wherein $D = \text{NCOOR}_c$ wherein R_c is C_{1-6} alkyl, or C_{3-7} cycloalkyl and each can be further substituted. Urethanes of this type can be prepared directly from intermediates depicted in Schemes 3 and 4 when $D = \text{NH}$. In certain reactions, use of a suitable nitrogen protecting group (such as, 'Boc, Cbz, Moz, Alloc, Fmoc and the like) may be necessary during further chemical modification of the core. Deprotection may be achieved using standard reagents familiar to one skilled in the art (these might include TFA, mineral acid, Palladium / hydrogen gas and the like in an alcoholic or ethereal solvent system chosen from methanol, ethanol, *tert*-butanol, THF, 1,4-dioxane, and the like). On occasion wherein the target molecule contains 2

protecting groups, an orthogonal protection strategy may be adopted. The deprotected secondary amine (D = NH) can subsequently be modified accordingly.

Scheme 6



Schemes 7 and 8 and 9 illustrate such chemistries wherein generation of a carbamate, urea or amide can be executed using an appropriate reaction in the presence of a base, for example, a tertiary amine base such as TEA, DIEA and the like, in an inert solvent system.


As illustrated in Scheme 7, urethane 19 can be obtained by a urethane reaction using R_cOCO- halide (wherein R_c is as described *supra*, and halide is chloro, bromo, or iodo, particularly useful is chloro) in an inert solvent with or without a base. Suitable bases include an alkali metal carbonate (such as, sodium carbonate, potassium carbonate, and the like), an alkali metal hydrogencarbonate (such as, sodium hydrogencarbonate, potassium hydrogencarbonate, and the like), an alkali hydroxide (such as, sodium hydroxide, potassium hydroxide, and the like), a tertiary amine (such as, *N,N*-diisopropylethylamine, triethylamine, *N*-methylmorpholine, and the like), or an aromatic amine (such as, pyridine, imidazole, poly-(4-vinylpyridine), and the like). The inert solvent includes lower halocarbon solvents (such as, dichloromethane, dichloroethane, chloroform, and the like), ethereal solvents (such as, tetrahydrofuran, dioxane, and the like), aromatic solvents (such as, benzene, toluene, and the like), or polar solvents (such as, *N,N*-dimethylformamide, dimethyl sulfoxide, and the like). Reaction temperature ranges from about -20°C to 120°C, preferably about 0°C to 100°C.

10

15

20

Scheme 7




As shown in Scheme 8a, the amine intermediate obtained from acidic deprotection of 30 can be functionalized to amides represented by species 32. Carbamate 20 is first reacted with 4N HCl in dioxane or alternatively TFA in dichloromethane and further reacted with a carboxylic acid (R_dCO_2H ,

25

wherein as used in **Scheme 8a**, R_d is Ar, or a C_{1-6} -alkylene-Ar; Ar can be substituted or unsubstituted and has the same meaning as described herein) with a dehydrating condensing agent in an inert solvent with or without a base to provide the amide **23** of the present invention. The dehydrating condensing agent includes dicyclohexylcarbodiimide (DCC), 1,3-diisopropylcarbodiimide (DIC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC•HCl), bromo-tris-pyrrolidino-phosphonium hexafluorophosphate (PyBroP), benzotriazoloyloxytris(dimethylamino)-phosphonium hexafluorophosphate (BOP), *O*-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU), or 1-cyclohexyl-3-methylpolystyrene-carbodiimide. The base includes a tertiary amine (such as, *N,N*-diisopropylethylamine, triethylamine, and the like). The inert solvent includes lower halocarbon solvents (such as, dichloromethane, dichloroethane, chloroform, and the like), ethereal solvents (such as, tetrahydrofuran, dioxane, and the like), nitrile solvents (such as, acetonitrile, and the like), amide solvents (*N,N*-dimethylformamide, *N,N*-dimethylacetamide, and the like) and mixtures thereof. Optionally, 1-hydroxybenzotriazole (HOBT), HOBT-6-carboxaamidomethyl polystyrene, or 1-hydroxy-7-azabenzotriazole (HOAT) can be used as a reactant agent. Reaction temperature ranges from about -20 °C to 50 °C, preferably about 0 °C to 40 °C.

Scheme 8



Alternatively, amides **32** of the present invention can be obtained by an amidation reaction using an acid halide (such as, R_dCOCl) and a base in an inert solvent (**Scheme 8a**). The base includes an alkali metal carbonate (such as, sodium carbonate, potassium carbonate, and the like), an alkali metal hydrogencarbonate (such as, sodium hydrogencarbonate, potassium hydrogencarbonate, and the

like), an alkali hydroxide (such as, sodium hydroxide or potassium hydroxide, and like), a tertiary amine (such as, *N,N*-diisopropylethylamine, triethylamine, *N*-methyldmorpholine, and the like), or an aromatic amine (such as, pyridine, imidazole, poly-(4-vinylpyridine), and the like). The inert solvent includes lower halocarbon solvents (such as, dichloromethane, dichloroethane, chloroform, and the like), etheral solvents (such as, tetrahydrofuran, dioxane, and the like), amide solvents (such as, *N,N*-dimethylacetamide, *N,N*-dimethylformamide, and the like), aromatic solvents (benzene, toluene, pyridine, and the like) and mixtures thereof. Reaction temperature ranges from about -20 °C to 50 °C, preferably about 0 °C to 40 °C.

Also illustrated in **Scheme 8**, amide **32** can be reacted with a reducing agent in an inert solvent to provide the amine **33** of the present invention. The reducing agent includes alkali metal aluminum hydrides (such as, lithium aluminum hydride, and the like), alkali metal borohydrides (such as, lithium borohydride, and the like), alkali metal trialkoxyaluminum hydrides (such as, lithium tri-*tert*-butoxyaluminum hydride, and the like), dialkylaluminum hydrides (such as, di-isobutylaluminum hydride, and the like), borane, dialkylboranes (such as, di-isoamyl borane, and the like), alkali metal trialkylboron hydrides (such as, lithium triethylboron hydride, and the like). The inert solvent includes etheral solvents (such as, tetrahydrofuran, dioxane, and the like), aromatic solvents (such as, toluene, and the like) and mixtures thereof. Reaction temperature ranges from about -78 °C to 200 °C, such as, about 50 °C to 120 °C.

Alternatively, the amine **33** of the present invention can be obtained by a reductive amination reaction using the acid deprotected secondary amine intermediate with an aldehyde ($R_6\text{CHO}$) and a reducing agent in an inert solvent with or without an acid. The reducing agent includes sodium triacetoxyborohydride, sodium cyanoborohydride, sodium borohydride, borane-pyridine complex, and the like. The inert solvent includes lower alkyl alcohol solvents (such as, methanol, ethanol, and the like), lower halocarbon solvents (such as, dichloromethane, dichloroethane, chloroform, and the like), etheral solvents (such as, tetrahydrofuran, dioxane, and the like), aromatic solvents (such as, benzene, toluene, and the like) and mixtures thereof. The acid includes an inorganic acid (such as, hydrochloric acid, sulfuric acid, and the like) or an organic acid (such as, acetic acid, and the like). Reaction temperature ranges from about -20 °C to 120 °C, preferably about 0 °C to 100 °C. In addition, this reaction can optionally be carried out under microwave conditions.

In an alternative manner, the intermediate amine product of acid deprotection of **30** can be alkylated directly with an alkylating agent, such as R_6 -halide (wherein R_6 is substituted or unsubstituted C_{1-6} alkyl, or substituted or unsubstituted C_{1-6} alkyl-Ar, and halide is chloro, bromo and iodo), in the presence of a base and in an inert solvent to provide amine **33**. The base includes an alkali metal carbonate (such as, sodium carbonate, potassium carbonate, and the like), an alkali metal hydride (such as, sodium hydride, potassium hydride, and the like), alkali metal alkoxide (such as, potassium *tert*-butoxide, sodium *tert*-butoxide, and the like); alkyl lithiums (such as, *tert*-butyl lithium, *n*-butyl lithium and the like). The inert solvents include, etheral solvents (such as,

tetrahydrofuran, dioxane), aromatic solvents (such as, benzene, toluene, and the like), amide solvents (such as, *N,N*-dimethylformamide, and the like) and mixtures thereof. Reaction temperature ranges from about -20 °C to 120 °C, preferably about 0 °C to 100 °C.


Also shown in Scheme 8 is the preparation of additional compounds of the invention via
5 alkylating the nitrogen of ureas represented by 32 with an alkyl-halide (wherein halide is chloro, bromo and iodo) in the presence of a base in an inert solvent to provide di-substituted urea. The base includes an alkali metal hydride (such as, sodium hydride, potassium hydride, and the like), alkali metal alkoxide (such as, potassium *tert*-butoxide, sodium *tert*-butoxide, and the like); alkyl lithiums (such as, *tert*-butyl lithium, *n*-butyl lithium and the like). The inert solvents include, ethereal solvents
10 (such as, tetrahydrofuran, dioxane), aromatic solvents (such as, benzene, toluene, and the like), amide solvents (such as, *N,N*-dimethylformamide, and the like) and mixtures thereof. Reaction temperature ranges from about -20 °C to 120 °C, preferably about 0 °C to 100 °C.

In addition, as illustrated in Scheme 9a, urea 34 can be obtained from deprotecting common intermediate 30 and allowing the amine (i.e., D = NH) to react with a variety isocyanates ($R_a\text{NCO}$,
15 wherein R_a has the same meaning as described herein) in an inert solvent with or without a base. Suitable bases include an alkali metal carbonate (such as, sodium carbonate, potassium carbonate, and the like), an alkali metal hydrogencarbonate (such as, sodium hydrogencarbonate, potassium hydrogencarbonate, and the like), an alkali hydroxide (such as, sodium hydroxide, potassium hydroxide, and the like), a tertiary amine (such as, *N,N*-diisopropylethylamine, triethylamine, *N*-methylmorpholine, and the like), or an aromatic amine (such as, pyridine, imidazole, and the like).
20 The inert solvent includes lower halocarbon solvents (such as, dichloromethane, dichloroethane, chloroform, and the like), ethereal solvents (such as, tetrahydrofuran, dioxane, and the like), aromatic solvents (such as, benzene, toluene, and the like), or polar solvents (such as, *N,N*-dimethylformamide, dimethyl sulfoxide, and the like). Reaction temperature ranges from about -20 °C to 120 °C,
25 preferably about 0 °C to 100 °C.

30

35


Scheme 9



Further, as illustrated in Scheme 9b, thiourea 35 can be obtained from deprotecting common intermediate 30 and allowing the amine (i.e., D = NH) to react with a variety of thioisocyanates ($R_a\text{NCS}$, wherein R_a has the same meaning as described herein) in an inert solvent with or without a base. Suitable bases include an alkali metal carbonate (such as, sodium carbonate, potassium carbonate, and the like), an alkali metal hydrogencarbonate (such as, sodium hydrogencarbonate, potassium hydrogencarbonate, and the like), an alkali hydroxide (such as, sodium hydroxide, potassium hydroxide, and the like), a tertiary amine (such as, *N,N*-diisopropylethylamine, triethylamine, *N*-methylmorpholine, and the like), or an aromatic amine (such as, pyridine, imidazole, and the like). The inert solvent includes lower halocarbon solvents (such as, dichloromethane, dichloroethane, chloroform, and the like), ethereal solvents (such as, tetrahydrofuran, dioxane, and the like), aromatic solvents (such as, benzene, toluene, and the like), or polar solvents (such as, *N,N*-dimethylformamide, dimethyl sulfoxide, and the like). Reaction temperature ranges from about -20 °C to 120 °C, preferably about 0 °C to 100 °C.


Scheme 10 illustrates the synthesis of para-alkyl sulfones (37) which are used as aryl building blocks in Scheme 4 of the present invention, wherein R_{10} - R_{13} have the same meaning as described herein. The common methods for preparing these sulfones include the oxidation of sulfides or the sulfonylation of arenes using aryl sulfonyl halides or aryl sulfonic acids in the presence of a strong acid catalyst (see for general reference: the Organic Chemistry of Sulfur; Oae S., Ed.; Plenum Press: New York, 1977). Optimal conversion to the optionally 2,5-disubstituted arene 37 was achieved thermally wherein Hal is preferably iodo using 5 mol % $(\text{CuOTf})_2\cdot\text{PhH}$ and 10 mol % *N,N'*-dimethylethylenediamine in DMSO by the method of Wang et al (see for reference Wang Z.; Baskin J. M., Org. Lett., 2002, 4, 25, 4423-4425). In some embodiments, R_{10} and R_{13} are each independently H, halogen, or C_{1-6} alkyl; R_{11} and R_{12} are both H; Hal = Br, I; and Q1 = OH, or NH_2 .

Scheme 10




Alternative standard organic synthetic methods may be used to introduce alternate substituents in to the Ar component. In one example wherein the linker atom is Q = N, the manipulation maybe carried out by protecting the aniline amino functionality using standard FmocCl and CbzCl protection deprotection steps familiar to one skilled in the art (Scheme 11, wherein R₁₀-R₁₃ have the same meaning as described herein) and subsequently using the deprotected aniline in subsequent steps such as those depicted in Scheme 4. Nitrile 39, maybe alternatively transformed in to amidines (see Table of compounds) by using hydroxylamine HCl followed by reduction using zinc / acetic acid. In some embodiments of the invention R₁₀ is halogen, and R₁₃ is H or halogen.

Scheme 11




Synthetic scheme 11.1, depicts some of organic synthetic strategies of the current invention for accessing advanced aromatic building blocks required for use in scheme 4c wherein R₁₀-R₁₃ are preferably halogen, alkoxy or short alkyl. Following incorporation in to analogues of the present invention via methodologies depicted in scheme 4c, intermediates such as those of type 38.3 may be deprotected through use of suitable silyl deprotection agents such as TBAF or HF. Resulting terminal alcohols may be optionally further modified (for ref see T. Matsui *et al.*, Biorg. Med. Chem, 10, 2002, 3787).

Scheme 11.1




Scheme 11.2



5

Synthesis of the 3,5-oxadiazolo variant is depicted in **Scheme 12**. Zinc(II)chloride catalyzed coupling of amidoxime **44** with 4-hydroxypiperidine, CNBr derived **46** yielded building block **47** after acidic workup, which was subsequently utilized in reaction sequences depicted as illustrated in **Scheme 3**.



10

In a preferred embodiment of the present invention a sulfonamide group may be introduced into the meta or para Ar position. This can be accomplished via several amenable synthetic multi step manipulations including the reaction of ammonia with sulfonyl chlorides (**Scheme 13A**) or alternatively sulfonamides can be obtained by reacting sulfinic acid salts with an electrophilic nitrogen source such as hydroxylamine-O-sulfonic acid or bis-(2,2,2-trichloroethyl)-azodicarboxylate.

15


Preferably 3-methoxy-3-oxapropane-1-sulfinate can serve as a sulfinate donor moiety through a simple alkylation and be subsequently removed via a beta-elimination reaction. Reaction of the resulting sulfinate with an electrophilic nitrogen source provides the primary sulfonamide analogue of the current invention. Such intermediates may be optionally further modified to amides such as those represented by general formula **49**. Acylsulfonamides of this type can be obtained by an

20

amidation reaction using an acid halide or anhydride (such as, $R_g\text{COCl}$ or $(R_g\text{CO})_2\text{O}$) and a base in an inert solvent (**Scheme 13C**). The base includes an alkali metal carbonate (such as, sodium carbonate, potassium carbonate, and the like), an alkali metal hydrogencarbonate (such as, sodium hydrogencarbonate, potassium hydrogencarbonate, and the like), an alkali hydroxide (such as, sodium

hydroxide or potassium hydroxide, and like), a tertiary amine (such as, *N,N*-diisopropylethylamine, triethylamine, *N*-methylmorpholine, and the like), or an aromatic amine (such as, pyridine, imidazole, poly-(4-vinylpyridine), and the like). The inert solvent includes lower halocarbon solvents (such as, dichloromethane, dichloroethane, chloroform, and the like), ethereal solvents (such as, tetrahydrofuran, dioxane, and the like), amide solvents (such as, *N,N*-dimethylacetamide, *N,N*-dimethylformamide, and the like), aromatic solvents (benzene, toluene, pyridine, and the like) and mixtures thereof. Reaction temperature ranges from about -20 °C to 50 °C, preferably about 0 °C to 40 °C.

Scheme 13



The compounds of the present invention may be prepared according to the general synthetic schemes as described herein as well as relevant published literature procedures that are used by one skilled in the art. Exemplary reagents and procedures for these reactions appear hereinafter in the working Examples. Protection and deprotection may be carried out by procedures generally known in the art (see, for example, Greene, T. W. and Wuts, P. G. M., *Protecting Groups in Organic Synthesis*, 3rd Edition, 1999 [Wiley]; incorporated herein by reference in its entirety).

The present invention also encompasses diastereomers as well as optical isomers, e.g. mixtures of enantiomers including racemic mixtures, as well as individual enantiomers and diastereomers, which arise as a consequence of structural asymmetry in certain compounds of Formula (I). Separation of the individual isomers or selective synthesis of the individual isomers is accomplished by application of various methods which are well known to practitioners in the art.

The present invention also encompasses diastereomers as well as optical isomers, e.g. mixtures of enantiomers including racemic mixtures, as well as individual enantiomers and diastereomers, which arise as a consequence of structural asymmetry in certain compounds of Formula (I). Separation of the individual isomers or selective synthesis of the individual isomers is accomplished by application of various methods which are well known to practitioners in the art.

INDICATIONS AND METHODS OF PROPHYLAXIS AND/OR TREATMENT

In addition to the foregoing beneficial uses for compounds of the present invention disclosed herein, compounds of the invention are useful in the treatment of additional diseases. Without limitation, these include the following.

The most significant pathologies in Type II diabetes are impaired insulin signaling at its target tissues ("insulin resistance") and failure of the insulin-producing cells of the pancreas to secrete an appropriate degree of insulin in response to a hyperglycemic signal. Current therapies to treat the latter include inhibitors of the β -cell ATP-sensitive potassium channel to trigger the release of endogenous insulin stores, or administration of exogenous insulin. Neither of these achieves accurate normalization of blood glucose levels and both carry the risk of inducing hypoglycemia. For these reasons, there has been intense interest in the development of pharmaceuticals that function in a glucose-dependent action, i.e. potentiators of glucose signaling. Physiological signaling systems which function in this manner are well-characterized and include the gut peptides GLP1, GIP and PACAP. These hormones act via their cognate G-protein coupled receptor to stimulate the production of cAMP in pancreatic β -cells. The increased cAMP does not appear to result in stimulation of insulin release during the fasting or preprandial state. However, a series of biochemical targets of cAMP signaling, including the ATP-sensitive potassium channel, voltage-sensitive potassium channels and the exocytotic machinery, are modified in such a way that the insulin secretory response to a postprandial glucose stimulus is markedly enhanced. Accordingly, agonists of novel, similarly functioning, β -cell GPCRs, including **RUP3**, would also stimulate the release of endogenous insulin and consequently promote normoglycemia in Type II diabetes.

It is also established that increased cAMP, for example as a result of GLP1 stimulation, promotes β -cell proliferation, inhibits β -cell death and thus improves islet mass. This positive effect on β -cell mass is expected to be beneficial in both Type II diabetes, where insufficient insulin is produced, and Type I diabetes, where β -cells are destroyed by an inappropriate autoimmune response.

Some β -cell GPCRs, including **RUP3**, are also present in the hypothalamus where they modulate hunger, satiety, decrease food intake, controlling or decreasing weight and energy expenditure. Hence, given their function within the hypothalamic circuitry, agonists or inverse agonists of these receptors mitigate hunger, promote satiety and therefore modulate weight.

It is also well-established that metabolic diseases exert a negative influence on other physiological systems. Thus, there is often the codevelopment of multiple disease states (e.g. type I diabetes, type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia, obesity or cardiovascular disease in "Syndrome X") or secondary diseases which clearly occur secondary to diabetes (e.g. kidney disease, peripheral neuropathy). Thus, it is expected that effective treatment of the diabetic condition will in turn be of benefit to such interconnected disease states.

In some embodiments of the present invention the metabolic-related disorder is hyperlipidemia, type 1 diabetes, type 2 diabetes mellitus, idiopathic type 1 diabetes (Type 1b), latent autoimmune diabetes in adults (LADA), early-onset type 2 diabetes (EOD), youth-onset atypical diabetes (YOAD), maturity onset diabetes of the young (MODY), malnutrition-related diabetes, gestational diabetes, coronary heart disease, ischemic stroke, restenosis after angioplasty, peripheral

vascular disease, intermittent claudication, myocardial infarction (e.g. necrosis and apoptosis), dyslipidemia, post-prandial lipemia, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose, metabolic acidosis, ketosis, arthritis, obesity, osteoporosis, hypertension, congestive heart failure, left ventricular hypertrophy, peripheral arterial disease, diabetic retinopathy, macular degeneration, cataract, diabetic nephropathy, glomerulosclerosis, chronic renal failure, diabetic neuropathy, metabolic syndrome, syndrome X, premenstrual syndrome, coronary heart disease, angina pectoris, thrombosis, atherosclerosis, myocardial infarction, transient ischemic attacks, stroke, vascular restenosis, hyperglycemia, hyperinsulinemia, hyperlipidemia, hypertriglyceridemia, insulin resistance, impaired glucose metabolism, conditions of impaired glucose tolerance, conditions of impaired fasting plasma glucose, obesity, erectile dysfunction, skin and connective tissue disorders, foot ulcerations and ulcerative colitis, endothelial dysfunction and impaired vascular compliance.

One aspect of the present invention pertains to methods for treatment of a metabolic-related disorder in an individual comprising administering to the individual in need of such treatment a therapeutically effective amount of a compound as described herein or a pharmaceutical composition thereof. In some embodiments the metabolic-related disorder is type I diabetes, type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia or syndrome X. In some embodiments the metabolic-related disorder is type II diabetes. In some embodiments the metabolic-related disorder is hyperglycemia. In some embodiments the metabolic-related disorder is hyperlipidemia. In some embodiments the metabolic-related disorder is hypertriglyceridemia. In some embodiments the metabolic-related disorder is type I diabetes. In some embodiments the metabolic-related disorder is dyslipidemia. In some embodiments the metabolic-related disorder is syndrome X. In some embodiments the individual is a mammal. In some embodiments the mammal is a human.

One aspect of the present invention pertains to methods of decreasing food intake of an individual comprising administering to the individual in need thereof a therapeutically effective amount of a compound of the present invention or pharmaceutical composition thereof. In some embodiments the individual is a mammal. In some embodiments the mammal is a human.

One aspect of the present invention pertains to methods of inducing satiety in an individual comprising administering to the individual in need of such treatment a therapeutically effective amount of a compound of the present invention or pharmaceutical composition thereof. In some embodiments the individual is a mammal. In some embodiments the mammal is a human.

One aspect of the present invention pertains to methods of controlling or decreasing weight gain of an individual comprising administering to the individual in need of such treatment a therapeutically effective amount of a compound of the present invention or pharmaceutical composition thereof. In some embodiments the individual is a mammal. In some embodiments the mammal is a human.

Some embodiments of the present invention pertain to methods wherein the human has a body mass index of about 18.5 to about 45. In some embodiments, the human has a body mass index of about 25 to about 45. In some embodiments, the human has a body mass index of about 30 to about 45. In some embodiments, the human has a body mass index of about 35 to about 45.

5 One aspect of the present invention pertains to methods of modulating a **RUP3** receptor in an individual comprising contacting the receptor with a compound of the present invention or pharmaceutical composition thereof. In some embodiments, the compound is an agonist. In some
10 embodiments, the compound is an inverse agonist. In some embodiments, the compound is an antagonist. In some embodiments, the modulation of the **RUP3** receptor is treatment of a metabolic-related disorder and complications thereof. In some embodiments, the metabolic-related disorder is
15 type I diabetes, type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia or syndrome X. In some
embodiments, the metabolic-related disorder is type II diabetes. In some embodiments, the
metabolic-related disorder is hyperglycemia. In some embodiments, the metabolic-related disorder is
hyperlipidemia. In some embodiments, the metabolic-related disorder is hypertriglyceridemia. In
some embodiments, the metabolic-related disorder is type I diabetes. In some embodiments, the
metabolic-related disorder is dyslipidemia. In some embodiments, the metabolic-related disorder is
syndrome X. In some embodiments, the individual is a mammal. In some embodiments, the mammal
is a human.

20 Some embodiments of the present invention include a method of modulating a **RUP3** receptor in an individual comprising contacting the receptor with a compound of the present invention wherein
the modulation of the **RUP3** receptor reduces food intake of the individual. In some embodiments the
individual is a mammal. In some embodiments the mammal is a human. In some embodiments the
human has a body mass index of about 18.5 to about 45. In some embodiments the human has a body
25 mass index of about 25 to about 45. In some embodiments the human has a body mass index of about
30 to about 45. In some embodiments the human has a body mass index of about 35 to about 45.

Some embodiments of the present invention include a method of modulating a **RUP3** receptor in an individual comprising contacting the receptor with a compound of the present invention wherein
the modulation of the **RUP3** receptor induces satiety in the individual. In some embodiments the
30 individual is a mammal. In some embodiments the mammal is a human. In some embodiments the
human has a body mass index of about 18.5 to about 45. In some embodiments the human has a body
mass index of about 25 to about 45. In some embodiments the human has a body mass index of about
30 to about 45. In some embodiments the human has a body mass index of about 35 to about 45.

Some embodiments of the present invention include a method of modulating a **RUP3** receptor
35 in an individual comprising contacting the receptor with a compound of the present invention wherein
the modulation of the **RUP3** receptor controls or reduces weight gain of the individual. In some
embodiments the individual is a mammal. In some embodiments the mammal is a human. In some

embodiments the human has a body mass index of about 18.5 to about 45. In some embodiments the human has a body mass index of about 25 to about 45. In some embodiments the human has a body mass index of about 30 to about 45. In some embodiments the human has a body mass index of about 35 to about 45.

5 One aspect of the present invention pertains to use of a compound as described herein, for production of a medicament for use in treatment of a metabolic-related disorder. In some embodiments, the metabolic-related disorder is type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia or syndrome X.

10 One aspect of the present invention pertains to use of a compound as described herein, for production of a medicament for use in decreasing food intake of an individual. In some embodiments, the individual is a mammal. In some embodiments, the mammal is a human. In some embodiments, the human has a body mass index of about 18.5 to about 45. In some embodiments, the human has a body mass index of about 25 to about 45. In some embodiments, the human has a body mass index of about 30 to about 45. In some embodiments, the human has a body mass index of about 35 to about 45.

15 One aspect of the present invention pertains to use of a compound as described herein, for production of a medicament for use of inducing satiety in an individual. In some embodiments, the individual is a mammal. In some embodiments, the mammal is a human. In some embodiments, the human has a body mass index of about 18.5 to about 45. In some embodiments, the human has a body mass index of about 25 to about 45. In some embodiments, the human has a body mass index of about 30 to about 45. In some embodiments, the human has a body mass index of about 35 to about 45.

20 One aspect of the present invention pertains to use of a compound as described herein, for production of a medicament for use in controlling or decreasing weight gain in an individual. In some embodiments, the individual is a mammal. In some embodiments, the mammal is a human. In some embodiments, the human has a body mass index of about 18.5 to about 45. In some embodiments, the human has a body mass index of about 25 to about 45. In some embodiments, the human has a body mass index of about 30 to about 45. In some embodiments, the human has a body mass index of about 35 to about 45.

25 One aspect of the present invention pertains to a compound, as described herein, for use in a method of treatment of the human or animal body by therapy.

 One aspect of the present invention pertains to a compound, as described herein, for use in a method of treatment of a metabolic-related disorder of the human or animal body by therapy.

35 One aspect of the present invention pertains to a compound, as described herein, for use in a method of decreasing food intake of the human or animal body by therapy.

One aspect of the present invention pertains to a compound, as described herein, for use in a method of inducing satiety of the human or animal body by therapy.

One aspect of the present invention pertains to a compound, as described herein, for use in a method of controlling or decreasing weight gain of the human or animal body by therapy.

5

PHARMACEUTICAL COMPOSITIONS

A further aspect of the present invention pertains to pharmaceutical compositions comprising one or more compounds of Formula (I) or any formula disclosed herein, and one or more pharmaceutically acceptable carriers. Some embodiments of the present invention pertain to pharmaceutical compositions comprising a compound of Formula (I) and a pharmaceutically acceptable carrier.

Some embodiments of the present invention include a method of producing a pharmaceutical composition comprising admixing at least one compound according to any of the compound embodiments disclosed herein and a pharmaceutically acceptable carrier.

Formulations may be prepared by any suitable method, typically by uniformly mixing the active compound(s) with liquids or finely divided solid carriers, or both, in the required proportions, and then, if necessary, forming the resulting mixture into a desired shape.

Conventional excipients, such as binding agents, fillers, acceptable wetting agents, tableting lubricants, and disintegrants may be used in tablets and capsules for oral administration. Liquid preparations for oral administration may be in the form of solutions, emulsions, aqueous or oily suspensions, and syrups. Alternatively, the oral preparations may be in the form of dry powder that can be reconstituted with water or another suitable liquid vehicle before use. Additional additives such as suspending or emulsifying agents, non-aqueous vehicles (including edible oils), preservatives, and flavorings and colorants may be added to the liquid preparations. Parenteral dosage forms may be prepared by dissolving the compound of the invention in a suitable liquid vehicle and filter sterilizing the solution before filling and sealing an appropriate vial or ampoule. These are just a few examples of the many appropriate methods well known in the art for preparing dosage forms.

A compound of the present invention can be formulated into pharmaceutical compositions using techniques well known to those in the art. Suitable pharmaceutically-acceptable carriers, outside those mentioned herein, are known in the art; for example, see Remington, The Science and Practice of Pharmacy, 20th Edition, 2000, Lippincott Williams & Wilkins, (Editors: Gennaro, A. R., et al.).

While it is possible that, for use in the prophylaxis or treatment, a compound of the invention may, in an alternative use, be administered as a raw or pure chemical, it is preferable however to present the compound or active ingredient as a pharmaceutical formulation or composition further comprising a pharmaceutically acceptable carrier.

The invention thus further provides pharmaceutical formulations comprising a compound of the invention or a pharmaceutically acceptable salt or derivative thereof together with one or more pharmaceutically acceptable carriers thereof and/or prophylactic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not
5 overly deleterious to the recipient thereof.

Pharmaceutical formulations include those suitable for oral, rectal, nasal, topical (including buccal and sub-lingual), vaginal or parenteral (including intramuscular, sub-cutaneous and intravenous) administration or in a form suitable for administration by inhalation, insufflation or by a transdermal patch. Transdermal patches dispense a drug at a controlled rate by presenting the drug for
10 absorption in an efficient manner with a minimum of degradation of the drug. Typically, transdermal patches comprise an impermeable backing layer, a single pressure sensitive adhesive and a removable protective layer with a release liner. One of ordinary skill in the art will understand and appreciate the techniques appropriate for manufacturing a desired efficacious transdermal patch based upon the needs of the artisan.

The compounds of the invention, together with a conventional adjuvant, carrier, or diluent, may thus be placed into the form of pharmaceutical formulations and unit dosages thereof, and in such form may be employed as solids, such as tablets or filled capsules, or liquids such as solutions, suspensions, emulsions, elixirs, gels or capsules filled with the same, all for oral use, in the form of suppositories for rectal administration; or in the form of sterile injectable solutions for parenteral
20 (including subcutaneous) use. Such pharmaceutical compositions and unit dosage forms thereof may comprise conventional ingredients in conventional proportions, with or without additional active compounds or principles, and such unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed.

For oral administration, the pharmaceutical composition may be in the form of, for example, a
25 tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are capsules, tablets, powders, granules or a suspension, with conventional additives such as lactose, mannitol, corn starch or potato starch; with binders such as crystalline cellulose, cellulose derivatives, acacia, corn starch or gelatins; with disintegrators such as corn starch, potato starch or sodium carboxymethyl-cellulose; and with lubricants such as talc or magnesium stearate. The active
30 ingredient may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable pharmaceutically acceptable carrier.

Compounds of the present invention or a solvate or physiologically functional derivative thereof can be used as active ingredients in pharmaceutical compositions, specifically as **RUP3**
35 receptor modulators. By the term "active ingredient" is defined in the context of a "pharmaceutical composition" and shall mean a component of a pharmaceutical composition that provides the primary

pharmacological effect, as opposed to an "inactive ingredient" which would generally be recognized as providing no pharmaceutical benefit.

5 The dose when using the compounds of the present invention can vary within wide limits, and as is customary and is known to the physician, it is to be tailored to the individual conditions in each individual case. It depends, for example, on the nature and severity of the illness to be treated, on the condition of the patient, on the compound employed or on whether an acute or chronic disease state is treated or prophylaxis is conducted or on whether further active compounds are administered in addition to the compounds of the present invention. Representative doses of the present invention include, but not limited to, about 0.001 mg to about 5000 mg, about 0.001 to about 2500 mg, about 10 0.001 to about 1000 mg, 0.001 to about 500 mg, 0.001 mg to about 250 mg, about 0.001 mg to 100 mg, about 0.001 mg to about 50 mg, and about 0.001 mg to about 25 mg. Multiple doses may be administered during the day, especially when relatively large amounts are deemed to be needed, for example 2, 3 or 4, doses. Depending on the individual and as deemed appropriate from the patient's physician or care-giver it may be necessary to deviate upward or downward from the doses described
15 herein.

The amount of active ingredient, or an active salt or derivative thereof, required for use in treatment will vary not only with the particular salt selected but also with the route of administration, the nature of the condition being treated and the age and condition of the patient and will ultimately be at the discretion of the attendant physician or clinician. In general, one skilled in the art
20 understands how to extrapolate in vivo data obtained in a model system, typically an animal model, to another, such as a human. Typically, animal models include, but are not limited to, the rodents diabetes models as described in Example 5, *infra* (as well as other animal models known in the art, such as those reported by Reed and Scribner in Diabetes, Obesity and Metabolism, 1, 1999, 75-86). In some circumstances, these extrapolations may merely be based on the weight of the animal model
25 in comparison to another, such as a mammal, preferably a human, however, more often, these extrapolations are not simply based on weights, but rather incorporate a variety of factors. Representative factors include the type, age, weight, sex, diet and medical condition of the patient, the severity of the disease, the route of administration, pharmacological considerations such as the activity, efficacy, pharmacokinetic and toxicology profiles of the particular compound employed,
30 whether a drug delivery system is utilized, on whether an acute or chronic disease state is being treated or prophylaxis is conducted or on whether further active compounds are administered in addition to the compounds of the Formula (I) and as part of a drug combination. The dosage regimen for treating a disease condition with the compounds and/or compositions of this invention is selected in accordance with a variety factors as cited above. Thus, the actual dosage regimen employed may
35 vary widely and therefore may deviate from a preferred dosage regimen and one skilled in the art will recognize that dosage and dosage regimen outside these typical ranges can be tested and, where appropriate, may be used in the methods of this invention.

The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example, as two, three, four or more sub-doses per day. The sub-dose itself may be further divided, e.g., into a number of discrete loosely spaced administrations. The daily dose can be divided, especially when relatively large amounts are administered as deemed
5 appropriate, into several, for example 2, 3 or 4, part administrations. If appropriate, depending on individual behavior, it may be necessary to deviate upward or downward from the daily dose indicated.

The compounds of the present invention can be administered in a wide variety of oral and parenteral dosage forms. It will be obvious to those skilled in the art that the following dosage forms
10 may comprise, as the active component, either a compound of the invention or a pharmaceutically acceptable salt of a compound of the invention.

For preparing pharmaceutical compositions from the compounds of the present invention, the selection of a suitable pharmaceutically acceptable carrier can be either solid, liquid or a mixture of both. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and
15 dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavouring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component.

20 In tablets, the active component is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted to the desired shape and size.

The powders and tablets may contain varying percentage amounts of the active compound. A representative amount in a powder or tablet may contain from 0.5 to about 90 percent of the active compound; however, an artisan would know when amounts outside of this range are necessary.
25 Suitable carriers for powders and tablets are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as carrier providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is thus in
30 association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid forms suitable for oral administration.

For preparing suppositories, a low melting wax, such as an admixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to
35 cool, and thereby to solidify.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

5 Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water-propylene glycol solutions. For example, parenteral injection liquid preparations can be formulated as solutions in aqueous polyethylene glycol solution. Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that 10 may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

15 The compounds according to the present invention may thus be formulated for parenteral administration (e.g. by injection, for example bolus injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in multi-dose containers with an added preservative. The pharmaceutical compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be 20 in powder form, obtained by aseptic isolation of sterile solid or by lyophilization from solution, for constitution with a suitable vehicle, e.g. sterile, pyrogen-free water, before use.

Aqueous formulations suitable for oral use can be prepared by dissolving or suspending the active component in water and adding suitable colorants, flavours, stabilizing and thickening agents, 25 as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, or other well known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before 30 use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

For topical administration to the epidermis the compounds according to the invention may be 35 formulated as ointments, creams or lotions, or as a transdermal patch.

Ointments and creams may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or

oily base and will in general also contain one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents.

Formulations suitable for topical administration in the mouth include lozenges comprising active agent in a flavored base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert base such as gelatin and glycerin or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Solutions or suspensions are applied directly to the nasal cavity by conventional means, for example with a dropper, pipette or spray. The formulations may be provided in single or multi-dose form. In the latter case of a dropper or pipette, this may be achieved by the patient administering an appropriate, predetermined volume of the solution or suspension. In the case of a spray, this may be achieved for example by means of a metering atomizing spray pump.

Administration to the respiratory tract may also be achieved by means of an aerosol formulation in which the active ingredient is provided in a pressurized pack with a suitable propellant. If the compounds of the Formula (I) or pharmaceutical compositions comprising them are administered as aerosols, for example as nasal aerosols or by inhalation, this can be carried out, for example, using a spray, a nebulizer, a pump nebulizer, an inhalation apparatus, a metered inhaler or a dry powder inhaler. Pharmaceutical forms for administration of the compounds of the Formula (I) as an aerosol can be prepared by processes well-known to the person skilled in the art. For their preparation, for example, solutions or dispersions of the compounds of the Formula (I) in water, water/alcohol mixtures or suitable saline solutions can be employed using customary additives, for example benzyl alcohol or other suitable preservatives, absorption enhancers for increasing the bioavailability, solubilizers, dispersants and others, and, if appropriate, customary propellants, for example include carbon dioxide, CFC's, such as, dichlorodifluoromethane, trichlorofluoromethane, or dichlorotetrafluoroethane; and the like. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by provision of a metered valve.

In formulations intended for administration to the respiratory tract, including intranasal formulations, the compound will generally have a small particle size for example of the order of 10 microns or less. Such a particle size may be obtained by means known in the art, for example by micronization. When desired, formulations adapted to give sustained release of the active ingredient may be employed.

Alternatively the active ingredients may be provided in the form of a dry powder, for example, a powder mix of the compound in a suitable powder base such as lactose, starch, starch derivatives such as hydroxypropylmethyl cellulose and polyvinylpyrrolidone (PVP). Conveniently the powder carrier will form a gel in the nasal cavity. The powder composition may be presented in unit dose form for example in capsules or cartridges of, e.g., gelatin, or blister packs from which the powder may be administered by means of an inhaler.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

Tablets or capsules for oral administration and liquids for intravenous administration are preferred compositions.

The compounds according to the invention may optionally exist as pharmaceutically acceptable salts including pharmaceutically acceptable acid addition salts prepared from pharmaceutically acceptable non-toxic acids including inorganic and organic acids. Representative acids include, but are not limited to, acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethenesulfonic, dichloroacetic, formic, fumaric, gluconic, glutamic, hippuric, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, oxalic, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, oxalic, p-toluenesulfonic and the like, such as those pharmaceutically acceptable salts listed in *Journal of Pharmaceutical Science*, 66, 2 (1977); incorporated herein by reference in its entirety.

The acid addition salts may be obtained as the direct products of compound synthesis. In the alternative, the free base may be dissolved in a suitable solvent containing the appropriate acid, and the salt isolated by evaporating the solvent or otherwise separating the salt and solvent. The compounds of this invention may form solvates with standard low molecular weight solvents using methods known to the skilled artisan.

Compounds of the present invention can be converted to "pro-drugs." The term "pro-drugs" refers to compounds that have been modified with specific chemical groups known in the art and when administered into an individual these groups undergo biotransformation to give the parent compound. Pro-drugs can thus be viewed as compounds of the invention containing one or more specialized non-toxic protective groups used in a transient manner to alter or to eliminate a property of the compound. In one general aspect, the "pro-drug" approach is utilized to facilitate oral absorption. A thorough discussion is provided in T. Higuchi and V. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S. Symposium Series; and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987, both of which are hereby incorporated by reference in their entirety.

Some embodiments of the present invention include a method of producing a pharmaceutical composition for "combination-therapy" comprising admixing at least one compound according to any of the compound embodiments disclosed herein, together with at least one known pharmaceutical agent as described herein and a pharmaceutically acceptable carrier.

In some embodiments the pharmaceutical agent is selected from the group consisting of: apolipoprotein-B secretion/microsomal triglyceride transfer protein (apo-B/MTP) inhibitors, MCR-4 agonists, cholecystokinin-A (CCK-A) agonists, serotonin and norepinephrine reuptake inhibitors (for example, sibutramine), sympathomimetic agents, β_3 adrenergic receptor agonists, dopamine agonists (for example, bromocriptine), melanocyte-stimulating hormone receptor analogs, cannabinoid 1 receptor antagonists [for example, SR141716: *N*-(piperidin-1-yl)-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide], melanin concentrating hormone antagonists, leptons (the OB protein), leptin analogues, leptin receptor agonists, galanin antagonists, lipase inhibitors (such as tetrahydrolipstatin, i.e., Orlistat), anorectic agents (such as a bombesin agonist),
 5 Neuropeptide-Y antagonists, thyromimetic agents, dehydroepiandrosterone or an analogue thereof, glucocorticoid receptor agonists or antagonists, orexin receptor antagonists, urocortin binding protein antagonists, glucagon-like peptide-1 receptor agonists, ciliary neurotrophic factors (such as Axokine™), human agouti-related proteins (AGRP), ghrelin receptor antagonists, histamine 3 receptor antagonists or reverse agonists, neuromedin U receptor agonists, noradrenergic anorectic agents (for
 10 example, phentermine, mazindol and the like), appetite suppressants (for example, bupropion) and the like. In further embodiments, the pharmaceutical agent is selected from the group consisting of orlistat, sibutramine, bromocriptine, ephedrine, leptin, and pseudoephedrine.

In some embodiments the pharmaceutical agent is selected from the group consisting of: sulfonylureas, meglitinides, biguanides, α -glucosidase inhibitors, peroxisome proliferators-activated
 20 receptor- γ (i.e., PPAR- γ) agonists, insulin, insulin analogues, HMG-CoA reductase inhibitors, cholesterol-lowering drugs (for example, fibrates that include: fenofibrate, bezafibrate, gemfibrozil, clofibrate and the like; bile acid sequestrants which include: cholestyramine, colestipol and the like; and niacin), antiplatelet agents (for example, aspirin and adenosine diphosphate receptor antagonists that include: clopidogrel, ticlopidine and the like), angiotensin-converting enzyme inhibitors,
 25 angiotensin II receptor antagonists and adiponectin.

It is noted that when the **RUP3** receptor modulators are utilized as active ingredients in a pharmaceutical composition, these are not intended for use only in humans, but in other non-human mammals as well. Indeed, recent advances in the area of animal health-care mandate that
 30 consideration be given for the use of active agents, such as **RUP3** receptor modulators, for the treatment of obesity in domestic animals (*e.g.*, cats and dogs), and **RUP3** receptor modulators in other domestic animals where no disease or disorder is evident (*e.g.*, food-oriented animals such as cows, chickens, fish, etc.). Those of ordinary skill in the art are readily credited with understanding the utility of such compounds in such settings.

35 COMBINATION THERAPY - PROPHYLAXIS AND TREATMENT

In the context of the present invention, a compound of Formula (I) or pharmaceutical composition thereof can be utilized for modulating the activity of **RUP3** receptor mediated diseases,

conditions and/or disorders as described herein. Examples of modulating the activity of **RUP3** receptor mediated diseases include the prophylaxis or treatment of metabolic related disorders such as, but not limited to, type I diabetes, type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia and syndrome X. Other examples of modulating the activity of **RUP3** receptor mediated diseases include the prophylaxis or treatment of obesity and/or overweight by decreasing food intake, inducing satiation (i.e., the feeling of fullness), controlling weight gain, decreasing body weight and/or affecting metabolism such that the recipient loses weight and/or maintains weight.

While the compounds of the invention can be administered as the sole active pharmaceutical agent (i.e., mono-therapy), they can also be used in combination with other pharmaceutical agents (i.e., combination-therapy) for the treatment of the diseases/conditions/disorders described herein. Therefore, another aspect of the present invention includes methods of prophylaxis and/or treatment of a metabolic related disorder or a weight related disorder, such as obesity, comprising administering to an individual in need of prophylaxis and/or treatment a therapeutically effective amount of a compound of the present invention, for example Formula (I), in combination with one or more additional pharmaceutical agent as described herein.

Suitable pharmaceutical agents that can be used in combination with the compounds of the present invention include anti-obesity agents such as apolipoprotein-B secretion/microsomal triglyceride transfer protein (apo-B/MTP) inhibitors, MCR-4 agonists, cholecystokinin-A (CCK-A) agonists, serotonin and norepinephrine reuptake inhibitors (for example, sibutramine), sympathomimetic agents, $\beta 3$ adrenergic receptor agonists, dopamine agonists (for example, bromocriptine), melanocyte-stimulating hormone receptor analogs, cannabinoid 1 receptor antagonists [for example, SR141716: N-(piperidin-1-yl)-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazole-3-carboxamide], melanin concentrating hormone antagonists, leptons (the OB protein), leptin analogues, leptin receptor agonists, galanin antagonists, lipase inhibitors (such as tetrahydrolipstatin, i.e., Orlistat), anorectic agents (such as a bombesin agonist), Neuropeptide-Y antagonists, thyromimetic agents, dehydroepiandrosterone or an analogue thereof, glucocorticoid receptor agonists or antagonists, orexin receptor antagonists, urocortin binding protein antagonists, glucagon-like peptide-1 receptor agonists, ciliary neurotrophic factors (such as Axokine™ available from Regeneron Pharmaceuticals, Inc., Tarrytown, NY and Procter & Gamble Company, Cincinnati, OH), human agouti-related proteins (AGRP), ghrelin receptor antagonists, histamine 3 receptor antagonists or reverse agonists, neuromedin U receptor agonists, noradrenergic anorectic agents (for example, phentermine, mazindol and the like) and appetite suppressants (for example, bupropion).

Other anti-obesity agents, including the agents set forth *infra*, are well known, or will be readily apparent in light of the instant disclosure, to one of ordinary skill in the art.

In some embodiments, the anti-obesity agents are selected from the group consisting of orlistat, sibutramine, bromocriptine, ephedrine, leptin, and pseudoephedrine. In a further

embodiment, compounds of the present invention and combination therapies are administered in conjunction with exercise and/or a sensible diet.

It will be understood that the scope of combination-therapy of the compounds of the present invention with other anti-obesity agents, anorectic agents, appetite suppressant and related agents is not limited to those listed above, but includes in principle any combination with any pharmaceutical agent or pharmaceutical composition useful for the treatment of overweight and obese individuals.

Other suitable pharmaceutical agents, in addition to anti-obesity agents, that can be used in combination with the compounds of the present invention include agents useful in the treatment of metabolic related disorders and/or concomitant diseases thereof. For example, but not limited to, congestive heart failure, type I diabetes, type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia, syndrome X, retinopathy, nephropathy and neuropathy. Prophylaxis or treatment of one or more of the diseases cited herein include the use of one or more pharmaceutical agents known in the art belonging to the classes of drugs referred to, but not limited to, the following: sulfonylureas, meglitinides, biguanides, α -glucosidase inhibitors, peroxisome proliferators-activated receptor- γ (i.e., PPAR- γ) agonists, insulin, insulin analogues, HMG-CoA reductase inhibitors, cholesterol-lowering drugs (for example, fibrates that include: fenofibrate, bezafibrate, gemfibrozil, clofibrate and the like; bile acid sequestrants which include: cholestyramine, colestipol and the like; and niacin), antiplatelet agents (for example, aspirin and adenosine diphosphate receptor antagonists that include: clopidogrel, ticlopidine and the like), angiotensin-converting enzyme inhibitors, angiotensin II receptor antagonists, adiponectin and the like. In accordance to one aspect of the present invention, a compound of the present can be used in combination with a pharmaceutical agent or agents belonging to one or more of the classes of drugs cited herein.

It will be understood that the scope of combination-therapy of the compounds of the present invention with other pharmaceutical agents is not limited to those listed herein, *supra* or *infra*, but includes in principle any combination with any pharmaceutical agent or pharmaceutical composition useful for the prophylaxis or treatment of diseases, conditions or disorders that are linked to metabolic related disorders.

Some embodiments of the present invention include methods of prophylaxis or treatment of a disease, disorder, condition or complication thereof as described herein, comprising administering to an individual in need of such prophylaxis or treatment a therapeutically effective amount or dose of a compound of the present invention in combination with at least one pharmaceutical agent selected from the group consisting of: sulfonylureas, meglitinides, biguanides, α -glucosidase inhibitors, peroxisome proliferators-activated receptor- γ (i.e., PPAR- γ) agonists, insulin, insulin analogues, HMG-CoA reductase inhibitors, cholesterol-lowering drugs (for example, fibrates that include: fenofibrate, bezafibrate, gemfibrozil, clofibrate and the like; bile acid sequestrants which include: cholestyramine, colestipol and the like; and niacin), antiplatelet agents (for example, aspirin and

adenosine diphosphate receptor antagonists that include: clopidogrel, ticlopidine and the like), angiotensin-converting enzyme inhibitors, angiotensin II receptor antagonists and adiponectin. In some embodiments, methods of the present invention include compounds of the present invention and the pharmaceutical agents are administered separately. In further embodiments, compounds of the present invention and the pharmaceutical agents are administered together.

Suitable pharmaceutical agents that can be used in conjunction with compounds of the present invention include sulfonylureas. The sulfonylureas (SU) are drugs which promote secretion of insulin from pancreatic β cells by transmitting signals of insulin secretion via SU receptors in the cell membranes. Examples of the sulfonylureas include glyburide, glipizide, glimepiride and other sulfonylureas known in the art.

Suitable pharmaceutical agents that can be used in conjunction with compounds of the present invention include the meglitinides. The meglitinides are benzoic acid derivatives represent a novel class of insulin secretagogues. These agents target postprandial hyperglycemia and show comparable efficacy to sulfonylureas in reducing HbA1c. Examples of meglitinides include repaglinide, nateglinide and other meglitinides known in the art.

Suitable pharmaceutical agents that can be used in conjunction with compounds of the present invention include the biguanides. The biguanides represent a class of drugs that stimulate anaerobic glycolysis, increase the sensitivity to insulin in the peripheral tissues, inhibit glucose absorption from the intestine, suppress of hepatic gluconeogenesis, and inhibit fatty acid oxidation. Examples of biguanides include phenformin, metformin, buformin, and biguanides known in the art.

Suitable pharmaceutical agents that can be used in conjunction with compounds of the present invention include the α -glucosidase inhibitors. The α -glucosidase inhibitors competitively inhibit digestive enzymes such as α -amylase, maltase, α -dextrinase, sucrase, etc. in the pancreas and or small intestine. The reversible inhibition by α -glucosidase inhibitors retard, diminish or otherwise reduce blood glucose levels by delaying the digestion of starch and sugars. Examples of α -glucosidase inhibitors include acarbose, N-(1,3-dihydroxy-2-propyl)valiolamine (generic name; voglibose), miglitol, and α -glucosidase inhibitors known in the art.

Suitable pharmaceutical agents that can be used in conjunction with compounds of the present invention include the peroxisome proliferators-activated receptor- γ (i.e., PPAR- γ) agonists. The peroxisome proliferators-activated receptor- γ agonists represent a class of compounds that activates the nuclear receptor PPAR- γ and therefore regulate the transcription of insulin-responsive genes involved in the control of glucose production, transport and utilization. Agents in the class also facilitate the regulation of fatty acid metabolism. Examples of PPAR- γ agonists include rosiglitazone, pioglitazone, tesaglitazar, netoglitazone, GW-409544, GW-501516 and PPAR- γ agonists known in the art.

Suitable pharmaceutical agents that can be used in conjunction with compounds of the present invention include the HMG-CoA reductase inhibitors. The HMG-CoA reductase inhibitors are agents also referred to as Statin compounds that belong to a class of drugs that lower blood cholesterol levels by inhibiting hydroxymethylglutaryl CoA (HMG-CoA) reductase. HMG-CoA reductase is the rate-limiting enzyme in cholesterol biosynthesis. The statins lower serum LDL concentrations by upregulating the activity of LDL receptors and are responsible for clearing LDL from the blood. Some representative examples the statin compounds include rosuvastatin, pravastatin and its sodium salt, simvastatin, lovastatin, atorvastatin, fluvastatin, cerivastatin, rosuvastatin, pitavastatin, BMS's "superstatin", and HMG-CoA reductase inhibitors known in the art.

Suitable pharmaceutical agents that can be used in conjunction with compounds of the present invention include the Fibrates. Fibrate compounds belong to a class of drugs that lower blood cholesterol levels by inhibiting synthesis and secretion of triglycerides in the liver and activating a lipoprotein lipase. Fibrates have been known to activate peroxisome proliferators-activated receptors and induce lipoprotein lipase expression. Examples of fibrate compounds include bezafibrate, beclobrate, binifibrate, ciprofibrate, clonofibrate, clofibrate, clofibric acid, etofibrate, fenofibrate, gemfibrozil, nicofibrate, pirifibrate, ronifibrate, simfibrate, theofibrate, and fibrates known in the art.

Suitable pharmaceutical agents that can be used in conjunction with compounds of the present invention include the angiotensin converting enzyme (ACE) inhibitors. The angiotensin converting enzyme inhibitors belong to the class of drugs that partially lower blood glucose levels as well as lowering blood pressure by inhibiting angiotensin converting enzymes. Examples of the angiotensin converting enzyme inhibitors include captopril, enalapril, alacepril, delapril; ramipril, lisinopril, imidapril, benazepril, ceronapril, cilazapril, enalaprilat, fosinopril, moveltopril, perindopril, quinapril, spirapril, temocapril,trandolapril, and angiotensin converting enzyme inhibitors known in the art.

Suitable pharmaceutical agents that can be used in conjunction with compounds of the present invention include the angiotensin II receptor antagonists. Angiotensin II receptor antagonists target the angiotensin II receptor subtype 1 (i.e., AT1) and demonstrate a beneficial effect on hypertension. Examples of angiotensin II receptor antagonists include losartan (and the potassium salt form), and angiotensin II receptor antagonists known in the art.

Other treatments for one or more of the diseases cited herein include the use of pharmaceutical agents known in the art belonging to the classes of drugs referred to, but not limited to, the following: amylin agonists (for example, pramlintide), insulin secretagogues (for example, GLP-1 agonists; exendin-4; insulinotropin (NN2211); dipeptyl peptidase inhibitors (for example, NVP-DPP-728), acyl CoA cholesterol acetyltransferase inhibitors (for example, Ezetimibe, eflucimibe, and like compounds), cholesterol absorption inhibitors (for example, ezetimibe, pamaqueside and like compounds), cholesterol ester transfer protein inhibitors (for example, CP-529414, JTT-705, CETi-1, and like compounds), microsomal triglyceride transfer protein inhibitors (for example, implitapide, and like compounds), cholesterol modulators (for example, NO-1886, and

like compounds), bile acid modulators (for example, GT103-279 and like compounds) and squalene synthase inhibitors.

Squalene synthesis inhibitors belong to a class of drugs that lower blood cholesterol levels by inhibiting synthesis of squalene. Examples of the squalene synthesis inhibitors include (S)- α -
5 [Bis[2,2-dimethyl-1-oxopropoxy)methoxy] phosphinyl]-3-phenoxybenzenebutanesulfonic acid, mono potassium salt (BMS-188494) and squalene synthesis inhibitors known in the art.

In accordance with the present invention, the combination can be used by mixing the respective active components either all together or independently with a physiologically acceptable carrier, excipient, binder, diluent, etc., as described herein above, and administering the mixture or
10 mixtures either orally or non-orally as a pharmaceutical composition. When a compound or a mixture of compounds of Formula (I) are administered as a combination therapy with another active compound the therapeutic agents can be formulated as a separate pharmaceutical compositions given at the same time or at different times, or the therapeutic agents can be given as a single composition.

15 OTHER UTILITIES

Another object of the present invention relates to radio-labeled compounds of Formula (I) that would be useful not only in radio-imaging but also in assays, both in vitro and in vivo, for localizing and quantitating the **RUP3** receptor in tissue samples, including human, and for identifying **RUP3** receptor ligands by inhibition binding of a radio-labeled compound. It is a further object of this
20 invention to develop novel **RUP3** receptor assays of which comprise such radio-labeled compounds.

The present invention embraces isotopically-labeled compounds of Formula (I) and any subgenera herein, such as but not limited to, Formula (Ia) through Formula (Is). An "isotopically" or "radio-labeled" compounds are those which are identical to compounds disclosed herein, but for the fact that one or more atoms are replaced or substituted by an atom having an atomic mass or mass
25 number different from the atomic mass or mass number typically found in nature (i.e., naturally occurring). Suitable radionuclides that may be incorporated in compounds of the present invention include but are not limited to ^2H (also written as D for deuterium), ^3H (also written as T for tritium), ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{18}F , ^{35}S , ^{36}Cl , ^{82}Br , ^{75}Br , ^{76}Br , ^{77}Br , ^{123}I , ^{124}I , ^{125}I and ^{131}I . The radionuclide that is incorporated in the instant radio-labeled compounds will depend on the specific
30 application of that radio-labeled compound. For example, for in vitro **RUP3** receptor labeling and competition assays, compounds that incorporate ^3H , ^{14}C , ^{82}Br , ^{125}I , ^{131}I , ^{35}S or will generally be most useful. For radio-imaging applications ^{11}C , ^{18}F , ^{125}I , ^{123}I , ^{124}I , ^{131}I , ^{75}Br , ^{76}Br or ^{77}Br will generally be most useful.

It is understood that a "radio-labeled" or "labeled compound" is a compound of Formula (I)
35 that has incorporated at least one radionuclide; in some embodiments the radionuclide is selected from the group consisting of ^3H , ^{14}C , ^{125}I , ^{35}S and ^{82}Br .

Certain isotopically-labeled compounds of the present invention are useful in compound and/or substrate tissue distribution assays. In some embodiments the radionuclide ^3H and/or ^{14}C isotopes are useful in these studies. Further, substitution with heavier isotopes such as deuterium (i.e., ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability (e.g., increased in vivo half-life or reduced dosage requirements) and hence may be preferred in some circumstances.

Isotopically labeled compounds of the present invention can generally be prepared by following procedures analogous to those disclosed in the Schemes *supra* and Examples *infra*, by substituting an isotopically labeled reagent for a non-isotopically labeled reagent. Other synthetic methods that are useful are discussed *infra*. Moreover, it should be understood that all of the atoms represented in the compounds of the invention can be either the most commonly occurring isotope of such atoms or the more scarce radio-isotope or nonradio-active isotope.

Synthetic methods for incorporating radio-isotopes into organic compounds are applicable to compounds of the invention and are well known in the art. These synthetic methods, for example, incorporating activity levels of tritium into target molecules, are as follows:

- 15 A. Catalytic Reduction with Tritium Gas - This procedure normally yields high specific activity products and requires halogenated or unsaturated precursors.
- B. Reduction with Sodium Borohydride [^3H] - This procedure is rather inexpensive and requires precursors containing reducible functional groups such as aldehydes, ketones, lactones, esters, and the like.
- 20 C. Reduction with Lithium Aluminum Hydride [^3H] - This procedure offers products at almost theoretical specific activities. It also requires precursors containing reducible functional groups such as aldehydes, ketones, lactones, esters, and the like.
- D. Tritium Gas Exposure Labeling - This procedure involves exposing precursors containing exchangeable protons to tritium gas in the presence of a suitable catalyst.
- 25 E. N-Methylation using Methyl Iodide [^3H] - This procedure is usually employed to prepare O-methyl or N-methyl (^3H) products by treating appropriate precursors with high specific activity methyl iodide (^3H). This method in general allows for higher specific activity, such as for example, about 70-90 Ci/mmol.

Synthetic methods for incorporating activity levels of ^{125}I into target molecules include:

- 30 A. Sandmeyer and like reactions - This procedure transforms an aryl or heteroaryl amine into a diazonium salt, such as a tetrafluoroborate salt, and subsequently to ^{125}I labeled compound using Na^{125}I . A represented procedure was reported by Zhu, D.-G. and co-workers in J. Org. Chem. 2002, 67, 943-948.
- B. Ortho ^{125}I iodination of phenols - This procedure allows for the incorporation of ^{125}I at the ortho position of a phenol as reported by Collier, T. L. and co-workers in J. Labeled Compd Radiopharm. 1999, 42, S264-S266.
- 35

C. Aryl and heteroaryl bromide exchange with ^{125}I – This method is generally a two step process. The first step is the conversion of the aryl or heteroaryl bromide to the corresponding tri-alkyltin intermediate using for example, a Pd catalyzed reaction [i.e. $\text{Pd}(\text{Ph}_3\text{P})_4$] or through an aryl or heteroaryl lithium, in the presence of a tri-alkyltinhalide or hexaalkylditin [e.g., $(\text{CH}_3)_3\text{SnSn}(\text{CH}_3)_3$].

5 A represented procedure was reported by Bas, M.-D. and co-workers in J. Labeled Compd Radiopharm. 2001, 44, S280-S282.

A radio-labeled **RUP3** receptor compound of Formula (I) can be used in a screening assay to identify/evaluate compounds. In general terms, a newly synthesized or identified compound (i.e., test compound) can be evaluated for its ability to reduce binding of the “radio-labeled compound of Formula (I)” to the **RUP3** receptor. Accordingly, the ability of a test compound to compete with the “radio-labeled compound of Formula (I)” for the binding to the **RUP3** receptor directly correlates to its binding affinity.

The labeled compounds of the present invention bind to the **RUP3** receptor. In one embodiment the labeled compound has an IC_{50} less than about 500 μM , in another embodiment the labeled compound has an IC_{50} less than about 100 μM , in yet another embodiment the labeled compound has an IC_{50} less than about 10 μM , in yet another embodiment the labeled compound has an IC_{50} less than about 1 μM , and in still yet another embodiment the labeled inhibitor has an IC_{50} less than about 0.1 μM .

Other uses of the disclosed receptors and methods will become apparent to those in the art based upon, inter alia, a review of this disclosure.

As will be recognized, the steps of the methods of the present invention need not be performed any particular number of times or in any particular sequence. Additional objects, advantages, and novel features of this invention will become apparent to those skilled in the art upon examination of the following examples thereof, which are intended to be illustrative and not intended to be limiting.

EXAMPLES

The examples are provided to further define the invention without, however, limiting the invention to the specifics of these examples.

30

Example 1

96- well Cyclic AMP membrane assay for RUP3

Materials:

35 1) Adenyl cyclase Activation Flashplate Assay kit from Perkin Elmer -- 96 wells (SMP004B) and ^{125}I tracer (NEX130) which comes with the kit. Keep in refrigerator, in a box, and do not expose the Flashplates to light.

- 2) Phosphocreatine – Sigma P-7936
- 3) Creatine Phosphokinase --- Sigma C-3755
- 4) GTP – Sigma G-8877
- 5) ATP- Sigma A-2383
- 5 6) IBMX – Sigma I-7018
- 7) Hepes – 1M solution in distilled water – Gibco #15630080
- 8) MgCl₂ – Sigma M-1028- 1M Solution
- 9) NaCl – Sigma – S6546 – 5M Solution
- 10) Bradford Protein Assay Kit – Biorad # 5000001
- 10 11) Proclin 300- Sigma #4-8126

Binding Buffer - filter through 45- micron Nalgene filter and keep in refrigerator. All buffers and membranes should be kept cold (in ice bucket) while performing assay.

20 mM Hepes, pH7.4

- 15 1 mM MgCl₂
- 100 mM NaCl

2X Regeneration Buffer (make in binding buffer):

20 mM Phosphocreatine (1.02 gm/200 ml binding buffer)

- 20 20 units Creatine phosphokinase (4 mg/200 ml)

20 uM GTP (make up 10.46 mg/ml in binding buffer and add 200 µL /200 ml)

0.2 mM ATP (22.04 mg/200 ml)

100 mM IBMX (44.4 mg IBMX dissolved in 1 ml 100% DMSO first and then add the entire amount to 200 ml of buffer).

25

Regeneration buffer can be aliquotted into 40-45 ml portions (in 50 ml sterile tubes) and kept frozen for up to 2 months. Simply put the tube in a beaker with room temperature water to thaw out the regeneration buffer on the day of the assay.

30

A. Assay procedure

- 1) Pipet 50 µL regeneration buffer in all 96 wells using Matrix 1250 8-channel pipettor.

- 2) Pipet 5 µL DMSO in columns 1 and columns 11 and 12.

35

- 3) Pipet 50 µL cAMP standards in columns 11 and 12 in this format: 50 pmole/well for row A, 25 pmole/well for row B, 12.5 pmol/well for row C, 5 picomol/well for row D, 2.5

pmole/well for row E, 1.25 pmole/well for row F, 0.5 pmole/well for row G, and 0 pmole/well (buffer only) for row H.

- 5 4) Pipet 5 μ L compounds from each well of a compound dilution plate, for IC₅₀s, using the following dilution scheme:

Well H: 400 μ M compound (final concentration of compound in reaction mix = $5/100 \times 400 \mu\text{M} = 20 \mu\text{M}$)

10 Well G: 1:10 dilution of Well H (i.e. 5 μ L compound from well H + 45 μ L 100% DMSO) (final concentration = 2 μ M)

Well F: 1:10 dilution of well G (final concentration = 0.2 μ M)

Well E: 1:10 dilution of well F (final concentration = 0.02 μ M)

Well D: 1:10 dilution of well E (final concentration = 0.002 μ M)

Well C: 1:10 dilution of well D (final concentration = 0.0002 μ M)

15 Well B: 1:10 dilution of well C (final concentration = 0.00002 μ M)

Well A: 1:10 dilution of well B (final concentration = 0.000002 μ M)

20 IC₅₀s or EC₅₀s are done in triplicate. One Flashplate can therefore be set up to handle 3 compounds. (i.e., columns 2, 3, and 4 are for compound #1, columns 5, 6, and 7 are for compound #2, and columns 8, 9, and 10 are for compound #3.)

- 25 5) Add 50 μ L of **RUP3** membranes to all wells in Columns 2 to 10. (Prior to the start of the assay, the frozen membrane pellets for both **RUP3** and CMV (cells transfected with an expression plasmid containing no **RUP3** sequences), are suspended in binding buffer, usually 1 ml binding buffer for 1 plate of membranes. The membranes are kept in ice all the time, and a polytron (Brinkmann polytron, model # PT-3100) is used (setting 6-7, for 15-20 seconds) to obtain a homogeneous membrane suspension.) Protein concentration is determined by Bradford protein assay kit using instructions given in the kit, using the standard supplied with the kit as a reference. The protein concentration of the membranes is adjusted with binding buffer, so that 50 μ L membranes = 15 μ g protein (i.e. 0.3 mg/ml protein).
- 30

- 35 6) In column 1, Wells A, B, C, and D, add 50 μ L **RUP3** membranes. To wells E, F, G, and H, add 50 μ L CMV membranes, (CMV membranes being of the same protein concentration as the **RUP3** membranes).

- 7) Incubate 1 hour at room temperature with agitation on a rotating platform shaker. Cover with foil while shaking.

- 8) After 1 hour, add (to all 96 wells), 100 μ L of the 125 I tracer in detection buffer supplied with the Flashplate kit plus proclin, made up in the following manner:

Pipet per 10 ml per Flashplate: 100 ml of detection buffer + 1 ml 125 I + 0.2 ml of Proclin (the proclin helps to stop the production of cAMP). Make a smaller quantity of detection buffer mix if you have fewer plates.

- 9) Shake the plates on a rotating platform shaker for 2 hours, covering plates with lead sheeting.

- 10) Seal the plates with the plastic film sealers provided with the Flashplate kit.

- 11) Count the plates using a TRILUX 1450 Microbeta Counter. See the door of the counter to determine which counting protocol to use.

- 12) Data is analyzed on the Arena Database according to the **RUP3** non-fusion, IC_{50} EC_{50} for 96-well cAMP membrane assay, and the compound numbers and the concentrations of compounds must be entered by the user.

B. Membrane Cyclase Criteria

- 1) Signal to Noise:

An acceptable signal-to-noise ratio for **RUP3** can vary from 4 to 6. The raw cpm's are approximately 1800 to 2500 for **RUP3** and 3500-4500 for CMV. The cpm (or ultimately pmoles of cAMP/well) cannot be outside the standard curve, and should not approach well A of the standard curve (50 pmole/well) and well H (no cAMP). Generally, the pmoles of cAMP produced by **RUP3** receptor are around 11 to 13 pmole/well (for 15 μ g/well protein), and for CMV are between 2 to 3 pmole/well (for 15 μ g protein /well).

- 2) Standard curve:

The slope should be linear and the error bars for duplicates should be very small. The receptor and CMV controls cannot be off scale of the standard curve, as described above. If the receptor controls are off the high end of the standard curve, i.e. 50 pmole/well or higher, one must repeat the experiment using less protein. However, such a case has not been observed with transiently transfected **RUP3** membranes (10 μ g DNA/15 cm plate, using 60 μ L Lipofectamine, and preparing membranes after 24 hour of transfection.)

- 3) The IC₅₀ or EC₅₀ curve should be at 100% (+ or – 20%) of control **RUP3** membranes at the top, and should go down to 0 (or up to 20%) at the bottom. The standard error of the triplicate determinations should be + or – 10%.

5

C. Stimulation of cAMP in HIT-T15 cells

HIT-T15 (ATCC CRL#1777) is an immortalized hamster insulin-producing cell line. These cells express **RUP3** and therefore can be used to assess the ability of **RUP3** ligands to stimulate or inhibit cAMP accumulation via its endogenously expressed receptor. In this assay, cells are grown to 80% confluence and then distributed into a 96-well Flashplate (50,000 cells/ well) for detection of cAMP via a "cAMP Flashplate Assay" (NEN, Cat # SMP004). Briefly, cells are placed into anti-cAMP antibody-coated wells that contain either vehicle, the test ligand(s) at a concentration of interest, or 1 uM forskolin. The latter is a direct activator of adenylyl cyclase and serves as a positive control for stimulation of cAMP in HIT-T15 cells. All conditions are tested in triplicate. After a 1 hour incubation to allow for stimulation of cAMP, a Detection Mix containing ¹²⁵I-cAMP is added to each well and the plate is allowed to incubate for another 1 hour. The wells are then aspirated to remove unbound ¹²⁵I-cAMP. Bound ¹²⁵I-cAMP is detected using a Wallac Microbeta Counter. The amount of cAMP in each sample is determined by comparison to a standard curve, obtained by placing known concentrations of cAMP in some wells on the plate.

A number of the compounds disclosed herein were screened using the above described assay. Representative compounds and their corresponding EC₅₀ values are shown in the Table 6 below:

TABLE 6

Compound	RUP3 (EC₅₀) (μM)
A1	0.020
A34	0.027
A35	0.059

D. Stimulation of insulin secretion in HIT-T15 cells

It is known that stimulation of cAMP in HIT-T15 cells causes an increase in insulin secretion when the glucose concentration in the culture media is changed from 3mM to 15 mM. Thus, **RUP3** ligands can also be tested for their ability to stimulate glucose-dependent insulin secretion (GSIS) in HIT-T15 cells. In this assay, 30,000 cells/well in a 12-well plate are incubated in culture media containing 3 mM glucose and no serum for 2 hours. The media is then changed; wells receive media containing either 3 mM or 15 mM glucose, and in both cases the media contains either vehicle (DMSO) or **RUP3** ligand at a concentration of interest. Some wells receive media containing 1 uM forskolin as a positive control. All conditions are tested in triplicate. Cells are incubated for 30

minutes, and the amount of insulin secreted into the media is determined by ELISA, using a kit from either Peninsula Laboratories (Cat # ELIS-7536) or Crystal Chem Inc. (Cat # 90060).

5 **E. Stimulation of insulin secretion in isolated rat islets**

As with HIT-T15 cells, it is known that stimulation of cAMP in isolated rat islets causes an increase in insulin secretion when the glucose concentration in the culture media is changed from 60 mg/dl to 300 mg/dl. **RUP3** is an endogenously expressed GPCR in the insulin-producing cells of rat islets. Thus, **RUP3** ligands can also be tested for their ability to stimulate GSIS in rat islet cultures.

10 This assay is performed as follows:

- A. Select 75-150 islet equivalents (IEQ) for each assay condition using a dissecting microscope. Incubate overnight in low-glucose culture medium. (Optional.)
- B. Divide the islets evenly into triplicate samples of 25-40 islet equivalents per sample. Transfer to 40 μ m mesh sterile cell strainers in wells of a 6-well plate with 5 ml of low
15 (60 mg/dl) glucose Krebs-Ringers Buffer (KRB) assay medium.
- C. Incubate 30 minutes (1 hour if overnight step skipped) at 37°C and 5% CO₂. Save the supernatants if a positive control for the RIA is desired.
- D. Move strainers with islets to new wells with 5ml/well low glucose KRB. This is the second pre-incubation and serves to remove residual or carryover insulin from the
20 culture medium. Incubate 30 minutes.
- E. Move strainers to next wells (Low 1) with 4 or 5 ml low glucose KRB. Incubate @ 37° C for 30 minutes. Collect supernatants into low-binding polypropylene tubes pre-labelled for identification and keep cold.
- F. Move strainers to high glucose wells (300mg/dl, which is equivalent to 16.7mM).
25 Incubate and collect supernatants as before. Rinse islets in their strainers in low-glucose to remove residual insulin. If the rinse is to be collected for analysis, use one rinse well for each condition (i.e. set of triplicates.)
- G. Move strainers to final wells with low-glucose assay medium (Low 2). Incubate and collect supernatants as before.
- H. Keeping cold, centrifuge supernatants at 1800 rpm for 5 minutes @ 4-8°C to remove
30 small islets/islet pieces that escape the 40mm mesh. Remove all but lower 0.5 – 1 ml and distribute in duplicate to pre-labelled low-binding tubes. Freeze and store at <-20° C until insulin concentrations can be determined.
- I. Insulin determinations are done as above, or by Linco Labs as a custom service, using
35 their rat insulin RIA (Cat. # RI-13K).

Example 2**A. RT-PCR analysis of RUP3 expression in human tissues (Figure 1A).**

5 RT-PCR was applied to determine the tissue distribution of **RUP3**. Oligonucleotides used for PCR had the following sequences:

ZC47: 5'-CATTGCCGGGCTGTGGTTAGTGTC-3' (forward primer), (SEQ ID NO:3);

ZC48: 5'-GGCATAGATGAGTGGGTTGAGCAG-3' (reverse primer), (SEQ ID NO:4);

and the human multiple tissue cDNA panels (MTC, Clontech) were used as templates (1 ng
10 cDNA per PCR amplification). Twenty-two (22) human tissues were analyzed. PCR was performed using Platinum PCR SuperMix (Life Technologies, Inc.; manufacture instructions were followed) in a 50 µl reaction by the following sequences: step 1, 95°C for 4 min; step 2, 95°C for 1 min; step 3, 60°C for 30 sec; step 4, 72°C for 1 min; and step 5, 72°C for 7 min. Steps 2 through 4 were repeated 35 times.

15 The resulting PCR reactions (15 µl) were loaded on a 1.5% agarose gel to analyze the RT-PCR products, and a specific 466 base-pair DNA fragment representing **RUP3** was specifically amplified from cDNA of pancreas origin. Low expression was also evident in subregions of brain.

B. cDNA Dot-Blot analysis of RUP3 expression in human tissues (Figure 1B).

20 Results from RT-PCR analysis were further confirmed in cDNA dot-blot analysis. In this assay, a dot-blot membrane containing cDNA from 50 human tissues (Clontech) was hybridized with a ³²P-radiolabelled DNA probe having sequences derived from human **RUP3**. Hybridization signals were seen in pancreas and fetal liver, suggesting these tissues express **RUP3**. No significant expression was detected in other tissues analyzed.

25

C. Analysis of RUP3 by RT-PCR with isolated human pancreatic islets of Langerhans (Figure 1C).

Further analysis of **RUP3** by RT-PCR with isolated human pancreatic islets of Langerhans showed robust expression of **RUP3** in islet cells but not in control samples.

30

D. Analysis of RUP3 expression with cDNAs of rat origin by RT-PCR (Figure 1D).

RUP3 expression was further analyzed with cDNAs of rat origin by RT-PCR technique. Tissue cDNAs used for this assay were obtained from Clontech except those for hypothalamus and islets, which were prepared in house. Concentrations of each cDNA sample were normalized via a control RT-PCR analysis of the house-keeping gene GAPDH before assaying for **RUP3** expression.
35 Oligonucleotides used for PCR had the following sequences:

rat **RUP3** ("rRUP3") forward: 5'-CATGGGCCCTGCACCTTCTTTG-3' (SEQ ID NO:5);
 rRUP3 reverse: 5'-GCTCCGGATGGCTGATGATAGTGA-3' (SEQ ID NO:6).

PCR was performed using Platinum PCR SuperMix (Life Technologies, Inc.; manufacture instructions were followed) in a 50 μ l reaction by the following sequences: step 1, 95°C for 4 min; step 2, 95°C for 1 min; step 3, 60°C for 30 sec; step 4, 72°C for 1 min; and step 5, 72°C for 7 min. Steps 2 through 4 were repeated 35 times.

The resulting PCR reactions (15 μ l) were loaded on a 1.5% agarose gel to analyze the RT-PCR products, and a specific 547 base-pair DNA fragment representing rat **RUP3** was specifically amplified from cDNA of pancreas origin, revealing a similar expression profile with human. Of particular note, robust expression was seen in isolated islets and hypothalamus.

Example 3

RUP3 protein expression is restricted to β cell lineage of pancreatic islets (Figure 2).

A. A polyclonal anti-RUP3 antibody was prepared in rabbits (Figure 2A).

Rabbits were immunized with an antigenic peptide with sequence derived from rat **RUP3** ("rRUP3"). The peptide sequence was RGPERTRESAYHIVTISHPELDG and shared 100% identity with mouse **RUP3** in the corresponding region. A cysteine residue was incorporated at the N-terminal end of this antigenic peptide to facilitate KLH crosslinking before injecting into rabbits. The resulting antisera ("anti-rRUP3") and the corresponding preimmune sera ("pre-rRUP3") were tested for immune reactivity to mouse **RUP3** in immunoblotting assays (lanes 1 through 4). In this assay, the GST-RUP3 fusion protein was readily recognized by the anti-rRUP3 antisera (lane 4), but not by the preimmune sera (lane 2). The immunoreactive signal could be efficiently eliminated when the immunoblotting assay was performed in the presence of excess antigenic peptide (lane 6).

B. RUP3 expression in insulin-producing β cells of pancreatic islets (Figure 2B).

Rat pancreas was perfused with 4% paraformaldehyde (PFA) in PBS and embedded in OCT embedding medium. Ten micron sections were prepared, fixed on glass slides, and immunostained with either pre-rRUP3 (Figure 2B, panel a) or with anti-rRUP3 antisera (Figure 2B, panels c and e) followed by secondary staining with donkey anti-rabbit IgG conjugated to the fluorochrome Cy-3. Each section was also co-immunostained with a monoclonal anti-insulin antibody (Santa Cruz, Figure 2B, panels b and d) in primary staining followed by a secondary staining with donkey anti-mouse IgG conjugated with FITC, or with a goat anti-glucagon antibody (Santa Cruz, Figure 2B, panel f) and donkey anti-goat IgG coupled to FITC. Immunofluorescent signals were examined under a fluorescent microscope. **RUP3** was found expressed in insulin producing cells (panels c and d), but not in glucagons producing cells (panels e and f). These data demonstrated that **RUP3** is expressed in β cells but not in β cells of the rat pancreatic islets. Analogous results were obtained when mouse pancreatic sections were investigated for **RUP3** expression.

Example 4**Functional Activities of RUP3 *In Vitro* (Figure 3).**

5 It was established that **RUP3** stimulates the production of cAMP by cotransfection of 293 cells with: (1) a CRE-Luciferase reporter, wherein the ability to stimulate the production of firefly luciferase depends on increased cAMP in cells, and (2) an expression plasmid encoding the human form of **RUP3** (Figure 3A). Note that cells co-transfected with an expression plasmid containing no **RUP3** sequences ("CMV" in Figure 3A) produce very little luciferase activity, whereas cells
10 transfected with an expression plasmid encoding **RUP3** ("**RUP3**" in Figure 3A) have at least a 10-fold increase in luciferase activity. This indicates that **RUP3** stimulates the production of cAMP when introduced into 293 cells. This property of **RUP3** is conserved across species, because hamster **RUP3** stimulates luciferase activity when introduced into 293 cells in a manner analogous to that described for human **RUP3** (Figure 3B).

15 It is established that, when cAMP is increased in insulin-producing cells of the pancreas, these cells exhibit an enhanced ability to secrete insulin when glucose concentrations rise. To test whether **RUP3** might impart enhanced glucose-dependent insulin release, retrovirus containing human **RUP3** was used to generate Tu6 cells that express high levels of **RUP3**. Tu6 cells produce insulin, but do not express appreciable levels of **RUP3** and do not normally exhibit an increase in insulin release
20 when increased glucose is present in the culture media. As shown in Figure 3C, Tu6 cells transduced with a control virus that contains no receptor are still able to produce insulin, but do not show an increase in insulin secretion when the concentration of glucose in the culture media is shifted from 1 mM to 16 mM. By contrast, Tu6 cells transduced with **RUP3**-containing retrovirus display significant glucose-dependent insulin secretion (Figure 3C).

25

Example 5***In vivo* effects of RUP3 agonists on glucose homeostasis in rats.****A. Oral Glucose tolerance test (oGTT)**

30 Male Sprague Dawley rats weighing approximately 200g-250g were fasted for 15 hours and randomly grouped (n=6) to receive a **RUP3** agonist (Compounds A78, A88 or A118) at 3, 10 or 30 mg/kg. Compounds were delivered orally via a gavage needle (p.o., volume 3 ml/kg). At time 0, levels of blood glucose were assessed using a glucometer (Elite XL, Bayer), and rats were administered either vehicle (20% hydroxypropyl-beta-cyclodextrin) or test compound. Thirty minutes
35 after administration of test compound, levels of blood glucose were again assessed, and rats were administered dextrose orally at a dose of 2g/kg. Blood glucose measurements were then taken 30 min, 60 min, and 120 min after this time. Table 7 shows the mean percentage inhibition of glucose excursion for each test compound, averaged across the six animals in the treatment group. These

results demonstrated that the **RUP3** agonists, Compounds **A78**, **A88** and **A118** lowered blood glucose after challenge with glucose.

TABLE 7

Mean % Inhibition of Glucose Excursion

Compound	% inhibition of glucose excursion, (dose, mg/kg)
A78	39%, (10)
A88	38%, (30)
A118	43%, (30)

5

Example 6**Generation of Tu6/ RUP3 Stable Lines**

To produce Tu6 cells that express **RUP3** at high levels, a retrovirus bearing an expression cassette for **RUP3** was generated. Briefly, **RUP3** coding sequence was cloned into the retroviral vector pLNCX2 (Clontech, Cat # 6102-1). The amphotropic packaging cell line PT-67 (Clontech, K1060-D) was then transfected with either the parental vector pLNCX2 or pLNCX2/**RUP3** using Lipofectamine and stable lines were established using guidelines provided by the PT-67 vendor. Retrovirus-containing supernatant was obtained by collecting media from the resultant stables according to the manufacturer's directions. Tu6 cells, in a 10 cm dish, were then infected with retrovirus by incubating in a solution of 1 ml viral supernatant/ 9 ml culture media containing 40 ug/ml polybrene for 24 hours. The medium was then changed to culture media containing 300 ug/ml G418. G418-resistant clones were ultimately created by virtue of the neomycin-resistance gene cassette present in the pLNCX2 vector, thus indicating the successful integration of retrovirus into the Tu6 genome. The expression of **RUP3** in the Tu6/**RUP3** G418-resistant colonies was confirmed by Northern blot.

20

Example 7**Insulin secretion, Tu6 Stables**

To measure insulin secretion from rodent insulin-producing cell lines, cells were first cultured overnight in serum-free, glucose-deficient media. The following morning, the cells were then placed in the same media supplemented with either 1 mM or 16 mM glucose. After an incubation of 4 hours, the media was collected and analyzed for insulin content using a Rat Insulin Enzyme-Immunoassay (EIA) System (Amersham Pharmacia Biotech, Cat. # RPN 2567). Typically, the assay was performed using multiple dilutions of sample media in order to ensure that the sample measurements fell within the boundaries of the standard curve (generated using known amounts of insulin), as recommended by the manufacturer.

30

Example 8**Receptor Binding Assay**

In addition to the methods described herein, another means for evaluating a test compound is by determining binding affinities to the **RUP3** receptor. This type of assay generally requires a radiolabelled ligand to the **RUP3** receptor. Absent the use of known ligands for the **RUP3** receptor and radiolabels thereof, compounds of Formula (I) can be labelled with a radioisotope and used in an assay for evaluating the affinity of a test compound to the **RUP3** receptor.

A radiolabelled **RUP3** compound of Formula (I) can be used in a screening assay to identify/evaluate compounds. In general terms, a newly synthesized or identified compound (i.e., test compound) can be evaluated for its ability to reduce binding of the "radiolabelled compound of Formula (I)" to the **RUP3** receptor. Accordingly, the ability to compete with the "radio-labelled compound of Formula (I)" or **Radiolabelled RUP3 Ligand** for the binding to the **RUP3** receptor directly correlates to its binding affinity of the test compound to the **RUP3** receptor.

15 ASSAY PROTOCOL FOR DETERMINING RECEPTOR BINDING FOR RUP3:**A. RUP3 RECEPTOR PREPARATION**

293 cells (human kidney, ATCC), transiently transfected with 10 ug human **RUP3** receptor and 60 µL Lipofectamine (per 15-cm dish), were grown in the dish for 24 hours (75% confluency) with a media change and removed with 10 ml/dish of Hepes-EDTA buffer (20mM Hepes + 10 mM EDTA, pH 7.4). The cells were then centrifuged in a Beckman Coulter centrifuge for 20 minutes, 17,000 rpm (JA-25.50 rotor). Subsequently, the pellet was resuspended in 20 mM Hepes + 1 mM EDTA, pH 7.4 and homogenized with a 50- ml Dounce homogenizer and again centrifuged. After removing the supernatant, the pellets were stored at -80°C, until used in binding assay. When used in the assay, membranes were thawed on ice for 20 minutes and then 10 mL of incubation buffer (20 mM Hepes, 1 mM MgCl₂, 100 mM NaCl, pH 7.4) added. The membranes were then vortexed to resuspend the crude membrane pellet and homogenized with a Brinkmann PT-3100 Polytron homogenizer for 15 seconds at setting 6. The concentration of membrane protein was determined using the BRL Bradford protein assay.

30 B. BINDING ASSAY

For total binding, a total volume of 50 µL of appropriately diluted membranes (diluted in assay buffer containing 50 mM Tris HCl (pH 7.4), 10 mM MgCl₂, and 1mM EDTA; 5-50 µg protein) is added to 96-well polypropylene microtiter plates followed by addition of 100 µL of assay buffer and 50 µL of **Radiolabelled RUP3 Ligand**. For nonspecific binding, 50 µL of assay buffer is added instead of 100 µL and an additional 50 µL of 10 µM cold **RUP3** is added before 50 µL of **Radiolabelled RUP3 Ligand** is added. Plates are then incubated at room temperature for 60-120 minutes. The binding reaction is terminated by filtering assay plates through a Microplate Devices

GF/C Unifilter filtration plate with a Brandell 96-well plate harvester followed by washing with cold 50 mM Tris HCl, pH 7.4 containing 0.9% NaCl. Then, the bottom of the filtration plate are sealed, 50 μ L of Optiphase Supermix is added to each well, the top of the plates are sealed, and plates are counted in a Trilux MicroBeta scintillation counter. For compound competition studies, instead of adding 100 μ L of assay buffer, 100 μ L of appropriately diluted test compound is added to appropriate wells followed by addition of 50 μ L of **Radiolabelled RUP3 Ligand**.

C. CALCULATIONS

The test compounds are initially assayed at 1 and 0.1 μ M and then at a range of concentrations chosen such that the middle dose would cause about 50% inhibition of a **Radio-RUP3 Ligand** binding (i.e., IC_{50}). Specific binding in the absence of test compound (B_0) is the difference of total binding (B_T) minus non-specific binding (NSB) and similarly specific binding (in the presence of test compound) (B) is the difference of displacement binding (B_D) minus non-specific binding (NSB). IC_{50} is determined from an inhibition response curve, logit-log plot of % B/B_0 vs concentration of test compound.

K_i is calculated by the Cheng and Prustoff transformation:

$$K_i = IC_{50} / (1 + [L]/K_D)$$

where $[L]$ is the concentration of a **Radio-RUP3 Ligand** used in the assay and K_D is the dissociation constant of a **Radio-RUP3 Ligand** determined independently under the same binding conditions.

CHEMISTRY

SYNTHESES OF COMPOUNDS OF THE PRESENT INVENTION

EXAMPLE 9

The compounds of the invention and their synthesis are further illustrated by the following examples. The following examples are provided to further define the invention without, however, limiting the invention to the particulars of these examples. The compounds described herein, *supra* and *infra*, are named according to the CS Chem Draw Ultra Version 7.0.1, AutoNom version 2.2. In certain instances common names are used and it is understood that these common names would be recognized by those skilled in the art.

Chemistry: Proton nuclear magnetic resonance (1H NMR) spectra were recorded on a Varian Mercury Vx-400 equipped with a 4 nucleus auto switchable probe and z-gradient or a Bruker Avance-400 equipped with a QNP (Quad Nucleus Probe) or a BBI (Broad Band Inverse) and z-gradient. Chemical shifts are given in parts per million (ppm) with the residual solvent signal used as reference. NMR abbreviations are used as follows: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad. Microwave irradiations were carried out using the Emyrs Synthesizer (Personal

Chemistry). Thin-layer chromatography (TLC) was performed on silica gel 60 F₂₅₄ (Merck), preparatory thin-layer chromatography (prep TLC) was performed on PK6F silica gel 60 A 1 mm plates (Whatman), and column chromatography was carried out on a silica gel column using Kieselgel 60, 0.063-0.200 mm (Merck). Evaporation was done in vacuo on a Buchi rotary evaporator. Celite 545 ® was used during palladium filtrations.

LCMS specs: 1) PC: HPLC-pumps: LC-10AD VP, Shimadzu Inc.; HPLC system controller: SCL-10A VP, Shimadzu Inc; UV-Detector: SPD-10A VP, Shimadzu Inc; Autosampler: CTC HTS, PAL, Leap Scientific; Mass spectrometer: API 150EX with Turbo Ion Spray source, AB/MDS Sciex; Software: Analyst 1.2. 2) Mac: HPLC-pumps: LC-8A VP, Shimadzu Inc; HPLC system controller: SCL-10A VP, Shimadzu Inc. UV-Detector: SPD-10A VP, Shimadzu Inc; Autosampler: 215 Liquid Handler, Gilson Inc; Mass spectrometer: API 150EX with Turbo Ion Spray source, AB/MDS Sciex Software: Masschrom 1.5.2.

Example 9.1: Preparation of 4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester (Compound A1)

4-Hydroxy-piperidine-1-carboxylic acid tert-butyl ester (3.03 mmol, 610 mg) and sodium hydride (10.6 mmol, 255 mg) were dissolved in dry THF (20 mL) and stirred for 30 minutes at room temperature. Then, (6-chloro-5-nitro-pyrimidin-4-yl)-(4-methanesulfonyl-phenyl)-amine (3.03 mmol, 1.0 g) was added. The reaction was stirred at room temperature for 30 minutes. Its progress was monitored by thin layer chromatography and LCMS. Sodium hydride was quenched with water and the desired compound was extracted in ethyl acetate. Organic solvents were evaporated in vacuo. Flash chromatography (Silica gel 60; 30/70 EtOAc/Hexanes) provided Compound A1 as a yellow solid (1.2 g, 68%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.10 (s, 1H), 8.33 (s, 1H), 7.90 (d, 2H), 7.79 (d, 2H), 5.51 (heptet, 1H), 3.58 (m, 2H), 3.46 (m, 2H), 2.97 (s, 3H), 1.84 (m, 4H), 1.36 (s, 9H). LCMS (ESI), m/z 494.4 (M+H⁺, 100%).

Example 9.2: Preparation of (4-Methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (Compound A2)

Compound A1 (1.42 mmol, 700 mg) was dissolved in a commercially available 4M HCl solution in 1,4-dioxane (25 mL). The mixture was stirred at 40°C for 1.0 hour. Removal of organic solvents in vacuo provided compound A2 as a yellow solid (580 mg, 100%). ¹H NMR (400 MHz, MeOH-d₄) δ (ppm): 8.29 (s, 1H), 7.81 (quartet, 4H), 5.56 (m, 1H), 3.21 (m, 4H), 3.00 (s, 3H), 2.07 (m, 4H). LCMS (ESI), m/z 394.1 (M+H⁺, 100%).

Example 9.3: Preparation of 1-{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-3,3-dimethyl-butan-1-one (Compound A3).

Compound A2 (0.12 mmol, 50 mg), 3,3-dimethyl-butyryl chloride (0.18 mmol, 24 mg) and triethylamine (0.63 mmol, 88 L) were dissolved in DMF and microwaved at 80°C for 5 minutes. The reaction mixture was quenched with water and extracted with ethyl acetate. Removal of organic solvents in vacuo provided Compound A3 as a yellow solid (46 mg, 78%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 9.98 (s, 1H), 8.20 (s, 1H), 7.77 (d, 2H), 7.64 (d, 2H), 5.44 (heptet, 1H), 3.71 (m, 1H), 3.48 (m, 3H), 2.75 (s, 3H), 2.11 (quartet, 2H), 1.76 (m, 4H), 0.85 (s, 9H). LCMS (ESI), m/z 492.4 (M+H⁺, 100%).

Example 9.4: Preparation of (4-Methanesulfonyl-phenyl)-[5-nitro-6-(1-thiophen-3-ylmethyl-piperidin-4-yloxy)-pyrimidin-4-yl]-thiophen-3-ylmethyl-amine (Compound A4).

Compound A2 (0.12 mmol, 50 mg), 3-chloromethyl-thiophene (0.12 mmol, 16 mg) and triethylamine (0.63 mmol, 88 μL) were dissolved in DMF and microwaved at 80°C for 10 minutes. The reaction mixture was quenched with water and extracted with ethyl acetate. Purification by HPLC provided Compound A4 as a yellow solid (24 mg, 34%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.35 (s, 1H), 7.93 (m, 2H), 7.79 (m, 2H), 7.22 (m, 1H), 7.19 (m, 2H), 7.08 (m, 1H), 6.98 (m, 1H), 6.84 (m, 1H), 5.68 (m, 1H), 4.08 (m, 4H), 3.36 (m, 2H), 3.04 (s, 3H), 2.86 (m, 2H), 2.34 (m, 2H), 2.07 (m, 2H). LCMS (ESI), m/z 586.1 (M+H⁺, 100%).

Example 9.5: Preparation of (4-Methanesulfonyl-phenyl)-[5-nitro-6-(1-pyridin-2-ylmethyl-piperidin-4-yloxy)-pyrimidin-4-yl]-amine (Compound A5).

Compound A2 (0.12 mmol, 50 mg), 2-chloromethyl-pyridine (0.12 mmol, 20 mg) and triethylamine (0.63 mmol, 88 μL) were dissolved in DMF and microwaved at 80°C for 10 minutes. The reaction mixture was quenched with water and extracted with ethyl acetate. Removal of organic solvent in vacuo provided Compound A5 pure as a yellow solid (27 mg, 47%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.09 (s, 1H), 8.51 (m, 1H), 8.32 (s, 1H), 7.89 (d, 2H), 7.78 (d, 2H), 7.60 (t, 1H), 7.36 (m, 1H), 7.13 (t, 1H), 5.40 (m, 1H), 3.65 (m, 2H), 3.07 (s, 3H), 2.72 (m, 2H), 2.45 (m, 2H), 2.02 (m, 2H), 1.91 (m, 2H). LCMS (ESI), m/z 484.5 (M+H⁺, 100%).

Example 9.6: Preparation of (4-Methanesulfonyl-phenyl)-[5-nitro-6-(1-pyridin-3-ylmethyl-piperidin-4-yloxy)-pyrimidin-4-yl]-amine (Compound A6).

Compound A2 (0.12 mmol, 50 mg), 3-chloromethyl-pyridine (0.12 mmol, 20 mg) and triethylamine (0.63 mmol, 88 μL) were dissolved in DMF and microwaved at 80°C for 10 minutes. The reaction mixture was quenched with water and extracted with ethyl acetate. Removal of organic solvent in vacuo provided Compound A6 pure as a yellow solid (39 mg, 66%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.09 (s, 1H), 8.45 (m, 2H), 8.33 (s, 1H), 7.90 (d, 2H), 7.78 (d, 2H), 7.65 (m, 1H),

7.21 (m, 1H), 5.41 (heptet, 1H), 3.52 (m, 2H), 3.01 (s, 3H), 2.65 (m, 2H), 2.47 (m, 2H), 1.98 (m, 2H), 1.94 (m, 2H). LCMS (ESI), m/z 484.3 (M+H⁺, 100%).

Example 9.7: Preparation of {6-[1-(3,3-Dimethyl-butyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine (Compound A7).

Compound A2 (0.20 mmol, 80 mg), and 3,3-dimethyl-butyraldehyde (0.24 mmol, 30 μ L) were dissolved in methanol (2 mL) and stirred for 5 minutes at room temperature. Then, added sodium borohydride (0.25 mmol, 8.7 mg) and stirred for 10 minutes at room temperature. The reaction mixture was quenched with saturated ammonium chloride solution (1 mL) followed by an extraction with dichloromethane. Removal of organic solvent in vacuo and purification by prep-LCMS provided Compound A7 as a yellow solid (12 mg, 13%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.17 (s, 1H), 8.36 (s, 1H), 7.90 (d, 2H), 7.78 (d, 2H), 5.71 (s broad, 1H), 3.51 (d, 2H), 3.08 (m, 2H), 2.97 (m, 5H), 2.38 (m, 2H), 2.14 (m, 2H), 1.60 (m, 2H), 0.85 (s, 9H). LCMS (ESI), m/z 478.3 (M+H⁺, 100%).

Example 9.8: Preparation of (4-Methanesulfonyl-phenyl)-{6-[1-(3-methyl-butyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-amine (Compound A8).

Compound A2 (0.15 mmol, 60 mg), and 3-methyl-butyraldehyde (0.15 mmol, 13 mg) were dissolved in methanol (2 mL) and stirred for 5 minutes at room temperature. Then, sodium borohydride (0.18 mmol, 6.3 mg) was added at 0°C. The reaction was complete immediately upon addition of sodium borohydride. The mixture was quenched with saturated ammonium chloride solution (1 mL) followed by an extraction with dichloromethane. Removal of organic solvent in vacuo and purification by HPLC provided Compound A8 as a yellow solid (25 mg, 36%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.17 (s, 1H), 8.35 (s, 1H), 7.90 (d, 2H), 7.80 (d, 2H), 5.72 (s broad, 1H), 3.65 (m, 2H), 3.11 (m, 2H), 2.96 (m, 5H), 2.40 (m, 2H), 2.15 (m, 2H), 1.60 (m, 3H), 0.85 (d, 6H). LCMS (ESI), m/z 464.4 (M+H⁺, 100%).

Example 9.9: Preparation of (4-Methanesulfonyl-phenyl)-[5-nitro-6-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yloxy)-pyrimidin-4-yl]-amine (Compound A9).

Compound A2 (0.13 mmol, 50 mg), and 2-bromo-pyridine (0.53 mmol, 53 μ L) were dissolved in DMF (1 mL) and triethylamine (0.46 mmol, 63 μ L). The reaction was heated in a microwave at 165°C for 40 minutes. The mixture was quenched with water and extracted with ethyl acetate. Removal of organic solvent in vacuo and purification by prep-TLC provided Compound A9 as a yellow solid (12 mg, 20%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.10 (s, 1H), 8.35 (s, 1H), 8.13 (m, 1H), 7.89 (d, 2H), 7.81 (d, 2H), 7.42 (m, 1H), 6.64 (d, 1H), 6.55 (m, 1H), 5.58 (heptet, 1H), 3.78 (m, 2H), 3.60 (m, 2H), 3.00 (s, 3H), 2.02 (m, 2H), 1.91 (m, 2H). LCMS (ESI), m/z 471.4 (M+H⁺, 100%).

Example 9.10: Preparation of 4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid ethyl ester (Compound A10).

Compound A2 (0.13 mmol, 50 mg), and ethyl chloroformate (0.13 mmol, 13 μ L) were dissolved in DMF (1 mL) and triethylamine (0.36 mmol, 50 μ L). The reaction was heated in a microwave at 80°C for 4 minutes. The mixture was quenched with water and extracted with ethyl acetate. Removal of organic solvent in vacuo provided Compound A10 as a yellow solid (50 mg, 89%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 9.97 (s, 1H), 8.20 (s, 1H), 7.77 (d, 2H), 7.66 (d, 2H), 5.41 (heptet, 1H), 3.96 (q, 2H), 3.47 (m, 4H), 2.88 (s, 3H), 1.72 (m, 4H), 1.08 (t, 3H). LCMS (ESI), m/z 466.3 (M+H⁺, 100%).

Example 9.11: Preparation of 1-{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-3,3-dimethyl-butan-2-one (Compound A11).

Compound A2 (0.12 mmol, 50 mg), and 1-bromo-3,3-dimethyl-butan-2-one (0.12 mmol, 16 μ L) were dissolved in DMF (1 mL) and triethylamine (0.36 mmol, 50 μ L). The reaction was heated in a microwave at 80°C for 4 minutes. The mixture was quenched with water and extracted with ethyl acetate. Removal of organic solvent in vacuo and purification by HPLC provided Compound A11 as a yellow solid (15 mg, 25%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.11 (s, 1H), 8.32 (s, 1H), 7.89 (d, 2H), 7.78 (d, 2H), 5.49 (s broad, 1H), 3.53 (s broad, 2H), 3.01 (s, 3H), 2.78 (m, 4H), 2.18 (m, 2H), 1.96 (m, 2H), 1.20 (s, 9H). LCMS (ESI), m/z 492.3 (M+H⁺, 100%).

Example 9.12: Preparation of {6-[1-(2-Ethoxy-ethyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine (Compound A12).

Compound A2 (0.13 mmol, 50 mg), and 1-bromo-2-ethoxy-ethane (0.65 mmol, 99 mg) were dissolved in DMF (1 mL) and triethylamine (0.91 mmol, 127 μ L). The reaction was heated in a microwave at 80°C for 20 minutes. The mixture was quenched with water and extracted with ethyl acetate. Removal of organic solvent in vacuo and purification by prep-TLC provided Compound A12 as a yellow solid (20 mg, 33%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.06 (s, 1H), 8.29 (s, 1H), 7.84 (d, 2H), 7.74 (d, 2H), 5.39 (s broad, 1H), 3.53 (m, 2H), 3.39 (q, 2H), 2.86 (s, 3H), 2.77 (m, 2H), 2.65 (m, 3H), 2.04 (m, 2H), 1.92 (m, 3H), 1.09 (m, 3H). LCMS (ESI), m/z 466.3 (M+H⁺, 100%).

Example 9.13: Preparation of 4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxymethyl]-piperidine-1-carboxylic acid tert-butyl ester (Compound A13).

4-Hydroxymethyl-piperidine-1-carboxylic acid tert-butyl ester (1.0 mmol, 226 mg) and sodium hydride (1.0 mmol, 25 mg) were dissolved in dimethyl acetamide (1.0 mL) and stirred for 30 minutes at room temperature. Then, (6-chloro-5-nitro-pyrimidin-4-yl)-(4-methanesulfonyl-phenyl)-

amine (0.21 mmol, 70 mg) was added. The reaction was stirred at 70°C for 20 minutes and the progress of the reaction was monitored by thin layer chromatography and LCMS. Sodium hydride was quenched with water and the desired compound was extracted in ethyl acetate. Organic solvents were evaporated in vacuo. Flash chromatography (Silica gel 60; 40/60 EtOAc/Hexanes) provided
5 Compound A13 as a yellow solid (10 mg, 10%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.21 (s, 1H), 8.41 (s, 1H), 7.97 (d, 2H), 7.86 (d, 2H), 4.39 (d, 2H), 4.17 (m, 2H), 3.07 (s, 3H), 2.76 (m, 2H), 1.83 (m, 2H), 1.59 (m, 1H), 1.45 (s, 9H), 1.30 (m, 2H). LCMS (ESI), m/z 408.2 (M+H⁺, 100%).

Example 9.14: Preparation of 4-{2-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-ethyl}-piperidine-1-carboxylic acid tert-butyl ester (Compound A14).

4-(2-Hydroxy-ethyl)-piperidine-1-carboxylic acid tert-butyl ester (1.0 mmol, 230 μL) and sodium hydride (1.0 mmol, 26 mg) were dissolved in dimethyl acetamide (1.0 mL) and stirred for 30 minutes at room temperature. Then, (6-chloro-5-nitro-pyrimidin-4-yl)-(4-methanesulfonyl-phenyl)-amine (0.21 mmol, 70 mg) was added. The reaction was stirred at 70°C for 20 minutes. Its progress
15 was monitored by thin layer chromatography and LCMS. Sodium hydride was quenched with water and the desired compound was extracted in ethyl acetate. Organic solvents were evaporated in vacuo. Flash chromatography (Silica gel 60; 40/60 EtOAc/Hexanes) provided Compound A14 as a yellow oil (90 mg, 82%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.26 (s, 1H), 8.40 (s, 1H), 7.98 (d, 2H), 7.86 (d, 2H), 4.51 (t, 2H), 4.09 (m, 2H), 3.73 (t, 2H), 3.07 (s, 3H), 2.72 (m, 2H), 1.76 (m, 1H), 1.55 (q, 2H),
20 1.46 (s, 9H), 1.15 (m, 2H). LCMS (ESI), m/z 422.2 (M+H⁺, 100%).

Example 9.15: Preparation of 3-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester (Compound A15).

3-Hydroxy-pyrrolidine-1-carboxylic acid tert-butyl ester (1.0 mmol, 197 mg) and sodium
25 hydride (1.0 mmol, 26 mg) were dissolved in THF (1.5 mL) and stirred for 30 minutes at room temperature. Then, (6-chloro-5-nitro-pyrimidin-4-yl)-(4-methanesulfonyl-phenyl)-amine (0.21 mmol, 70 mg) was added. The reaction was stirred at 0°C for 30 minutes. Its progress was monitored by thin layer chromatography and LCMS. Sodium hydride was quenched with water and the desired compound was extracted in ethyl acetate. Organic solvents were evaporated in vacuo. Flash
30 chromatography (Silica gel 60; 50/50 EtOAc/Hexanes) provided Compound A15 as a yellow oil (60 mg, 60%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.18 (s, 1H), 8.47 (s, 1H), 7.98 (d, 2H), 5.78 (m, 2H), 5.78 (m, 1H), 4.46 (m, 2H), 3.08 (s, 3H), 2.26 (m, 2H), 1.63 (m, 2H), 1.48 (s, 9H). LCMS (ESI), m/z 480.4 (M+H⁺, 100%).

Example 9.16: Preparation of 3-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxymethyl]-pyrrolidine-1-carboxylic acid tert-butyl ester (Compound A16).

3-Hydroxymethyl-pyrrolidine-1-carboxylic acid tert-butyl ester (0.65 mmol, 131 mg) and sodium hydride (1.3 mmol, 31 mg) were dissolved in N,N-dimethyl acetamide (1.5 mL) and stirred for 30 minutes at room temperature. Then, (6-chloro-5-nitro-pyrimidin-4-yl)-(4-methanesulfonyl-phenyl)-amine (0.26 mmol, 84 mg) was added. The reaction was stirred at 70°C for 30 minutes. Its progress was monitored by thin layer chromatography and LCMS. Sodium hydride was quenched with water and the desired compound was extracted in ethyl acetate. Organic solvents were evaporated in vacuo. Flash chromatography (Silica gel 60; 50/50 EtoAc/Hexanes) provided Compound A16 as a yellow solid (96 mg, 54%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.21 (s, 1H), 8.41 (s, 1H), 7.97 (d, 2H), 7.86 (d, 2H), 4.52 (m, 2H), 3.49 (m, 2H), 3.11 (s, 3H), 2.75 (m, 1H), 1.84 (m, 2H), 1.65 (m, 2H), 1.46 (s, 9H). LCMS (ESI), m/z 394.1 (M+H⁺, 100%).

Example 9.17: Preparation of 3-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-ylloxymethyl]-pyrrolidine-1-carboxylic acid tert-butyl ester (Compound A17).

(S)-3-Hydroxymethyl-pyrrolidine-1-carboxylic acid tert-butyl ester (0.65 mmol, 131 mg) and sodium hydride (1.3 mmol, 31 mg) were dissolved in N,N-dimethyl acetamide (1.5 mL) and stirred for 30 minutes at room temperature. Then, (6-chloro-5-nitro-pyrimidin-4-yl)-(4-methanesulfonyl-phenyl)-amine (0.26 mmol, 84 mg) was added. The reaction was stirred at 70°C for 30 minutes. Its progress was monitored by thin layer chromatography and LCMS. Sodium hydride was quenched with water and the desired compound was extracted in ethyl acetate. Organic solvents were evaporated in vacuo. Flash chromatography (Silica gel 60; 50/50 EtoAc/Hexanes) provided Compound A17 as a yellow solid (26 mg, 15%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 10.22 (s, 1H), 8.41 (s, 1H), 7.97 (d, 2H), 7.87 (d, 2H), 4.52 (m, 2H), 3.49 (m, 2H), 3.09 (s, 3H), 2.75 (m, 1H), 1.97 (m, 2H), 1.67 (m, 2H), 1.49 (s, 9H). LCMS (ESI), m/z 394.1 (M+H⁺, 100%).

Example 9.18: Preparation of 4-[5-Cyano-6-(6-methylsulfanyl-pyridin-3-ylamino)-pyrimidin-4-yl]oxy]-piperidine-1-carboxylic acid tert-butyl ester (Compound A18).

To a solution of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester (304 mg, 1.51 mmol) in DMF was added sodium hydride (36 mg, 1.51 mmol) and allowed the resulting mixture was allowed to stir at room temperature. After 30 minutes, 4-chloro-6-(6-methylsulfanyl-pyridin-3-ylamino)-pyrimidine-5-carbonitrile was added and the resulting mixture was heated at 70°C for 1 hour. Worked up with Ethyl acetate, sodium bicarbonate, dried with magnesium sulfate and evaporated to afford a white solid as Compound A18 (80.0 mg, 59.8%). ¹H NMR 400MHz DMSO-d₆ δ (ppm): 9.90 (s, 1H), 8.54 (d, 1H), 8.40 (s, 1H, pyrimidine), 7.78 (m, 1H), 7.29 (d, 1H), 5.75 (s, 3H), 5.35 (m, 1H), 3.58 (m, 2H), 3.27 (m, 2H), 1.93 (m, 2H), 1.63 (m, 2H), 1.38 (s, 9H), LCMS (ESI) for C₂₁H₂₆ClN₆OS: m/z 443.4 (M + H⁺, 100%).

The intermediate 4-chloro-6-(6-methylsulfanyl-pyridin-3-ylamino)-pyrimidine-5-carbonitrile was prepared in the following manner:

A. 4,6-Dichloro-pyrimidine-5-carbaldehyde

Phosphorus oxychloride (200 mL, 2184.8mmol) was added drop wise (via additional funnel) to DMF cooled to 0°C. After for 1 hour, 4,6 dihydroxypyrimidine (50.0 g, 446.1 mmol) was added and the mixture was allowed to warm to room temperature. The resulting heterogeneous mixture was refluxed for 3 hours. The volatiles were removed at reduce pressure, and the residue was poured in ice water and extract with CHCl₃/Et₂O, wash with sodium bicarbonate and concentrate under high vacuum. Final product was purified by silica plug using CH₂Cl₂ to afford a yellow solid (54.0 g), ¹H NMR 400MHz CDCl₃ δ (ppm): 10.3 (s, 1H, aldehyde), 8.7 (s, 1H, pyrimidine).

B. 4,6-Dichloro-pyrimidine-5-carbonitrile

4,6-Dichloro-pyrimidine-5-carbaldehyde (15.0 g, 84.75 mmol, 1.0 equivalent) was dissolved in ethyl acetate (150 mL), mixed with hydroxylamine hydrochloride in water (30mL) and added sodium acetate. Reaction was at left room temperature for 1.5 hours. Worked up with ethyl acetate, sodium bicarbonate, dried with magnesium sulfate, rotovaped and dried and high vacuum to afford a white solid (14.593 g). The white solid (iminohydroxy intermediate) was added to thionyl chloride (100 mL) at 0°C with stirring and allowed to warm to room temperature for 3 hours. The reaction was quenched in ice (500g), and the precipitated was filtered off, washed with cold water, and dried under high vacuum to afford a white solid as product (10.739g, 72.8%). ¹H NMR 400MHz CDCl₃ δ (ppm): 8.95 (s, 1H, pyrimidine).

C. 4-Chloro-6-(6-methylsulfanyl-pyridin-3-ylamino)-pyrimidine-5-carbonitrile

6-Methylsulfanyl-pyridin-3-ylamine (500.0 mg, 3.57 mmol, 1.0 equivalent) in DMF (1mL) was added drop wise to a suspension of 4,6-dichloro-pyrimidine-5-carbonitrile (616.9 mg, 3.57 mmol, 1.0equivalent), potassium carbonate (542.1mg, 3.92mmol, 1.1equivalent) at 0°C under stirring. The reaction was left reacting at room temperature for 1.5 hours. Product was crystallized using ethyl acetate, hexane getting a yellow solid as product (650.00mg, 65.62%). LCMS (ESI) for C₁₁H₈CIN₅S: m/z 278.0 (M + H⁺, 100%).

Example 9.19: Preparation of 4-[5-Cyano-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester (Compound A19).

To a solution of Compound A18 (52.0 mg, 0.11 mmol) in CH₂Cl₂ (5 mL) was added mCPBA (101.5 mg, 0.59 mmol) and the resulting mixture was heated to reflux. After 30 minutes, the mixture was worked up with water (basic conditions using ammonium hydroxide, pH=10), dichloromethane, and sodium bicarbonate, dried with magnesium sulfate and evaporated to afford Compound A19 as a white solid (24.9mg, 43.9%). ¹H NMR 400MHz CDCl₃ δ (ppm): 8.92 (d, 1H), 8.52 (d, 1H), 8.46 (s, 1H, pyrimidine), 8.10 (d, 1H), 7.47 (s, 1H), 5.45 (m, 1H), 3.77 (m, 2H), 3.37 (m, 2H), 3.24 (m, 3H), 1.98 (m, 2H), 1.84 (m, 2H), 1.48 (s, 9H), LCMS (ESI) for C₂₁H₂₆CIN₆OS: m/z 474.9 (M + H⁺, 100%).

Example 9.20: Preparation of [6-(1-Hexyl-piperidin-4-yloxy)-5-nitro-pyrimidin-4-yl]-(4-methanesulfonyl-phenyl)-amine (Compound A20).

General procedure; alkoxide substitution of (6-Chloro-5-nitro-pyrimidin-4-yl)-(4-methanesulfonyl-phenyl)-amine: In a 16 mL reaction vial was placed sodium hydride (25 mg, 60% in oil, 0.625 mmol) and 1.5 mL of THF. 1-Hexyl-piperidin-4-ol (30 mg, 0.162 mmol) was added to the suspension and the mixture was stirred for 20 min under N₂ at room temperature, followed by the addition of (6-chloro-5-nitro-pyrimidin-4-yl)-(4-methanesulfonyl-phenyl)-amine (41mg, 0.125mmol). After stirring overnight under N₂ at room temperature, all of the starting chloropyrimidines was completely converted as indicated by LCMS. The reaction mixture was then concentrated under vacuum and purified by preparative HPLC to give Compound A20. ¹H NMR (CDCl₃, 400MHz) δ 0.89 (m, 2H), 1.37 (m, 6H), 1.80 (m, 2H), 2.21 (m, 2H), 2.56 (m, 2H), 3.03 (m, 2H), 3.08 (s, 3H), 3.18 (m, 2H), 3.56 (m, 2H), 5.79 (m, 1H), 7.86 (d, 2H), 7.98 (d, 2H), 8.40 (s, 1H), 10.23 (s, 1H), 12.5 (s, 1H). Exact mass calculated for C₂₂H₃₁N₅O₅S 477.20, found 478.4 (MH⁺).

Example 9.21: Preparation of [6-(1-Cyclopropylmethyl-piperidin-4-yloxy)-5-nitro-pyrimidin-4-yl]-(4-methanesulfonyl-phenyl)-amine (Compound A21).

Compound A21 was prepared in a similar manner as described above using N-cyclopropyl-4-hydroxy-piperidine. ¹H NMR (CDCl₃, 400 MHz) δ 0.43 (m, 2H), 0.82 (m, 2H), 1.18 (m, 1H), 2.26 (m, 2H), 2.56 (m, 2H), 3.01 (m, 2H), 3.08 (s, 3H), 3.25 (m, 2H), 3.69 (m, 2H), 5.80 (m, 1H), 7.87 (d, 2H), 7.97 (d, 2H), 8.44 (s, 1H), 10.24 (s, 1H), 12.0 (s, 1H). Exact mass calculated for C₂₀H₂₅N₅O₅S 447.16, found 448.3 (MH⁺).

Example 9.22: Preparation of 4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A22).

General procedure for the syntheses of carbamates, pyridinamides, and sulfonamides. In a 16 mL reaction vessel was placed (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (42 mg, 0.1 mmol), triethylamine (90 μl) and DMF (1.5 mL) was added to completely dissolve the solid material. Isopropyl chloroformate (0.15 mL, 1.0 M in toluene) was added to the solution and the mixture was stirred 30 min under N₂ at room temperature. After all of the starting amine was completely converted as indicated by LCMS, the reaction was stopped by quenching with water. The reaction mixture was concentrated under vacuum and purified by preparative HPLC to give Compound A22, ¹H NMR (CDCl₃, 400MHz) δ 1.26 (d, 6H), 1.89 (m, 2H), 1.93 (m, 2H), 3.07 (s, 3H), 3.63 (m, 2H), 3.67 (m, 2H), 4.94 (m, 1H), 5.61 (m, 1H), 7.87 (d, 2H), 7.97 (d, 2H), 8.40 (s, 1H), 10.18 (s, 1H). Exact mass calculated for C₂₀H₂₅N₅O₇S 479.15, found 480.4 (MH⁺).

Example 9.23: Preparation of 4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid 2-isopropyl-5-methyl-cyclohexyl ester (Compound A23).

Compound A23 was prepared in a similar manner as described above using (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (15 mg, 0.035 mmol), triethylamine (50 μ l), menthyl chloroformate (10 mg, 0.046 mmol), DMF (0.6 mL), ^1H NMR (CDCl_3 , 400 MHz) δ 0.81 (d, 3H), 0.92 (d, 6H), 1.06 (m, 1H), 1.10 (m, 1H), 1.41 (m, 1H), 1.51 (m, 1H), 1.67 (m, 2H), 1.94 (m, 4H), 2.08 (m, 2H), 3.08 (s, 3H), 3.65 (m, 4H), 4.58 (m, 1H), 5.60 (m, 1H), 7.86 (d, 2H), 7.97 (d, 2H), 8.41 (s, 1H), 10.18 (s, 1H). Exact mass calculated for $\text{C}_{27}\text{H}_{37}\text{N}_5\text{O}_7\text{S}$ 575.24, found 576.4 (MH^+).

Example 9.24: Preparation of {4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-pyridin-3-yl-methanone (Compound A24).

Compound A24 was prepared in a similar manner as described above using (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (16 mg, 0.037 mmol), triethylamine (50 μ l), nicotinoyl chloride (10 mg, 0.046 mmol), DMF (1 mL), ^1H NMR (CDCl_3 , 400 MHz) δ 1.95 (m, 2H), 2.14 (m, 2H), 3.07 (s, 3H), 3.55 (m, 1H), 3.65 (m, 2H), 4.13 (m, 1H), 5.72 (m, 1H), 7.40 (m, 1H), 7.79 (m, 1H), 7.87 (d, 2H), 7.97 (d, 2H), 8.41 (s, 1H), 8.70 (m, 2H), 10.20 (s, 1H). Exact mass calculated for $\text{C}_{22}\text{H}_{22}\text{N}_6\text{O}_6\text{S}$ 498.13, found 499.3 (MH^+).

Example 9.25: Preparation of (2-Chloro-pyridin-3-yl)-{4-[6-(4-methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-methanone (Compound A25).

Compound A25 was prepared in a similar manner as described above using (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (16 mg, 0.037 mmol), triethylamine (50 μ l), 2-chloro-nicotinoyl chloride (10 mg, 0.046 mmol), DMF (1 mL), ^1H NMR (CDCl_3 , 400 MHz) δ 1.91 (m, 1H), 2.00 (m, 1H), 2.14 (m, 2H), 3.08 (s, 3H), 3.36 (m, 1H), 3.65 (m, 2H), 4.13 (m, 1H), 4.31 (m, 1H), 5.72 (m, 1H), 7.36 (m, 1H), 7.69 (m, 1H), 7.87 (d, 2H), 7.97 (d, 2H), 8.41 (m, 1H), 8.47 (m, 1H), 10.20 (s, 1H). Exact mass calculated for $\text{C}_{22}\text{H}_{21}\text{ClN}_6\text{O}_6\text{S}$ 532.09, found 533.3 (MH^+).

Example 9.26: Preparation of {4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-pyridin-2-yl-methanone (Compound A26).

Compound A26 was prepared in a similar manner as described above using (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (16 mg, 0.037 mmol), triethylamine (50 μ l), pyridine-2-carbonyl chloride (10 mg, 0.046 mmol) [Pyridine-2-carbonyl chloride was prepared by refluxing picolinic acid with SOCl_2 for 3 hours and worked up in general method], DMF (1 mL), ^1H NMR (CDCl_3 , 400 MHz) δ 1.95 (m, 2H), 2.13 (m, 2H), 3.07 (s, 3H), 3.65 (m, 2H), 3.79 (m, 2H), 4.13 (m, 1H), 5.72 (m, 1H), 7.37 (m, 1H), 7.67 (m,

1H), 7.81 (m, 1H), 7.87 (d, 2H), 7.97 (d, 2H), 8.41 (s, 1H), 8.60 (m, 2H), 10.19 (s, 1H). Exact mass calculated for $C_{22}H_{22}N_6O_6S$ 498.13, found 499.3 (MH^+).

5 **Example 9.27: Preparation of (4-Methanesulfonyl-phenyl)-[6-(1-methanesulfonyl-piperidin-4-yloxy)-5-nitro-pyrimidin-4-yl]-amine (Compound A27).**

Compound A27 was prepared in a similar manner as described above using (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (15 mg, 0.035 mmol), triethylamine (50 μ l), methanesulfonyl chloride (10 mg, 0.087 mmol), DMF (1 mL), 1H NMR ($CDCl_3$, 400 MHz) δ 2.13 (m, 4H), 2.85 (s, 3H), 3.08 (s, 3H), 3.31 (m, 2H), 3.57 (m, 10 2H), 4.13 (m, 1H), 5.69 (m, 1H), 7.87 (d, 2H), 7.98 (d, 2H), 8.42 (s, 1H), 10.21 (s, 1H). Exact mass calculated for $C_{17}H_{21}N_5O_7S_2$ 471.09, found 472.3 (MH^+).

Example 9.28: Preparation of (4-Methanesulfonyl-phenyl)-[5-nitro-6-[1-(propane-1-sulfonyl)-piperidin-4-yloxy]-pyrimidin-4-yl]-amine (Compound A28).

15 Compound A28 was prepared in a similar manner as described above using (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (15 mg, 0.035 mmol), triethylamine (20 μ l), propane-1-sulfonyl chloride (8 mg, 0.056 mmol), DMF (0.6 mL), 1H NMR ($CDCl_3$, 400 MHz) δ 1.09 (t, 3H), 1.90 (m, 2H), 2.07 (m, 4H), 2.95 (m, 2H), 3.08 (s, 3H), 3.40 (m, 2H), 3.57 (m, 2H), 5.67 (m, 1H), 7.87 (d, 2H), 7.98 (d, 2H), 8.41 (s, 1H), 10.21 (s, 20 1H). Exact mass calculated for $C_{19}H_{25}N_5O_7S_2$ 499.12, found 500.3 (MH^+).

Example 9.29: Preparation of {6-[1-(Butane-1-sulfonyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine (Compound A29).

Compound A29 was prepared in a similar manner as described above using (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (15 mg, 0.035 mmol), triethylamine (20 μ l), butane-1-sulfonyl chloride (8 mg, 0.056 mmol), DMF (0.6 mL), 1H NMR ($CDCl_3$, 400 MHz) δ 0.98 (t, 3H), 1.51 (m, 2H), 1.83 (m, 2H), 2.07 (m, 4H), 2.97 (m, 2H), 3.08 (s, 3H), 3.40 (m, 2H), 3.58 (m, 2H), 5.68 (m, 1H), 7.87 (d, 2H), 7.98 (d, 2H), 8.41 (s, 1H), 10.21 (s, 1H). Exact mass calculated for $C_{20}H_{27}N_5O_7S_2$ 513.14, found 514.4 (MH^+).

30

Example 9.30: Preparation of (4-Methanesulfonyl-phenyl)-[5-nitro-6-[1-(thiophene-2-sulfonyl)-piperidin-4-yloxy]-pyrimidin-4-yl]-amine (Compound A30).

Compound A30 was prepared in a similar manner as described above using (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (15 mg, 0.035 mmol), triethylamine (20 μ l), thiophene-2-sulfonyl chloride (9 mg, 0.049 mmol), DMF (0.6 mL), 1H NMR ($CDCl_3$, 400 MHz) δ 2.07 (m, 4H), 3.08 (s, 3H), 3.14 (m, 2H), 3.41 (m, 2H), 5.53

35

(m, 1H), 7.31 (m, 1H), 7.50 (m, 1H), 7.55 (m, 1H), 7.83 (m, 2H), 7.95 (m, 2H), 8.37 (s, 1H), 10.14 (s, 1H). Exact mass calculated for $C_{20}H_{21}N_5O_7S_3$ 539.06, found 540.2 (MH^+).

Example 9.31: Preparation of (4-Methanesulfonyl-phenyl)-{6-[1-(1-methyl-1H-imidazole-4-sulfonyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-amine (Compound A31).

Compound A31 was prepared in a similar manner as described above using (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (15 mg, 0.035 mmol), triethylamine (20 μ L), 1-methyl-1H-imidazole-4-sulfonyl chloride (9 mg, 0.050 mmol), DMF (0.6 mL), 1H NMR ($CDCl_3$, 400 MHz) δ 2.07 (m, 4H), 3.08 (s, 3H), 3.32 (m, 2H), 3.53 (m, 2H), 3.79 (s, 3H), 5.55 (m, 1H), 7.46 (s, 1H), 7.53 (s, 1H), 7.85 (d, 2H), 7.97 (d, 2H), 8.38 (s, 1H), 10.16 (s, 1H). Exact mass calculated for $C_{20}H_{23}N_7O_7S_2$ 537.11, found 538.4 (MH^+).

Example 9.32: Preparation of {6-[1-(2,4-Dimethyl-thiazole-5-sulfonyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine (Compound A32).

Compound A32 was prepared in a similar manner as described above using (4-methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (i.e., Compound A2) (15 mg, 0.035 mmol), triethylamine (20 μ L), 2,4-dimethyl-thiazole-5-sulfonyl chloride (10 mg, 0.047 mmol), DMF (0.6 mL), 1H NMR ($CDCl_3$, 400 MHz) δ 2.09 (m, 4H), 2.67 (s, 3H), 2.75 (s, 3H), 3.08 (s, 3H), 3.21 (m, 2H), 3.50 (m, 2H), 5.58 (m, 1H), 7.85 (d, 2H), 7.97 (d, 2H), 8.38 (s, 1H), 10.14 (s, 1H). Exact mass calculated for $C_{21}H_{24}N_6O_7S_3$ 568.09, found 569.4 (MH^+).

Example 9.33: Preparation of 4-[5-Cyano-6-(3-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester (Compound A33).

Compound A33 was prepared in a similar manner as described in Example 9.1 as a yellow solid (78%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.48 (s, 9H), 1.80-1.86 (m, 2H), 1.90-1.98 (m, 2H), 3.23 (s, 3H), 3.34-3.40 (m, 2H), 3.73-3.78 (m, 2H), 5.44-5.46 (m, 1H), 7.34-7.37 (m, 2H), 7.92-7.96 (m, 1H), 8.04-8.07 (m, 1H), 8.55 (s, 1H). Exact mass calculated for $C_{22}H_{26}FN_5O_5S$ 491.1, found 492.3 (MH^+).

Example 9.34: Preparation of 4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester (Compound A34).

Compound A34 was prepared in a similar manner as described in Example 9.1 as a yellow solid (287 mg, 93%). 1H NMR 400 MHz $CDCl_3$ δ (ppm): 10.3 (s, NH), 8.69 (t, 1H), 8.45 (s, 1H), 7.78 (t, 2H), 5.60 (m, 1H), 3.64-3.61 (m, 2H), 3.56 (m, 2H), 3.09 (s, 3H), 1.97 (m, 2H), 1.88-1.84 (m, 2H), 1.48 (s, 9H). Exact mass calculated for $C_{21}H_{26}FN_5O_7S$ 511.15, LCMS (ESI) m/z 534.3 ($M+H^++Na$, 100%).

Example 9.35: Preparation of 4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester (Compound A35).

Compound A35 was prepared in a similar manner as described in Example 9.1 as a white solid (1.930 g, 72%). ¹H NMR 400MHz CDCl₃ δ (ppm): 8.51 (s, 1H), 7.96 (d, 2H), 7.86 (d, 2H), 7.37 (s, NH), 5.44 (m, 1H), 3.78-3.73 (m, 2H), 3.40-3.33 (m, 2H), 3.07 (s, 3H), 1.99 (m, 2H), 1.85-1.82 (m, 2H), 1.48 (s, 9H). Exact mass calculated for C₂₂H₂₇N₅O₅S 473.17, LCMS (ESI) m/z 474.1 (M+H⁺, 100%).

Example 9.36: Preparation of 4-[6-(6-Methanesulfonyl-pyridin-3-ylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester (Compound A36).

Compound A36 was prepared in a similar manner as described in Example 9.1 as a yellow solid (1.848 g, 76%). ¹H NMR 400MHz CDCl₃ δ (ppm): 10.2 (s, NH), 8.92 (s, 1H), 8.43 (d, 1H), 8.42 (s, 1H), 8.13 (d, 1H), 5.61 (m, 1H), 3.64-3.61 (m, 2H), 3.56-3.51 (m, 2H), 3.24 (s, 3H), 1.96 (m, 2H), 1.91-1.88 (m, 2H), 1.48 (s, 9H). Exact mass calculated for C₂₀H₂₆N₆O₇S 494.16, LCMS (ESI) m/z 495.1 (M+H⁺, 100%).

Example 9.37: Preparation of 4-[5-Acetyl-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester (Compound A37).

4-Hydroxy-piperidine-1-carboxylic acid tert-butyl ester (3.2 mmol, 633 mg) and sodium hydride (3.2 mmol, 76 mg) were dissolved in N,N-dimethyl acetamide (1.5 mL) and stirred for 30 minutes at room temperature. Subsequently, compound 1-[4-chloro-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-5-yl]-ethanone (0.63 mmol, 207 mg) was added. The reaction was stirred at 70°C for 30 minutes. Progress of the reaction was monitored by thin layer chromatography and LCMS. Sodium hydride was carefully quenched with water and the desired compound was extracted with ethyl acetate. Organic solvents were evaporated in vacuo and purified by flash chromatography (Silica gel 60; 50/50 EtoAc/Hexanes) to afford Compound A37 as a yellow solid (156 mg, 50%). ¹H NMR (400 MHz, CDCl₃) δ (ppm): 12.19 (s, 1H), 8.95 (s, 1H), 8.56 (d, 1H), 8.44 (s, 1H), 8.07 (d, 1H), 5.56 (h, 1H), 3.82 (m, 2H), 3.31 (m, 2H), 3.23 (s, 3H), 2.70 (s, 3H), 2.11 (m, 2H), 1.85 (m, 2H), 1.48 (s, 9H). LCMS (ESI), m/z 492.4 (M+H⁺, 100%).

Example 9.38: Preparation of 4-[5-Amino-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester (Compound A38).

Compound A34 was subjected to hydrogenation conditions, H₂ in the presence of 10% Pd/C and ethyl acetate to provide compound A38 as a yellow solid (503 mg, 89%). ¹H NMR 400MHz CDCl₃ δ (ppm): 8.63 (t, 1H), 8.18 (s, 1H), 7.72 (d, 1H), 7.69 (d, 1H), 7.16 (s, NH), 5.32 (m, 1H), 3.82 (m, 2H), 3.30-3.24 (m, 2H), 3.05 (s, 3H), 2.03 (m, 2H), 1.76 (m, 2H), 1.48 (s, 9H). Exact mass calculated for C₂₁H₂₈F N₅O₅S 481.18, LCMS (ESI) m/z 482.3 (M+H⁺, 100%).

Example 9.39: Preparation of 4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A39).

Compound A39 was obtained in a similar manner as described in Example 9.10 as a solid (80%). ¹H NMR (CDCl₃, 400 MHz) δ 1.26 (d, 6H), 1.82-1.86 (m, 2H), 1.90-1.99 (m, 2H), 3.07 (s, 3H), 3.39-3.45 (m, 2H), 3.76-3.82 (m, 2H), 4.94 (sept, 1H), 5.44-5.48 (m, 1H), 7.37 (s, 1H), 7.85-7.87 (m, 2H), 7.95-7.97 (m, 2H), 8.52 (s, 1H). Exact mass calculated for C₂₁H₂₅N₅O₅S 459.2, found 460.2 (MH⁺).

Example 9.40: Preparation of 4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid ethyl ester (Compound A40).

Compound A40 was obtained in a similar manner as described in Example 9.10 as a solid (75%). ¹H NMR (CDCl₃, 400 MHz) δ 1.28 (t, 3H), 1.82-1.86 (m, 2H), 1.90-1.99 (m, 2H), 3.07 (s, 3H), 3.39-3.45 (m, 2H), 3.76-3.82 (m, 2H), 4.16 (q, 2H), 5.44-5.48 (m, 1H), 7.37 (s, 1H), 7.85-7.87 (m, 2H), 7.95-7.97 (m, 2H), 8.52 (s, 1H). Exact mass calculated for C₂₀H₂₃N₅O₅S 445.1, found 446.2 (MH⁺).

Example 9.41: Preparation of 4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isobutyl ester (Compound A41).

Compound A41 was obtained in a similar manner as described in Example 9.10 as a solid (76%). ¹H NMR (CDCl₃, 400 MHz) δ 0.95 (d, 6H), 1.82-1.86 (m, 2H), 1.90-1.99 (m, 2H), 3.07 (s, 3H), 3.42-3.48 (m, 2H), 3.76-3.82 (m, 2H), 3.89 (d, 2H), 5.44-5.48 (m, 1H), 7.37 (s, 1H), 7.85-7.87 (m, 2H), 7.95-7.97 (m, 2H), 8.52 (s, 1H). Exact mass calculated for C₂₂H₂₇N₅O₅S 473.2, found 474.3 (MH⁺).

Example 9.42: Preparation of 4-(4-Methanesulfonyl-phenylamino)-6-[1-(tetrahydro-furan-2-carbonyl)-piperidin-4-yloxy]-pyrimidine-5-carbonitrile (Compound A42).

Compound A42 was obtained in a similar manner as described in Example 9.24 as a solid (75%). ¹H NMR (CDCl₃, 400 MHz) δ 1.87-2.06 (m, 8H), 2.31-2.34 (m, 1H), 3.07 (s, 3H), 3.49-3.50 (m, 1H), 3.74-3.99 (m, 4H), 4.64 (t, 1H), 5.54-5.56 (m, 1H), 7.40-7.42 (m, 1H), 7.85-7.88 (m, 2H), 7.95-7.97 (m, 2H), 8.52 (s, 1H). Exact mass calculated for C₂₂H₂₅N₅O₅S 471.2, found 472.2 (MH⁺).

Example 9.43: Preparation of 4-[1-(3,3-Dimethyl-2-oxo-butyl)-piperidin-4-yloxy]-6-(4-methanesulfonyl-phenylamino)-pyrimidine-5-carbonitrile (Compound A43).

Compound A43 was obtained in a similar manner as described in Example 9.5 as a solid (70%). ¹H NMR (CDCl₃, 400 MHz) δ 1.17 (s, 9H), 1.95-1.99 (m, 2H), 2.00-2.11 (m, 2H), 2.48-2.52 (m, 2H), 2.70-2.75 (m, 2H), 3.07 (s, 3H), 3.48 (s, 2H), 5.44-5.48 (m, 1H), 7.37 (s, 1H), 7.85-7.87 (m,

2H), 7.95-7.97 (m, 2H), 8.52 (s, 1H). Exact mass calculated for $C_{23}H_{29}N_5O_4S$ 471.2, found 472.2 (MH^+).

Example 9.44: Preparation of 4-(4-Methanesulfonyl-phenylamino)-6-[1-(pyridine-3-carbonyl)-piperidin-4-yloxy]-pyrimidine-5-carbonitrile (Compound A44).

Compound A44 was obtained in a similar manner as described in Example 9.24 as a solid (88%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.80-2.14 (m, 4H), 3.07 (s, 3H), 3.40-4.01 (m, 4H), 5.56-5.60 (m, 1H), 7.38-7.44 (m, 2H), 7.79-7.81 (m, 1H), 7.85-7.87 (m, 2H), 7.95-7.97 (m, 2H), 8.52 (s, 1H), 8.70 (s, 1H). Exact mass calculated for $C_{23}H_{22}N_6O_4S$ 478.1, found 479.3 (MH^+).

Example 9.45: Preparation of 4-(1-Formyl-piperidin-4-yloxy)-6-(4-methanesulfonyl-phenylamino)-pyrimidine-5-carbonitrile (Compound A45).

Compound A45 was obtained in a similar manner as described in Example 9.24 as a solid (60%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.93-2.07 (m, 4H), 3.07 (s, 3H), 3.42-3.48 (m, 1H), 3.66-3.76 (m, 3H), 5.56-5.60 (m, 1H), 7.36 (s, 1H), 7.85-7.87 (m, 2H), 7.96-7.98 (m, 2H), 8.13 (s, 1H), 8.53 (s, 1H). Exact mass calculated for $C_{18}H_{19}N_5O_4S$ 401.1, found 402.4 (MH^+).

Example 9.46: Preparation of 4-(4-Methanesulfonyl-phenylamino)-6-[1-(pyridine-2-carbonyl)-piperidin-4-yloxy]-pyrimidine-5-carbonitrile (Compound A46).

Compound A46 was obtained in a similar manner as described in Example 9.24 as a solid (23%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.90-2.14 (m, 4H), 3.07 (s, 3H), 3.46-3.48 (m, 1H), 3.69-3.97 (m, 3H), 5.56-5.60 (m, 1H), 7.47 (s, 1H), 7.54-7.58 (m, 1H), 7.70-7.72 (m, 1H), 7.85-7.87 (m, 2H), 7.95-7.97 (m, 2H), 8.01-8.03 (m, 1H), 8.52 (s, 1H), 8.73-8.74 (m, 1H). Exact mass calculated for $C_{23}H_{22}N_6O_4S$ 478.1, found 479.2 (MH^+).

Example 9.47: Preparation of 4-[5-Cyano-6-(2-fluoro-4-isopropylamino-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A63)

4-[5-Cyano-6-(2-fluoro-4-iodo-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (250 mg, 0.48 mmol), isopropylamine (408 μ L, 4.8 mmol), proline (99 mg, 0.86 mmol), copper iodide (92 mg, 0.48 mmol) and potassium carbonate (152 mg, 1.1 mmol) were mixed together in DMSO (4 mL). The reaction vessel was heated in a microwave at 80°C for 1.0 hour. Progress of the reaction was monitored by TLC and LCMS. Purification by HPLC afforded compound A63 as a white solid (50 mg, 23%). 1H NMR ($CDCl_3$, 400MHz) δ 8.37 (s, 1H), 7.99 (t, 1H), 7.12 (s, 1H), 6.93 (t, 2H), 5.38 (h, 1H), 4.87 (h, 1H), 3.72 (m, 2H), 3.52 (m, 1H), 1.91 (m, 2H), 1.77 (m, 2H), 1.92 (m, 2H), 1.26 (d, 6H), 1.13 (d, 6H). Exact mass calculated for $C_{23}H_{29}FN_6O_3$ 456.51, found 457.1 (MH^+).

Example 9.48: Preparation of 4-[5-Cyano-6-(2-fluoro-4-propylamino-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A64)

4-[5-Cyano-6-(2-fluoro-4-iodo-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (250 mg, 0.48 mmol), *n*-propylamine (408 μ L, 4.8 mmol), proline (99 mg, 0.86 mmol), copper iodide (92 mg, 0.48 mmol) and potassium carbonate (152 mg, 1.1 mmol) were mixed together in DMSO (4 mL). The reaction vessel was heated in a microwave at 80°C for 30 minutes. Progress of the reaction was monitored by TLC and LCMS. Purification by HPLC afforded compound A64 as a white solid (80 mg, 37%). ¹H NMR (CDCl₃, 400MHz) δ 9.37 (s, 1H), 8.22 (s, 1H), 6.91 (t, 1H), 6.91 (m, 2H), 5.27 (h, 1H), 4.71 (h, 1H), 3.53 (m, 2H), 3.23 (m, 2H), 2.91 (m, 2H), 1.85 (m, 2H), 1.54 (m, 4H), 1.13 (d, 6H), 0.88 (t, 3H). Exact mass calculated for C₂₃H₂₉FN₆O₃ 456.51, found 457.4 (MH⁺).

Example 9.49: Preparation of 4-[5-Cyano-6-(2-fluoro-4-propoxy-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A65)

A mixture of 4-[5-cyano-6-(2-fluoro-4-iodo-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (250 mg, 0.48 mmol), propan-1-ol (2 mL, excess), copper iodide (9.1 mg, .048 mmol), 1,10-phenanthroline (18.1 mg, 0.096 mmol) and cesium carbonate (313 mg, 0.96 mmol) in dioxane (3.5 mL) was heated under microwave irradiation for 30 min at 90°C. The crude mixture was concentrated in *vacuo* and purified by HPLC to provide compound A65 as a white solid (10 mg, 12%). Exact mass calculated for C₂₃H₂₈FN₅O₄ 457.50, found 458.8 (MH⁺).

Example 9.50: Preparation of 4-[5-Cyano-6-(6-propyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A66)

In a 25mL round-bottomed flask fitted with a condenser and N₂ inlet was placed 4-[6-(6-chloro-pyridin-3-ylamino)-5-cyano-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (100 mg, 1.3 mmol), *n*-propylzinc bromide (0.5M in THF, 0.72 mL), tetrakis (triphenylphosphino)palladium (28 mg, 0.024 mmol), and THF (3.5 mL). The reaction mixture was refluxed overnight under N₂ atmosphere. The product was purified by preparative HPLC. ¹H NMR (CDCl₃, 400MHz) δ 1.03 (t, 3H), 1.26 (d, 6H), 1.85 (m, 4H), 1.98 (m, 2H), 3.04 (t, 2H), 3.44 (m, 2H), 3.77 (m, 2H), 4.94 (m, 1H), 5.46 (m, 1H), 7.57 (d, 1H), 8.34 (s, 1H), 8.51 (s, 1H), 8.56 (d, 1H), 9.42 (s, 1H). Exact mass calculated for C₂₂H₂₈N₆O₃ 424.22, found 425.2 (MH⁺).

Example 9.51: Preparation of 4-[5-Cyano-6-[4-(2-dimethylamino-ethylsulfanyl)-2-fluoro-phenylamino]-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A67)

In a microwave reaction tube was placed 4-[5-cyano-6-(2-fluoro-4-iodo-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (10 mg, 0.19 mmol), 2-dimethylamino-ethanethiol (27 mg, 0.19 mmol), di-*m*-bromobis (tri-*t*-butylphosphino)dipalladium (I)

(8 mg, 0.0095 mmol), sodium t-butoxide (55 mg, 0.57 mmol), and DMSO (0.5 mL). The reaction mixture was heated at 120°C under microwave for 4 hours. The resulting mixture was filtered through a syringe filter and purified by preparative HPLC. ¹H NMR (CDCl₃, 400MHz) δ 1.26 (d, 6H), 1.86 (m, 2H), 2.00 (m, 2H), 2.86 (s, 6H), 3.20 (m, 2H), 3.30 (m, 2H), 3.43 (m, 2H), 3.78 (m, 2H), 4.94 (m, 1H), 5.44 (m, 1H), 7.22 (s, 1H), 7.24 (s, 1H), 7.35 (s, 1H), 8.11 (t, 1H), 8.45 (s, 1H). Exact mass calculated for C₂₄H₃₁FN₆O₃S 502.22, found 503.2 (MH⁺).

Example 9.52: Preparation of 4-{5-Cyano-6-[4-(2-dimethylamino-ethanesulfonyl)-2-fluoro-phenylamino]-3-oxy-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A68)

In a 50 mL round-bottomed flask immersed in an ice-bath was placed a stir bar, 4-{5-cyano-6-[4-(2-dimethylamino-ethylsulfanyl)-2-fluoro-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (25 mg, 0.04 mmol) and CH₂Cl₂ (15 mL). *m*CPBA (20 mg, 0.089 mmol) dissolved in CH₂Cl₂ (2 mL) was added dropwise at 0°C. The resulting mixture was stirred at 0°C for 1 h and subsequently quenched with sodium bisulfite solution. The organic layer was separated. The aqueous layer was extracted with CH₂Cl₂. The combined organic extracts were dried and concentrated under vacuum to give the crude product. The crude was purified by preparative HPLC. Exact mass calculated for C₂₄H₃₁FN₆O₆S 550.20, found 551.2 (MH⁺).

Example 9.53: Preparation of 4-[5-Cyano-6-(2-fluoro-4-morpholin-4-yl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A71)

A mixture of 4-[5-cyano-6-(2-fluoro-4-iodo-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (60 mg, 0.114 mmol), morpholine (50 μL, 0.571 mmol), CuI (21 mg, 0.114 mmol), proline (23 mg, 0.205 mmol) and potassium carbonate (36 mg, 0.262 mmol) in DMSO (1 mL) was heated in microwave for 30 minutes at 80°C. The mixture was purified by HPLC to give Compound A71 as a solid (25.1 mg, 45%). ¹H NMR (CDCl₃, 400 MHz) δ (ppm): 8.31 (s, 1H), 7.61 (t, 1H), 7.08 (s, 1H), 6.69 (m, 2H), 5.35 (m, 1H), 4.86 (m, 1H), 3.82 (m, 4H), 3.68 (m, 2H), 3.38 (m, 2H), 3.19 (m, 4H), 1.90 (m, 2H), 1.75 (m, 2H), 1.18 (d, 6H). Exact mass calculated for C₂₄H₂₉FN₆O₄ 484.22, found 485.2 (MH⁺).

30

Example 9.54: Preparation of 4-[5-Cyano-6-(4-dimethylamino-2-fluoro-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A73)

Compound A73 was prepared in a similar procedure as described in Example 9.53 as a brownish solid (20 mg, 39.6%). ¹H NMR (CDCl₃, 400 MHz) δ (ppm): 8.41 (s, 1H), 7.97 (t, 1H), 7.37 (s, 1H), 6.99 (m, 2H), 5.42 (m, 1H), 4.92 (m, 1H), 3.73 (m, 2H), 3.44 (m, 2H), 3.09 (s, 6H), 1.95 (m, 2H), 1.85 (m, 2H), 1.23 (d, 6H). Exact mass calculated for C₂₂H₂₇FN₆O₃ 442.49, found 443.3 (MH⁺).

35

Example 9.55: Preparation of 4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A75)

Compound A75 was obtained as a tan solid (HCl salt, 219 mg, 21%). ¹H NMR (MeOH-d₄, 400 MHz) δ 1.17-1.18 (d, 6H), 1.66-1.78 (m, 2H), 1.87-2.01 (m, 2H), 2.12 (s, 3H), 3.10 (s, 3H), 3.18-3.234 (m, 1H), 3.36 (m, 2H), 3.53-3.73 (m, 2H), 5.28-5.39 (m, 1H), 7.73-7.88 (m, 3H), 8.25 (s, 1H). Exact mass calculated for C₂₁H₂₇FN₄O₅S 466.17, found 467.5 (MH⁺).

Example 9.56: Preparation of 4-[6-(2-Fluoro-4-iodo-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A77).

Step 1: Preparation of the 4-(6-Chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester.

To a solution of 4-hydroxy-piperidine-1-carboxylic acid isopropyl ester (6.26 g, 33.4 mmol) and 4,6-dichloro-5-methyl-pyrimidine (5.45 g, 33.4 mmol) in 100 mL THF, 1M potassium *tert*-butoxide in THF (40 mL, 40 mmol) were added slowly by syringe pump. After 1 hour, everything had been added and mixture was concentrated. Residue was extracted with methylene chloride and water. Organic phases were dried over magnesium sulfate, filtered, and concentrated to give 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester as a pale yellow solid (10.3 g, 98%). ¹H NMR (CDCl₃, 400 MHz) δ 1.22-1.24 (d, 6H), 1.74-1.81 (m, 2H), 1.95-2.04 (m, 2H), 2.24 (s, 3H), 3.40-3.45 (m, 2H), 3.74-3.81 (m, 2H), 4.90-4.98 (m, 1H), 5.31-5.37 (m, 1H), 8.40 (s, 1H).

Step 2: Preparation of 4-[6-(2-Fluoro-4-iodo-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A77).

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (2.58 g, 8.22 mmol), palladium acetate (185 mg, 0.82 mmol), biphenyl-3-yl-di-*tert*-butyl-phosphane (25 mg, 0.08 mmol), sodium *tert*-butoxide (2.4 g, 21.2 mmol), and 4-iodo-2-fluoro aniline (2.0 g, 8.4 mmol) in 15 mL dioxane was heated in microwave for 1 hour at 120°C. Solids were filtered off and mixture was purified by column chromatography and precipitating out of hexane/AcOEt to give compound A77 as a tanned solid (1.99 g, 47%). ¹H NMR (CDCl₃, 400 MHz) δ 1.15-1.16 (d, 6H), 1.61-1.71 (m, 2H), 1.85-1.90 (m, 2H), 1.99 (s, 3H), 3.27-3.33 (m, 2H), 3.63-3.66 (m, 2H), 4.82-4.85 (m, 1H), 5.20-5.23 (m, 1H), 6.35-6.36 (d, 1H), 7.33-7.36 (m, 2H), 8.08-8.13 (m, 1H), 8.22 (s, 1H). Exact mass calculated for C₂₀H₂₄F₁N₄O₃ 514.09, found 515.2 (MH⁺).

Example 9.57: Preparation of 4-[6-(2-Fluoro-4-morpholin-4-yl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A79)

Compound A79 was obtained in a similar manner as described in Example 9.47 as a white solid (HCl salt, 401 mg, 38%). ¹H NMR (MeOH-d₄, 400 MHz) δ 1.03-1.05 (d, 6H), 1.53-1.68 (m, 3H), 1.79-1.88 (m, 2H), 1.98 (s, 3H), 3.05-3.09 (m, 3H), 3.15-3.25 (m, 2H), 3.49-3.57 (m, 3H), 3.62-

3.65 (m, 4H), 4.69-4.63 (m, 1H), 5.24-5.28 (m, 1H), 6.74-6.80 (m, 1H), 7.08-7.12 (m, 1H), 8.06 (s, 1H). Exact mass calculated for $C_{25}H_{32}FN_5O_4$ 473.24, found 474.7 (MH^+).

Example 9.58: Preparation of 4-[6-(2,5-Difluoro-4-propoxy-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A80)

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (330 mg, 1.05 mmol), palladium acetate (23.6 mg, 0.01 mmol), biphenyl-3-yl-di-tert-butyl-phosphane (4 mg, 0.013 mmol), sodium tert-butoxide (330 mg, 3.43 mmol), and 2,5-difluoro-4-propoxy-phenylamine (HCl salt, 235 mg, 1.05 mmol) in 15 mL dioxane was heated in microwave for 1 hour at 120°C. Mixture was purified by HPLC and treated with THF to give compound A80 as a white solid (HCl salt, 140 mg, 27%). 1H NMR ($CDCl_3$, 400 MHz) δ 0.88-0.92 (t, 3H), 1.08-1.09 (d, 6H), 1.62-1.73 (m, 4H), 1.83-1.91 (m, 2H), 2.02 (s, 3H), 3.22-3.30 (m, 2H), 3.53-3.60 (m, 2H), 3.88-3.91 (t, 2H), 4.70-4.74 (m, 1H), 5.29-5.30 (m, 1H), 6.99-7.04 (m, 1H), 7.10-7.15 (m, 1H), 8.12 (s, 1H). Exact mass calculated for $C_{23}H_{30}F_2N_4O_4$ 464.22, found 465.4 (MH^+).

15

Example 9.59: Preparation of 4-[6-(2-Fluoro-4-propylamino-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A81)

A mixture of 4-[6-(2-fluoro-4-iodo-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (100 mg, 196 mmol), L-proline (45 mg, 0.39 mmol), copper iodide (37.6 mg, 0.198 mmol), and propylamine (321 μ l, 3.91 mmol) in 4 mL DMSO was heated in microwave for 1 hour at 80°C. Mixture was purified by HPLC to give Compound A81 as a white solid (TFA salt, 108.6 mg, 99%). 1H NMR ($MeOH-d_4$, 400 MHz) δ 0.80-0.84 (t, 3H), 1.06-1.07 (d, 6H), 1.44-1.50 (m, 2H), 1.57-1.62 (m, 2H), 1.79-1.97 (m, 2H), 1.97 (s, 3H), 2.90-2.93 (m, 2H), 3.20-3.29 (m, 2H), 3.54-3.58 (m, 2H), 4.68-4.72 (m, 1H), 5.23-5.26 (m, 1H), 6.37-6.41 (m, 1H), 6.96-7.00 (m, 1H), 8.01 (s, 1H). Exact mass calculated for $C_{22}H_{32}FN_5O_3$ 445.25, found 446.3 (MH^+).

25

Example 9.60: Preparation of 4-[6-[2-Fluoro-4-(2-methoxy-ethylamino)-phenylamino]-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A82)

A mixture of 4-[6-(2-fluoro-4-iodo-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (100 mg, 196 mmol), L-proline (45 mg, 0.39 mmol), copper iodide (37.6 mg, 0.198 mmol), and 2-methoxyethylamine (340 μ l, 3.91 mmol) in 4 mL DMSO was heated in microwave for 1 hour at 80°C. Mixture was purified by HPLC to give Compound A82 as a white solid (TFA salt, 101.7 mg, 90%). 1H NMR ($MeOH-d_4$, 400 MHz) δ 1.21-1.22 (d, 6H), 1.71-1.78 (m, 2H), 1.96-2.01 (m, 2H), 2.13 (s, 3H), 3.34 (s, 3H), 3.34-3.45 (m, 2H), 3.53-3.56 (t, 2H), 3.70-3.73 (m, 2H), 4.81-4.87 (m, 1H), 5.38-5.42 (m, 1H), 6.64-6.57 (m, 1H), 7.09-7.13 (m, 1H), 8.17 (s, 1H). Exact mass calculated for $C_{22}H_{32}FN_5O_3$ 461.24, found 462.4 (MH^+).

35

Example 9.61: Preparation of 4-(6-{2-Fluoro-4-[(tetrahydro-furan-2-ylmethyl)-amino]-phenylamino}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (Compound A83)

A mixture of 4-[6-(2-fluoro-4-iodo-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (100 mg, 196 mmol), L-proline (45 mg, 0.39 mmol), copper iodide (37.6 mg, 0.198 mmol), and C-(tetrahydro-furan-2-yl)-methylamine (404 μ l, 3.91 mmol) in 4 mL DMSO was heated in microwave for 1 hour at 80°C. Mixture was purified by HPLC to give Compound A83 as a white solid (TFA salt, 119 mg, 100%). ^1H NMR (MeOH- d_4 , 400 MHz) δ 1.51-1.53 (d, 6H), 1.90-2.10 (m, 3H), 2.12-2.38 (m, 5H), 2.43 (s, 3H), 3.40-3.49 (m, 2H), 3.75-3.83 (m, 2H), 3.98-4.06 (m, 3H), 4.10-4.19 (m, 1H), 4.30-4.38 (m, 1H), 5.13-5.17 (m, 1H), 5.69-5.72 (m, 1H), 6.85-6.87 (m, 2H), 7.37-7.41 (m, 1H), 8.47 (s, 1H). Exact mass calculated for $\text{C}_{22}\text{H}_{32}\text{FN}_5\text{O}_3$ 487.26, found 488.3 (MH^+).

Example 9.62: Preparation of 4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethylamino)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A84)

A mixture of 4-[6-(2-fluoro-4-iodo-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (100 mg, 196 mmol), L-proline (45 mg, 0.39 mmol), copper iodide (37.6 mg, 0.198 mmol), and 2-methanesulfonyl-ethylamine (307 μ l, 2.5 mmol) in 4 mL DMSO was heated in microwave for 1 hour at 80°C. Mixture was purified by HPLC to give Compound A84 as a white solid (TFA salt, 52.9 mg, 44%). ^1H NMR (MeOH- d_4 , 400 MHz) δ 1.17-1.18 (d, 6H), 1.68-1.72 (m, 2H), 1.91-1.95 (m, 2H), 2.08 (s, 3H), 2.93 (s, 3H), 3.28-3.37 (m, 4H), 3.58-3.67 (m, 4H), 4.78-4.82 (m, 1H), 5.32-5.36 (m, 1H), 6.48-6.53 (s, 2H), 7.06-7.10 (m, 1H), 8.11 (s, 1H). Exact mass calculated for $\text{C}_{23}\text{H}_{32}\text{FN}_5\text{O}_5\text{S}$ 509.21, found 510.5 (MH^+).

Example 9.63: Preparation of 4-(6-{2-Fluoro-4-[(2-methanesulfonyl-ethyl)-methyl-amino]-phenylamino}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (Compound A85)

A mixture of 4-[6-(2-fluoro-4-iodo-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (100 mg, 196 mmol), L-proline (45 mg, 0.39 mmol), copper iodide (37.6 mg, 0.198 mmol), and (2-Methanesulfonyl-ethyl)-methyl-amine (268 μ l, 1.95 mmol) in 4 mL DMSO was heated in microwave for 3 hours at 80°C and for 2 hours at 90°C. Mixture was purified by HPLC to give Compound A85 as a white solid (TFA salt, 22.4 mg, 18%). ^1H NMR (MeOH- d_4 , 400 MHz) δ 1.04-1.06 (d, 6H), 1.51-1.62 (m, 2H), 1.78-1.89 (m, 2H), 1.95 (s, 3H), 2.79 (s, 3H), 2.82 (s, 3H), 3.19-3.23 (m, 4 H), 3.50-3.60 (m, 2H), 3.68-3.72 (t, 2H), 4.66-4.70 (m, 1H), 5.19-5.22 (m, 1H), 6.50-6.53 (m, 2H), 7.01-7.06 (m, 1H), 7.96 (s, 1H). Exact mass calculated for $\text{C}_{24}\text{H}_{34}\text{FN}_5\text{O}_5\text{S}$ 523.23, found 524.4 (MH^+).

Example 9.64: Preparation of 4-[6-(4-Bromo-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A86)

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (1.03 g, 3.28 mmol), palladium acetate (74 mg, 0.33 mmol), biphenyl-3-yl-di-tert-butyl-phosphane (9.7 mg, 0.033 mmol), sodium tert-butoxide (708 mg, 7.36 mmol), and 4-bromo-2,5-difluoro-phenylamine (706 mg, 3.39 mmol) in 15 mL dioxane was heated in microwave for 1 hour at 120°C. Solids were filtered off and mixture was purified by column chromatography (hexane/AcOEt) and crystallized from hexane/AcOEt to give compound A86 as a tanned solid (652 mg, 41%). ¹H NMR (CDCl₃, 400 MHz) δ 1.04-1.05 (d, 6H), 1.50-1.61 (m, 2H), 1.74-1.82 (m, 2H), 1.89 (s, 3H), 3.16-3.22 (m, 2H), 3.51-3.60 (m, 2H), 4.69-4.76 (m, 1H), 5.09-5.15 (m, 1H), 6.34-6.36 (m, 1H), 7.07-7.11 (m, 1H), 8.15 (s, 1H), 8.34-8.38 (m, 1H). Exact mass calculated for C₂₀H₂₃BrF₂N₄O₃ 484.09, found 485.2 (MH⁺).

Example 9.65: Preparation of 4-[6-(4-Cyano-2-fluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A87)

Compound A87 was obtained as a tanned solid (TFA salt, 387.1 mg, 28%). ¹H NMR (MeOH-d₄, 400 MHz) δ 1.118-1.221 (d, J = 6.32 Hz, 6H), 1.608-1.724 (m, 2H), 1.859-1.966 (m, 2H), 2.064 (s, 3H), 3.289-3.404 (m, 2H), 3.607-3.727 (m, 2H), 4.73-4.82 (m, 1H), 5.220-5.310 (m, 1H), 7.409 (d, 1H), 7.545 (d, 1H), 7.954-8.031 (t, J = 8.08 Hz, 1H), 8.145 (s, 1H). Exact mass calculated for C₂₁H₂₄FN₅O₃ 413.19, found 414.4 (MH⁺).

Example 9.66: Preparation of 4-[6-(4-Cyano-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A88)

Compound A88 was obtained as a white solid (TFA salt, 309.8 mg, 22%). ¹H NMR (DMSO-d₆, 400 MHz) δ 1.18-1.20 (d, J = 6.32 Hz, 6H), 1.57-1.66 (m, 2H), 1.89-1.94 (m, 2H), 2.1 (s, 3H), 3.30-3.35 (m, 2H), 3.59-3.65 (m, 1H), 4.73-4.82 (m, J = 6.32 Hz, 2H), 5.24-5.30 (m, J = 3.79 Hz, 1H), 7.88-7.93 (dd, J = 11.37, 6.57 Hz, 1H), 7.93-7.98 (dd, J = 10.36, 6.06 Hz, 1H), 8.31 (s, 1H), 8.72 (s, 1H). Exact mass calculated for C₂₁H₂₃F₂N₅O₃ 431.18, found 432.3 (MH⁺).

Example 9.67: Preparation of 4-[6-(2,5-Difluoro-4-morpholin-4-yl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A89)

A mixture of 4-[6-(4-bromo-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (645 mg, 1.33 mmol), L-proline (306 mg, 2.66 mmol), copper iodide (253 mg, 1.33 mmol), potassium carbonate (211 mg, 1.53 mmol), and morpholine (2.3 mL, 26 mmol) in 15 mL DMSO was heated in microwave for 18 hours at 80°C. Mixture was purified by HPLC to give Compound A89 as a tanned solid (HCl salt, 251 mg, 30%). ¹H NMR (MeOH-d₄, 400 MHz) δ 1.05-1.07 (d, 6H), 1.52-1.63 (m, 2H), 1.80-1.89 (m, 2H), 1.99 (s, 3H), 2.94-2.96 (m, 4H),

3.21-3.29 (m, 2H), 3.54-3.70 (m, 6H), 5.22-5.29 (m, 1H), 6.82-6.86 (m, 1H), 7.03-7.08 (m, 1H), 8.10 (s, 1H). Exact mass calculated for $C_{24}H_{31}F_2N_5O_4$ 491.23, found 492.5 (MH^+).

Example 9.68: Preparation of 4-[6-(6-Chloro-2-methyl-pyridin-3-ylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A90)

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (1.546 g, 4.93 mmol), palladium acetate (110 mg, 0.49 mmol), biphenyl-3-yl-di-tert-butyl-phosphane (18.5 mg, 0.062 mmol), sodium tert-butoxide (1.20 g, 12.5 mmol), and 6-chloro-2-methyl-pyridin-3-ylamine (709 mg, 4.97 mmol) in 15 mL dioxane was heated in microwave for 2 hour at 120°C. Solids were filtered off and mixture was purified to give Compound A90 as a tanned solid (640 mg, 31%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.11-1.12 (d, 6H), 1.52-1.62 (m, 2H), 1.81-1.89 (m, 2H), 1.98 (s, 3H), 2.23 (m, 3H), 3.21-3.30 (m, 2H), 3.59-3.70 (m, 3H), 5.14-5.17 (m, 1H), 6.83-6.91 (m, 1H), 7.14-7.16 (d, 1H), 7.55-7.57 (d, 1H), 7.87 (s, 1H). Exact mass calculated for $C_{20}H_{26}ClN_5O_3$ 419.17, found 419.9 (MH^+).

Example 9.69: Preparation of 4-[5-(4,5-Dihydro-1H-imidazol-2-yl)-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A92)

To a solution of zinc chloride (28 mg, 0.149 mmol) and 4-[5-cyano-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (1 g, 2.09 mmol) in chlorobenzene (15 mL), ethane-1,2-diamine (0.100 mL, 1.463 mmol) was added. The mixture was heated under reflux for 24h. LCMS indicated desired product. The crude was concentrated under *vacuo* and purified by HPLC to afford compound A92 as a yellow solid (303 mg, 23%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.23 (d, 6H), 1.68-1.77 (m, 2H), 2.05-2.09 (m, 2H), 3.07 (s, 3H), 3.16-3.23 (m, 2H), 3.84-3.92 (m, 2H), 4.07 (s, 4H), 4.87-4.92 (m, 1H), 5.42-5.47 (m, 1H), 7.50-7.62 (m, 2H), 7.79-7.83 (m, 1H), 8.35 (s, 1H). Exact mass calculated for $C_{23}H_{29}FN_6O_5S$ 520.19, found 521.5 (MH^+).

Example 9.70: Preparation of (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidin-4-yl}-amine (Compound A93)

A mixture of (6-chloro-5-methyl-pyrimidin-4-yl)-(2-fluoro-4-methanesulfonyl-phenyl)-amine (HCl salt, 1.76 g, 5.0 mmol) and 1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-ol (1.05 g, 5.0 mmol) in anhydrous THF (10 mL) was treated with potassium *t*-butoxide (20 mL, 20 mmol), placed under inert atmosphere, and refluxed for 4 hours, at which point the reaction had stalled at 60% conversion. The reaction mixture was cooled, quenched with water (30 mL), and extracted with ether (2 X 50 mL). The combined organic extract was rinsed with water (20 mL), followed by brine (20 mL), and was dried over $MgSO_4$. After solvent removal, the residue was rinsed with boiling ether (2

X 20 mL), and the combined rinses were set aside to cool. Crystallization yielded a white solid (91% pure by LCMS) which was triturated in hot ether and filtered hot to furnish Compound A93 as a white solid in >95% purity (731 mg, 30% yield). This material was taken up in CH₂Cl₂ (10 mL), to which was added 1 N HCl/ether (1.5 mL). Upon solvent removal, a light gray foam was obtained (800 mg):
5 ¹H NMR (DMSO-*d*₆) δ 10.26 (brs, 1 H), 8.77 (s, 1 H), 8.22 (s, 1 H), 7.83-7.71 (m, 3 H), 5.33 (m, 1 H), 3.75 (m, 2 H), 3.57 (m, 2 H), 3.37 (s, 3 H), 2.83 (m 1 H), 2.13 (s, 3 H), 2.04 (m, 2 H), 1.78 (m, 2 H), 1.20 (d, 6 H, *J* = 6.9 Hz), MS *m/z* 491.2 (M⁺).

10 **Example 9.71: Preparation of 4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A95)**

General procedure of coupling alcohol to aryl halides: A mixture of 4-[6-(2-fluoro-4-iodo-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (103 mg, 0.2 mmole), cesium carbonate (130 mg, 0.4 mmole), copper iodide (8 mg, 0.04 mmole), and 1, 10-phenanthroline (14 mg, 0.08 mmole) in 2-methanesulfonyl-ethanol (3 mL) was heated under
15 microwave irradiation at 150°C for 1 hour. The crude mixture was purified by HPLC to provide Compound A95 as a yellow solid (3 mg, 3%). Exact mass calculated for C₂₃H₃₁FN₄O₆S 510.2, found 511.3 (MH⁺).

20 **Example 9.72: Preparation of 4-[6-(2-Fluoro-4-propoxy-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A94)**

Compound A94 was obtained in a similar manner as described in Example 9.71 as a solid (38 mg, 84%). ¹H NMR (CDCl₃, 400 MHz) δ 1.04 (t, 3H), 1.26 (d, 6H), 1.75-1.84 (m, 7H), 1.97-2.02 (m, 2H), 3.35-3.41 (m, 2H), 3.74-3.77 (m, 2H), 3.91 (t, 2H), 4.93 (sept, 1H), 5.37-5.40 (m, 1H), 6.67-6.72 (m, 2H), 7.27-7.30 (m, 1H), 8.32 (s, 1H), 9.30 (s, 1H). Exact mass calculated for C₂₃H₃₁FN₄O₄ 446.2, found 447.3 (MH⁺).
25

Example 9.73: Preparation of 4-{6-[2-Fluoro-4-(2-methoxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A96)

Compound A96 was obtained in a similar manner as described in Example 9.71 as a tan solid
30 (76 mg, 83%). ¹H NMR (CDCl₃, 400 MHz) δ 1.25 (d, *J* = 6.3 Hz, 6H), 1.74-1.79 (m, 2H), 1.81 (s, 3H), 1.97-2.05 (m, 2H), 3.35-3.41 (m, 2H), 3.45 (s, 3H), 3.75-3.77 (m, 4H), 4.10-4.12 (m, 2H), 4.93 (sep, *J* = 6.3 Hz, 1H), 5.36-5.41 (m, 1H), 6.72-6.75 (m, 2H), 7.36 (t, *J* = 9.1 Hz, 1H), 8.31 (s, 1H), 9.15 (s, NH). Exact mass calculated for C₂₃H₃₁FN₄O₅ 462.2, found 463.5 (MH⁺).

35 **Example 9.74: Preparation of 4-{6-[2-Fluoro-4-(2-isopropoxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A97)**

Compound A97 was obtained in a similar manner as described in Example 9.71 as a yellow solid (86 mg, 88%). ¹H NMR (CDCl₃, 400 MHz) δ 1.21 (d, J = 6.1 Hz, 6H), 1.25 (d, J = 6.3 Hz, 6H), 1.75-1.79 (m, 2H), 1.80 (s, 3H), 1.97-2.02 (m, 2H), 3.35-3.42 (m, 2H), 3.70 (sep, J = 6.3 Hz, 1H), 3.76 (dd, J = 4.0 Hz, 4.8 Hz, 4H), 4.09 (t, J = 4.8 Hz, 2H), 4.93 (sep, J = 6.3 Hz, 1H), 5.37-5.41 (m, 1H), 6.73 (dd, J = 11.6 Hz, 2H), 7.26 (t, J = 8.6 Hz, 1H), 8.32 (s, 1H), 9.36 (s, NH). Exact mass calculated for C₂₅H₃₅FN₄O₅ 490.3, found 491.4 (MH⁺).

Example 9.75: Preparation of 4-[6-(6-Chloro-4-methyl-pyridin-3-ylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A98)

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (1.80 g, 5.74 mmol), palladium acetate (155 mg, 0.69 mmol), biphenyl-3-yl-di-tert-butyl-phosphane (21.5 mg, 0.072 mmol), sodium tert-butoxide (1.38 g, 14.4 mmol), and 6-chloro-4-methyl-pyridin-3-ylamine (838 mg, 5.80 mmol) in 20 mL dioxane was heated in microwave for 1 hour at 120°C. Solids were filtered off and mixture was purified by column chromatography (hexane/AcOEt) to give Compound A98 as a tanned solid (702 mg, 29%). ¹H NMR (CDCl₃, 400 MHz) δ 1.24-1.26 (d, 6H), 1.72-1.81 (m, 2H), 1.95-2.02 (m, 2H), 2.10 (s, 3H), 2.27 (s, 3H), 3.37-3.43 (m, 2H), 3.74-3.77 (m, 2H), 4.90-4.97 (m, 1H), 5.29-5.34 (m, 1H), 5.91 (s, 1H), 7.00 (s, 1H), 8.22 (s, 1H), 8.57 (s, 1H). Exact mass calculated for C₂₀H₂₆ClN₅O₃ 419.17, found 420.4 (MH⁺).

Example 9.76: Preparation of 4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-(N-hydroxycarbamimidoyl)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A99)

4-[5-Cyano-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (0.5g, 1.04 mmol) was dissolved in the mixture ethanol/water (30mL/14mL) and heated to 80°C. Hydroxylamine hydrochloride (7.22g, 104 mmol) and potassium carbonate (14.5 g, 105 mmol) were slowly added and the mixture kept stirring at 80°C for 1h. The crude was filtered and the retrieved solid was thoroughly washed with acetonitrile. The filtrate was concentrated under reduced pressure, yielding a yellow solid residue, which was purified by HPLC to afford Compound A99 (0.51g, 78%). Exact mass calculated for C₂₁H₂₇FN₆O₆S 510.17, found 511.2 (MH⁺).

Example 9.77: Preparation of 4-[5-Carbamimidoyl-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A100)

4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-(N-hydroxycarbamimidoyl)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (0.510 g, 0816 mmol) was dissolved in acetic acid glacial (20 mL) and zinc dust (1 g, 16.32 mmol) was added. The reaction mixture was heated at 70°C

for 40 min. The crude was filtered, the filtrate was concentrated under reduced pressure and the residue was purified by HPLC to afford Compound A100 (43 mg, 8.65%). Exact mass calculated for $C_{21}H_{27}FN_6O_5S$ 494.17, found 495.5 (MH^+).

5 **Example 9.78: Preparation of 4-{6-[2-Fluoro-4-(tetrahydro-furan-2-ylmethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A101)**

Compound A101 was obtained in a similar manner as described in Example 9.71 as a solid (35 mg, 24%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.15 (d, 6H), 1.66-1.73 (m, 5H), 1.87-2.02 (m, 6H), 3.27-3.34 (m, 2H), 3.66-3.89 (m, 6H), 4.19-4.21 (m, 1H), 4.85 (sept, 1H), 5.28-5.30 (m, 1H), 6.64-10 6.67 (m, 2H), 7.32 (t, 1H), 8.22 (s, 1H), 8.90 (s, 1H). Exact mass calculated for $C_{25}H_{33}FN_4O_5$ 488.2, found 489.5 (MH^+).

Example 9.79: Preparation of 4-{6-[6-(2-Methoxy-ethoxy)-2-methyl-pyridin-3-ylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A103)

15 A mixture of 4-[6-(6-chloro-2-methyl-pyridin-3-ylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (507 mg, 1.21 mmol) and potassium carbonate (1.62 g, 12 mmol) in 4.5 mL 2-methoxyethanol was heated in microwave for 16.5 hours at 180°C. Mixture was purified by HPLC to give Compound A103 as a tanned solid (HCl salt, 103.5 mg, 17%). 1H NMR (MeOH- d_4 , 400 MHz) δ 1.15-1.17 (d, 6H), 1.68-1.74 (m, 2H), 1.92-1.96 (m, 2H), 2.12 (s, 3H), 2.44 20 (s, 3H), 3.28-3.37 (m, 5H), 3.64-3.70 (m, 4H), 4.43-4.46 (m, 2H), 4.74-4.79 (m, 1H), 5.35-5.39 (m, 1H), 6.92-6.94 (d, 1H), 7.70-7.73 (d, 1H), 8.16 (s, 1H). Exact mass calculated for $C_{23}H_{33}N_5O_5$ 459.25, found 460.5 (MH^+).

25 **Example 9.80: Preparation of 4-{6-[6-(2-Methoxy-ethoxy)-4-methyl-pyridin-3-ylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A104)**

A mixture of 4-[6-(6-chloro-4-methyl-pyridin-3-ylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (353 mg, 0.84 mmol) and potassium carbonate (1.1 g, 7.96 mmol) in 4 mL 2-methoxyethanol was heated in microwave for 17 hours at 180°C. Mixture was purified by HPLC to give Compound A104 as a tanned solid (HCl salt, 61.8 mg, 15%). 1H NMR 30 (MeOH- d_4 , 400 MHz) δ 1.07-1.08 (d, 6H), 1.60-1.65 (m, 2H), 1.83-1.87 (m, 2H), 2.05 (s, 3H), 2.15 (s, 3H), 3.21-3.32 (m, 5H), 3.54-3.61 (m, 4H), 4.34-4.36 (m, 2H), 4.67-4.73 (m, 1H), 5.27-5.31 (m, 1H), 6.98 (s, 1H), 8.04 (s, 1H), 8.09 (s, 1H). Exact mass calculated for $C_{23}H_{33}N_5O_5$ 459.25, found 460.3 (MH^+).

35 **Example 9.81: Preparation of 4-{6-[2-Fluoro-4-(2-isopropoxy-ethylsulfamoyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A106)**

A mixture of 4-amino-3-fluoro-N-(2-isopropoxy-ethyl)-benzenesulfonamide (116 mg, 0.42 mmol), 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (100 mg, 0.3 mmol), palladium acetate (3 mg, .017 mmol), biphenyl-2-yl-di-tert-butyl-phosphane (7.1 mg, .034 mmol), and sodium *t*-butoxide (87 mg, 0.90 mmol) in dioxane (2 mL) was heated under microwave irradiation for 60 min at 150°C. The crude mixture was concentrated in vacuo and purified by HPLC to provide Compound A106 as a brown solid (50 mg, 22%). ¹H NMR (CDCl₃, 400 MHz) δ 1.04 (d, 6H), 1.19 (d, 6H), 1.67-1.78 (m, 2H), 1.89-1.99 (m, 5H), 3.05 (t, 2H), 3.30-3.40 (m, 4H), 3.42-3.52 (m, 1H), 3.66-3.76 (m, 1H), 4.87 (h, 1H), 5.19-5.38 (m, 2H), 7.58 (t, 3H), 7.90-7.98 (s broad, 1H), 8.24 (t, 1H), 8.35 (s, 1H). Exact mass calculated for C₂₃H₃₆FN₅O₆S 553.65, found 554.6 (MH⁺).

Example 9.82: Preparation of 4-{6-[2,5-Difluoro-4-(N-hydroxycarbamimidoyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A107)
4-[6-(4-Carbamoyl-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A108)

A mixture of 4-[6-(4-cyano-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (TFA salt, 181 mg, 0.332 mmol), hydroxylamine (283.8 mg, 4.08 mmol), and potassium carbonate (283.9 mg, 2.05 mmol) in 6 mL EtOH/H₂O (2:1 v/v) was stirred at 75°C for 45 min. The crude mixture was purified by HPLC to give Compound A107 as an oil (TFA salt, 111 mg, 58%) and Compound A108 as an oil as a by-product (TFA salt, 74 mg, 40%). ¹H NMR of A107 (DMSO-d₆, 400 MHz) δ 1.19-1.20 (d, J = 6.32 Hz, 6H), 1.61-1.63 (m, 2H), 1.88-1.95 (m, 2H), 2.1 (s, 3H), 3.30-3.37 (m, 2H), 3.61-3.63 (m, 2H), 4.75-4.82 (m, J = 6.32 Hz, 1H), 5.25-5.29 (m, J = 3.79 Hz, 1H), 7.64-7.68 (dd, J = 10.36, 6.32 Hz, 1H), 7.72-7.76 (dd, J = 11.62, 6.32 Hz, 1H), 8.23 (s, 1H), 8.66 (s, 1H), 9.11 (s, 1H). Exact mass calculated for A107, C₂₁H₂₆F₂N₆O₄ 464.2, found 465.5 (MH⁺) and for A108, C₂₁H₂₅F₂N₅O₄ 449.19, found 450.3 (MH⁺).

Example 9.83: Preparation of 4-[6-(4-Carbamimidoyl-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A110)

A mixture of Compound A107 (TFA salt, 107.5 mg, 0.186 mmol) and zinc dust (242.6 mg, 3.71 mmol) in acetic acid (3 mL) was stirred at 75°C for 45 min. The crude mixture was purified by HPLC to provide Compound A110 as a solid (TFA salt, 97.4 mg, 93%). ¹H NMR (CDCl₃, 400 MHz) δ 1.24-1.26 (d, J = 6.32 Hz, 6H), 1.76-1.78 (m, 2H), 1.97-1.98 (m, 2H), 2.14 (s, 3H), 3.37-3.43 (m, 2H), 3.75-3.77 (m, 2H), 4.89-4.96 (m, 1H), 5.33-5.37 (m, 1H), 7.16 (s, 1H), 7.48 (s, 1H), 7.59-7.64 (dd, J = 10.61, 6.82 Hz, 1H), 8.36 (s, 1H), 8.62-8.68 (m, 1H), 10.42 (s, 2H). Exact mass calculated for C₂₁H₂₆F₂N₆O₃ 448.2, found 449.2 (MH⁺).

Example 9.84: Preparation of 4-{6-[4-(2-Ethoxy-ethoxy)-2-fluoro-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A111)

Compound A111 was obtained in a similar manner as described in Example 9.71 as a brown solid (52 mg, 55%). ¹H NMR (CDCl₃, 400 MHz) δ 1.25 (t, J = 7.1 Hz, 3H), 1.25 (d, J = 6.3 Hz, 6H), 1.75-1.79 (m, 2H), 1.79 (s, 3H), 1.97-2.02 (m, 2H), 3.35-3.42 (m, 2H), 3.61 (q, J = 7.1 Hz, 2H), 3.75-3.76 (m, 2H), 3.80 (t, J = 4.8 Hz, 2H), 4.11 (t, J = 4.8 Hz, 2H), 4.93 (sep, J = 6.3 Hz, 1H), 5.36-5.40 (m, 1H), 6.72 (d, J = 2.02 Hz, 1H), 6.75 (d, J = 2.02 Hz, 1H), 7.32 (t, J = 8.6 Hz, 1H), 8.30 (s, 1H), 9.41 (s, NH). Exact mass calculated for C₂₄H₃₃FN₄O₅ 476.2, found 477.4 (MH⁺).

Example 9.85: Preparation of 4-{6-[2-Fluoro-4-(tetrahydro-pyran-4-yloxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A112)

Compound A112 was obtained in a similar manner as described in Example 9.71 as an orange solid (71 mg, 49%). ¹H NMR (CDCl₃, 400 MHz) δ 1.26 (d, J = 6.1 Hz, 6H), 1.78-1.86 (m, 4H), 1.90 (s, 3H), 1.99-2.07 (m, 4H), 3.37-3.44 (m, 2H), 3.62-3.68 (m, 2H), 3.76-3.79 (m, 2H), 3.98-4.04 (m, 2H), 4.49 (m, 1H), 4.94 (sep, J = 6.1 Hz, 1H), 5.42-5.44 (m, 1H), 6.72-6.74 (m, 1H), 6.74-6.76 (m, 1H), 7.25 (t, J = 8.8 Hz, 1H), 8.37 (s, 1H), 8.75 (s, NH). Exact mass calculated for C₂₅H₃₃FN₄O₅ 488.2, found 489.5 (MH⁺).

Example 9.86: Preparation of 4-{6-[2-Fluoro-4-(2-hydroxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A113)

Compound A113 was obtained in a similar manner as described in Example 9.71 as an orange solid (76 mg, 84%). ¹H NMR (CDCl₃, 400 MHz) δ 1.25 (d, J = 6.3 Hz, 6H), 1.75-1.79 (m, 2H), 1.84 (s, 3H), 1.97-2.02 (m, 2H), 3.35-3.42 (m, 2H), 3.74-3.78 (m, 2H), 3.98 (t, J = 4.6 Hz, 2H), 4.09 (t, J = 4.6 Hz, 2H), 4.93 (sep, J = 6.3 Hz, 1H), 5.36-5.40 (m, 1H), 6.72-6.74 (m, 1H), 6.75 (s, 1H), 7.35 (t, J = 9.1 Hz, 1H), 8.31 (s, 1H), 9.15 (s, NH). Exact mass calculated for C₂₂H₂₉FN₄O₅ 448.2, found 449.3 (MH⁺).

Example 9.87: Preparation of 4-{6-[2-Fluoro-4-(pyridin-2-ylmethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A117)

Compound A117 was obtained in a similar manner as described in Example 9.71 as a white solid (11 mg, 11%). Exact mass calculated for C₂₆H₃₀FN₅O₄ 495.2, found 496.3 (MH⁺).

Example 9.88: Preparation of 4-[2-(2-Fluoro-4-methanesulfonyl-phenylamino)-3-methyl-pyridin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A118)

Step 1: Preparation of 2,4-dichloro-3-methyl-pyridine.

1.6 M n-Butyl lithium in hexanes (3.75 mL, 6.0 mmol) and anhydrous THF (5 mL) were added to a flame-dried flask under nitrogen atmosphere. This solution was cooled to -78C, 2,4-

Dichloro-pyridine was added dropwise while stirring and the mixture was stirred at -78°C for 30 min after which time methyl iodide (0.374 mL, 6.0 mmol) was added dropwise at -78°C. This mixture was stirred at -78°C for 1 h under nitrogen atmosphere after which time glacial AcOH (0.114 mL, 2.0 mmol) was added to give a reaction mixture pH (wet pH paper) of 5-6. The reaction mixture was dissolved in Et₂O (100 mL), the organic layer was washed with water (10 mL), then brine (10 mL), dried with MgSO₄, and the solvent was evaporated in vacuo to give an oil which was purified by flash chromatography using hexanes:CH₂Cl₂ (50:50 v/v) to hexanes:CH₂Cl₂:EtOAc (50:47:3 v/v/v) to give 2,4-dichloro-3-methyl-pyridine as a white solid (589 mg, 72%). It was noted that 2,4-dichloro-3-methyl-pyridine readily sublimates in vacuo. ¹H NMR (CD₃OD, 400 MHz) δ 8.15 (d, 1H), 7.46 (d, 1H), 2.50 (s, 3H). LRMS calculated for C₆H₅Cl₂N: 160.98, found: (MH)⁺ 161.9.

Step 2: Preparation of 4-(2-chloro-3-methyl-pyridin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester.

4-Hydroxy-piperidine-1-carboxylic acid isopropyl ester (0.496 mL, 2.90 mmol) was dissolved in anhydrous dimethylacetamide (DMA, 5 mL), NaH (60% oil dispersion, 116 mg, 2.90 mmol) was added and this mixture was stirred at 23°C for 45 min, then this mixture was added dropwise to 2,4-dichloro-3-methyl-pyridine, which was dissolved in anhydrous DMA (4 mL). This mixture was stirred at 23°C for 2 h then heated at 50°C for 15 h, after which time the mixture was diluted with Et₂O (140 mL), washed with water (14 mL), then brine twice (2 x 14 mL). The organic layer was separated, dried with MgSO₄, and the solvent was evaporated in vacuo to give an oil which was purified by flash chromatography using hexanes-EtOAc, 75:25, v/v, then hexanes-EtOAc, 50:50, v/v, to give 4-(2-chloro-3-methyl-pyridin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester as a solid (223 mg, 27%). ¹H NMR (CDCl₃, 400 MHz) δ 8.09 (d, 1H), 6.69 (d, 1H), 4.91 (m, 1H), 4.60 (m, 1H), 3.61 (m, 2H), 3.52 (m, 2H), 2.24 (s, 3H), 1.91 (m, 2H), 1.80 (m, 2H), 1.24 (d, 6H). LRMS calculated for C₁₅H₂₁ClN₂O₃: 312.12, Found: (MH)⁺ 313.4.

Step 3: Preparation of 4-[2-(2-fluoro-4-methanesulfonyl-phenylamino)-3-methyl-pyridin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A118).

The free base form of Compound A118 was prepared in a similar manner as described in Example 9.64 with modifications, wherein Pd₂dba₃ was used instead of Pd (OAc)₂, toluene instead of dioxane, and the reaction was heated for 4 h instead of 2 h. Furthermore, no workup was performed and the reaction mixture was applied directly to flash chromatography using hexanes:CH₂Cl₂:EtOAc (10:30:60, v/v/v) to give the free base form of Compound A118 as a solid (166 mg, 51%). ¹H NMR (CDCl₃, 400 MHz) δ 1.24 (d, J = 6.2 Hz, 6H), 1.86 (m, 2H), 2.00 (m, 2H), 2.05 (s, 3H), 3.05 (s, 3H), 3.50 (m, 2H), 3.70 (m, 2H), 4.75 (septet, J = 6.3 Hz, 1H), 4.92 (m, 1H), 6.74 (d, J = 6.1 Hz, 1H), 7.65 (m, 3H), 8.00 (d, J = 6.5 Hz, 1H). LRMS calculated for C₂₂H₂₈FN₃O₅S: 465.17, found: 466.5 (MH)⁺.

Example 9.89: Preparation of 1-[4-(1-Benzyl-azetidin-3-yloxy)-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-5-yl]-ethanone (Compound A61).

Compound A61 was prepared in a manner similar as described in Example 9.37 using 1-benzyl-azetidin-3-ol. Exact mass calculated for $C_{22}H_{23}N_5O_4S$ 453.15, found 489.6 (MH⁺).

Example 9.90: Preparation of 4-[5-Acetyl-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isobutyl ester (Compound A60).

A mixture of 1-[4-(6-methanesulfonyl-pyridin-3-ylamino)-6-(piperidin-4-yloxy)-pyrimidin-5-yl]-ethanone (48 mg, 0.11 mmol), isobutylchloroformate (14.0 μ L, 0.11 mmol), and triethylamine (45 μ L, 0.34 mmol) in DMF (1.0 mL) was heated under microwave irradiation for 3 minutes at 80°C. The crude mixture was purified by HPLC to provide Compound A60 as a white solid (35 mg, 65%).
¹H NMR (CDCl₃, 400 MHz) δ 0.97 (d, 6H), 1.82-1.92 (m, 2H), 2.10-2.19 (m, 2H), 2.70 (s, 3H), 3.22 (s, 3H), 3.37 (m, 2H), 3.89-3.96 (m, 5H), 5.59 (h, 1H), 8.10 (d, 1H), 8.49-8.57 (m, 2H), 8.92 (d, 2H). Exact mass calculated for $C_{22}H_{29}N_5O_6S$ 491.18, found 492.3 (MH⁺).

Example 9.91: Preparation of 4-[5-Methyl-6-(4-methyl-6-morpholin-4-yl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A102).

A solution of 4-[6-(6-chloro-4-methyl-pyridin-3-ylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (223 mg, 0.53 mmol) in 4.5 mL morpholine was reacted under microwave irradiation at 180°C for 16 hours. Mixture was concentrated and purified by HPLC to give Compound A102 as a white solid (200 mg, 74%).
¹H NMR (MeOH-d₄, 400 MHz) δ 1.16-1.18 (d, 6H), 1.64-1.71 (m, 2H), 1.89-1.98 (m, 2H), 2.10 (s, 3H), 2.28 (s, 3H), 3.31-3.38 (m, 2H), 3.61-3.69 (m, 6H), 3.78-3.80 (m, 4H), 4.77-4.82 (m, 1H), 5.28-5.35 (m, 1H), 7.34 (s, 1H), 7.98 (s, 1H), 8.16 (s, 1H). Exact mass calculated for $C_{24}H_{34}N_6O_4$ 470.26, found 471.4 (MH⁺).

Example 9.92: Preparation of 4-[5-Methyl-6-(2-methyl-6-morpholin-4-yl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A91).

A solution of 4-[6-(6-chloro-2-methyl-pyridin-3-ylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (613 mg, 1.46 mmol) in 15 mL morpholine was reacted under microwave irradiation at 180°C for 14 hours. Mixture was concentrated and purified by HPLC to give Compound A91 as a white solid (427 mg, 58%).
¹H NMR (MeOH-d₄, 400 MHz) δ 1.03-1.05 (d, 6H), 1.51-1.60 (m, 2H), 1.78-1.85 (m, 2H), 1.98 (s, 3H), 2.29 (s, 3H), 3.19-3.25 (m, 2H), 3.54-3.58 (m, 6H), 3.65-3.67 (m, 4H), 4.65-4.70 (m, 1H), 5.20-5.25 (m, 1H), 7.08-7.10 (d, 1H), 7.68-7.71 (d, 1H), 8.07 (s, 1H). Exact mass calculated for $C_{24}H_{34}N_6O_4$ 470.26, found 471.3 (MH⁺).

Example 9.93: Preparation of 4-[5-Amino-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound A120).

Mixture of 4-[6-(2-fluoro-4-methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (197 mg, 0.3 mmole), Zn Dust (2.4 mmole, 8 eq) and

1 mL of Sat NH_4Cl solution in 2 mL THF and 2 mL H_2O was stirred at room temperature for 25 minutes. Zn Dust was filtered off by celite and washed with ethyl acetate. Crude was purified by column chromatography (Hexane/Ethyl Acetate = 1/2, R_f = 0.44) to give compound **A120** as a yellow oil (100 mg, 71%). ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ 1.19 (d, 6H), 1.62-1.68 (m, 2H), 1.88-1.93 (m, 2H), 3.23 (s, 3H), 3.33-3.39 (m, 2H), 3.64-3.70 (m, 2H), 4.77 (sep, 1H), 5.28-5.29 (m, 1H), 7.68 (d, 1H), 7.77 (d, 1H), 7.88 (s, 1H), 8.06 (t, 1H), 8.41 (sb, NH). Exact mass calculated for $\text{C}_{20}\text{H}_{26}\text{FN}_5\text{O}_5\text{S}$ 467.2, found 468.5 (MH^+).

Example 9.94: Preparation of 1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butan-1-one (Compound A114).

General Procedure of Amide Formation

Dissolve HBTU (1.2 eq, 24 mg) in DMF (0.5 mL), and add butyric acid (1.2 eq, 5.8 μL) followed by diisopropyl ethyl amine (2.2eq, 20.3 μL). After approximately 3 min, (2-fluoro-4-methanesulfonyl-phenyl)-[5-methyl-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine (0.053 mmol) was added, and stirred at room temperature overnight. Reactions were filtered through 0.1 μm syringe filter and purified by Prep-LCMS. Fractions were frozen and lyophilized to solid product. Exact Mass: 450.2, found: 451.3 (MH^+).

Example 9.95: Preparation of 1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-pentan-1-one (Compound A115).

Compound **A115** was prepared in a similar manner as described in Example 9.94. Exact Mass: 464.2, found: 465.4 (MH^+).

Example 9.96: Preparation of 1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-butan-1-one (Compound A116).

Compound **A116** was prepared in a similar manner as described in Example 9.94. Exact Mass: 464.2, found: 465.6 (MH^+).

Example 9.97: Preparation of 4-{6-[2,5-Difluoro-4-(2-methoxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound A105).

Compound **A105** was obtained in a similar manner as described in Example 9.71 as a solid (TFA salt, 222.5 mg, 16%). ^1H NMR (CDCl_3 , 400 MHz) δ 1.23-1.25 (d, J = 6.32 Hz, 6H), 1.72-1.79 (m, 2H), 1.88 (s, 3H), 1.95-2.00 (m, 2H), 3.34-3.41 (m, 2H), 3.44 (s, 3H), 3.71-3.77 (m, 4H), 4.14-4.16 (m, 2H), 4.87-4.96 (sep, J = 6.32 Hz, 1H), 5.31-5.37 (m, 1H), 6.79-6.84 (dd, J = 11.62, 7.58 Hz, 1H), 7.50-7.55 (dd, J = 11.62, 7.58 Hz, 1H), 8.31 (s, 1H), 8.56 (s, 1H). Exact mass calculated for $\text{C}_{23}\text{H}_{30}\text{F}_2\text{N}_4\text{O}_5$ 480.22, found 481.3 (MH^+).

EXAMPLE 10**SYNTHESES OF COMPOUNDS OF THE PRESENT INVENTION**

Example 10.1: Preparation of 4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-ylamino]-piperidine-1-carboxylic acid tert-butyl ester (Compound B1).

General Procedure for the Addition of Amine to pyrimidine: (6-Chloro-5-nitro-pyrimidin-4-yl)-(4-methanesulfonyl-phenyl)-amine (132 mg, 0.4 mmole), 4-amino-piperidine-1-carboxylic acid tert-butyl ester (0.4 mmole, 1 eq) and K_2CO_3 (0.4 mmole, 1 eq) were dissolved in DMF, and the mixture was stirred at 60°C for 1 hour. Final product was precipitated out with water to provide Compound B1 as a yellow solid (152 mg, 77%). 1H NMR (400 MHz $CDCl_3$) δ (ppm): 10.8 (s, 1H), 9.18 (d, 1H), 8.17 (s, 1H), 7.90 (d, 2H), 7.85 (d, 2H), 4.39-4.32 (m, 1H), 4.02 (m, 2H), 3.01 (s, 3H), 2.95-2.90 (m, 2H), 2.00 (m, 2H), 1.57-1.50 (m, 2H), 1.46 (s, 9H). Exact mass calculated for $C_{21}H_{28}N_6O_6S$ 492.18, LCMS (ESI) m/z 493.4 ($M+H^+$, 100%).

Example 10.2: Preparation of N-(4-Methanesulfonyl-phenyl)-5-nitro-N'-piperidin-4-yl-pyrimidine-4,6-diamine (Compound B2).

General Deprotection Procedure: A mixture of Compound B1 and 4 M HCl in dioxane was stirred at 40°C overnight and concentrated. Excess HCl was evaporated with isopropyl alcohol provided Compound B2 as yellow solid (261 mg, 97%). 1H NMR (400 MHz $CDCl_3$) δ (ppm): 10.9 (s, 1H), 8.96 (d, 2H), 8.17 (s, 1H), 7.84 (d, 4H), 4.40-4.37 (m, 1H), 3.25-3.22 (m, 2H), 3.16 (s, 3H), 3.01-2.93 (m, 2H), 2.04-2.01 (m, 2H), 1.88-1.78 (m, 2H). Exact mass calculated for $C_{16}H_{20}N_6O_4S$ 392.13, LCMS (ESI) m/z 393.1 ($M+H^+$, 100%).

Example 10.3: Preparation of 1-{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-ylamino]-piperidin-1-yl}-ethanone (Compound B3).

General Procedure of Acetylating: Mixture of B2 and acetyl chloride was stirred at 180°C for 2 hours in microwave to provide Compound B3 as yellow solid (10 mg, 18%). 1H (400 MHz $CDCl_3$) δ (ppm): 9.06 (d, 1H), 8.07 (s, 1H), 7.78 (d, 2H), 7.70 (d, 2H), 4.42-4.37 (m, 1H), 4.35-4.30 (m, 1H), 3.74-3.71 (m, 1H), 3.19-3.13 (m, 1H), 2.89 (s, 3H), 2.82-2.76 (m, 1H), 2.04 (s, 3H), 2.00-1.97 (m, 2H), 1.46-1.37 (m, 2H). Exact mass calculated for $C_{18}H_{22}N_6O_5S$ 434.14, LCMS (ESI) m/z 435.4 ($M+H^+$, 100%).

Example 10.4: Preparation of 1-{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-ylamino]-piperidin-1-yl}-2,2-dimethyl-propan-1-one (Compound B4).

Compound B4 was prepared in a similar manner as described above as yellow solid (7 mg, 11%). 1H NMR (400 MHz $CDCl_3$) δ (ppm): 9.16 (d, 1H), 8.17 (s, 1H), 7.89 (d, 2H), 7.82 (d, 2H), 4.48-4.42 (m, 1H), 4.35-4.32 (m, 2H), 3.07-3.04 (m, 2H), 3.00 (s, 3H), 2.10-2.08 (m, 2H), 1.55-1.46 (m,

2H), 1.24 (s, 9H). Exact mass calculated for $C_{21}H_{28}N_6O_5S$ 476.18, LCMS (ESI) m/z 477.3 ($M+H^+$, 100%).

5 **4-({[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yl]-isopropyl-amino}-methyl)-piperidine-1-carboxylic acid tert-butyl ester (Compound B5).**

Compound B5 was prepared in a similar manner as described in Example 10.1 as a solid (24 mg, 23%). Exact mass calculated for $C_{26}H_{38}FN_5O_4S$ 535.2, found 536.4 (MH^+).

10 **EXAMPLE 11**

Syntheses of compounds of the present invention

Example 11.1: Preparation of 4-[6-(2-Fluoro-4-morpholin-4-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C3).

15 A mixture of 4-[6-(4-bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C130, 500 mg, 1.07 mmole), morpholine (121 mg, 1.39 mmole), palladium acetate (3 mg, 0.011 mmole), biphenyl-2-yl-di-tert-butyl-phosphane (4 mg, 0.012 mmole) and sodium *t*-butoxide (257 mg, 2.14 mmole) in dioxane (3 mL) was heated under microwave irradiation at 150°C for 1 hours. The crude mixture was purified by HPLC to provide Compound C3 as a yellow oil (235 mg, 46%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.28 (d, $J = 6.3$ Hz, 6H), 1.81-1.85 (m, 2H), 1.99-2.04 (m, 2H), 2.20 (s, 3H), 3.44-3.47 (m, 2H), 3.49-3.51 (m, 4H), 3.73-3.78 (m, 2H), 4.08-4.10 (m, 2H), 4.95 (sep, $J = 6.3$ Hz, 1H), 5.35-5.37 (m, 1H), 7.25 (d, $J = 10.1$ Hz, 2H), 7.32 (t, $J = 8.6$ Hz, 1H), 8.26 (s, 1H). Exact mass calculated for $C_{24}H_{31}FN_4O_5$ 474.2, found 475.4 (MH^+).

25 **Example 11.2: Preparation of (6-Amino-pyridin-3-yl)-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-methanone (Compound C5).**

6-Amino-nicotinic acid (21.5 mg, 0.155 mmol), HATU (59 mg, 0.155 mmol) and triethylamine (0.05 mL, 0.359 mmol) were mixed in DMF and stirred at rt for 20 min. 4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-(piperidin-4-yloxy)-pyrimidine was then added and the mixture stirred at rt for 2h. The crude was purified by HPLC to afford Compound C5 as a yellow solid (67 mg, 90.7%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.88-1.87 (m, 2H), 1.99-2.01 (m, 2H), 2.15 (s, 3H), 3.03 (s, 3H), 3.58-3.60 (m, 2H), 3.77-3.78 (m, 2H), 5.37-5.41 (m, 1H), 6.78-6.82 (d, 1H), 7.33-7.38 (m, 1H), 7.69-7.74 (m, 2H), 7.87-7.96 (m, 2H), 8.14 (s, 1H). Exact mass calculated for $C_{23}H_{24}FN_5O_5S$ 501.15, found 502.4 (MH^+).

35 **Example 11.3: Preparation of 4-[5-Ethyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C6).**

Step1: Preparation of 4-(6-chloro-5-ethyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester.

To a solution of 4,6-dichloro-5-ethyl-pyrimidine (1 g, 5.65 mmol) and 4-hydroxy-piperidine-1-carboxylic acid isopropyl ester (1.05 g, 5.65 mmol) in dry THF under nitrogen at 0°C, potassium *tert*-butoxide (1M solution in THF, 6.78 mL) was added dropwise. The reaction was stirred at rt for 30 min. The mixture was quenched with water and extracted with EtOAc (3x). The organic layer was washed with water, sat. NH₄Cl and brine, followed by drying over sodium sulfate and concentration under *vacuo*. The resulting oil was purified by HPLC to afford 4-(6-chloro-5-ethyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (0.74 g, 39.8%) as a colorless oil. Exact mass calculated for C₁₅H₂₂ClN₃O₃ 327.13, found 328.2 (MH⁺).

Step 2: Preparation of 4-[5-ethyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C6).

4-(6-Chloro-5-ethyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (50 mg, 0.152 mmol), 2-fluoro-4-methanesulfonyl-phenol (43.5 mg, 0.228 mmol) and sodium hydride (60% dispersion in mineral oil, 7.28 mg, 0.182 mmol) were dissolved in DMSO (2 mL) and the mixture was heated under microwave irradiation for 1 h at 150°C. The crude was quenched with water and extracted with ethyl acetate. The organic layer was concentrated and the residue was purified by HPLC to afford Compound C6 (20.3 mg, 27.6%) as a white powder. ¹H NMR (CDCl₃, 400 MHz) δ 1.13 (t, J=7.33 Hz, 3H), 1.18 (d, J=6.32 Hz, 6H), 1.72-1.76 (m, 2H), 1.89-1.94 (d, 2H), 2.64 (q, J=7.33 Hz, 2H), 3.02 (s, 3H), 3.35-3.41 (m, 2H), 3.63-3.66 (m, 2H), 4.85-4.88 (m, 1H), 5.25-5.32 (m, 1H), 7.35-7.37 (m, 1H), 7.69-7.74 (m, 2H), 8.13 (s, 1H). Exact mass calculated for C₂₂H₂₈FN₃O₆S 481.17, found 482.4 (MH⁺).

Example 11.4: Preparation of 4-{6-[6-(2-Isopropoxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C10).

A mixture of 4-[6-(6-chloro-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (60 mg, 0.143 mmol), palladium acetate (12 mg, 0.05 mmol), 2,8,9-triisobutyl-2,5,8,9-tetraaza-1-phospha-bicyclo [3,3,3] undecane (5 mL, 0.015 mmol), 2-isopropoxy-ethylamine (35 µL, 0.28 mmol), and sodium *tert*-butoxide in 1.5 mL dioxane were heated under microwave irradiation for 1 hour at 120°C. Mixture was purified by HPLC to give Compound C10 as a tanned solid (TFA salt, 44.1 mg, 51%). ¹H NMR (MeOH-d₄, 400 MHz) δ 1.01-1.03 (d, 6H), 1.09-1.11 (d, 6H), 1.58-1.63 (m, 2H), 1.82-1.90 (m, 2H), 2.05 (s, 3H), 2.20 (s, 3H), 3.22-3.30 (m, 2H), 3.46-3.63 (m, 7H), 4.70-4.75 (m, 1H), 5.21-5.27 (m, 1H), 6.82-6.85 (d, 1H), 7.60-7.62 (d, 1H), 8.01 (s, 1H). Exact mass calculated for C₂₅H₃₇N₅O₅ 487.28, found 488.6 (MH⁺).

Example 11.5: Preparation of 4-{6-[6-(2-Hydroxy-ethylsulfanyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C12).

A mixture of 4-[6-(6-chloro-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (835 mg, 1.98 mmol) and potassium carbonate (305 mg, 2.2 mmol) in 3 mL of 2-mercapto-ethanol was stirred in microwave at 80°C. After 17 hours, mixture was continued to be stirred at 100°C for 30 minutes and then 120°C for 30 minutes. Mixture was purified by HPLC and column chromatography (hexane/AcOEt) to give Compound C12 as a white solid (16.4 mg, 2%). ¹H NMR (CDCl₃, 400 MHz) δ 1.07-1.09 (d, J = 6.3 Hz, 6H), 1.68-1.78 (m, 2H), 1.90-1.99 (m, 2H), 2.12 (s, 3H), 2.29 (s, 3H), 3.23-3.25 (t, J = 5.1 Hz, 2H), 3.32-3.38 (m, 2H), 3.68-3.71 (m, 2H), 3.91-3.94 (t, J = 4.9 Hz, 2H), 4.84-4.90 (m, 1H), 5.24-5.30 (m, 1H), 7.12-7.21 (m, 2H), 8.12 (s, 1H). Exact mass calculated for C₂₂H₃₀N₄O₅S 462.19, found 463.3 (MH⁺).

Example 11.6: Preparation of 4-[5-Methyl-6-(2-methyl-6-pentyl-pyridin-3-yloxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C15).

To a solution of 4-[6-(6-chloro-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (75.1 mg, 0.140 mmol) and iron (III)acetylacetonate (3.1 mg, 0.0088 mmol) in 1 mL THF and 0.1 mL NMP, 2M pentylmagnesium bromide solution in diethylether (135 µl, 0.275 mol) were added. After stirring at room temperature for several hours, mixture was purified by HPLC to give Compound C15 as an oil (TFA salt, 1.8 mg, 2%). Exact mass calculated for C₂₅H₃₆N₄O₄ 456.27, found 457.4 (MH⁺).

Example 11.7: Preparation of 1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butan-2-one (Compound C93).

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-(piperidin-4-yloxy)-pyrimidine hydrochloride salt (42 mg, 0.1 mmol), 1-bromo-butan-2-one (0.1 mmol, 1eq), and triethylamine (0.2 mmol, 2eq) were dissolved in DMF (1 mL) and then stirred at room temperature overnight. The crude was filtered and then purified via prep-LCMS 5-95% to provide Compound C93 as an oil (39.6 mg, 88 %). Exact mass calculated for C₂₁H₂₆FN₃O₅S 451.2, found 452.3 (MH⁺).

Example 11.8: Preparation of 4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(2-pyridin-3-yl-ethyl)-piperidin-4-yloxy]-pyrimidine (Compound C18).

A mixture of 4-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-(piperidin-4-yloxy)-pyrimidine (100 mg, 0.24 mmol), toluene-4-sulfonic acid 2-pyridin-3-yl-ethyl ester (133 mg, 0.48 mmol), and triethylamine (167 µL, 1.2 mmol) in DMF (2 mL) was heated under microwave irradiation for 60 min at 150°C. The crude mixture was concentrated in vacuo and purified by HPLC to provide Compound C18 as an oil (15 mg, 13%). ¹H NMR (CDCl₃, 400 MHz) δ 2.15 (s, 3H), 2.18-2.38 (m, 4H), 3.04 (s, 3H), 3.07-3.22 (m, 2H), 3.29-3.43 (m, 4H), 3.44-3.65 (m, 2H), 5.43-5.51 (m, 1H), 7.36 (t, 1H), 7.67-7.79 (m, 3H), 8.11 (s, 1H), 8.35 (d, 1H), 8.58 (d, 1H), 8.91 (s, 1H). Exact mass calculated for C₂₄H₂₇FN₄O₄S 486.56, found 487.4 (MH⁺).

Example 11.9: Preparation of 2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-trifluoromethoxy-phenyl)-ethanone (Compound C21).

Compound C21 was prepared using a similar procedure as described in Example 11.7 and
5 purified by preparative HPLC. ¹H NMR (CDCl₃, 400MHz) δ 2.22 (s, 3H), 2.33 (m, 2H), 2.48 (m, 2H), 3.14 (s, 3H), 3.69 (m, 4H), 4.78 (s, 2H), 5.58 (m, 1H), 7.34 (d, 2H), 7.44 (t, 1H), 7.79 (m, 2H), 7.99 (d, 2H), 8.22 (s, 1H). Exact mass calculated for C₂₆H₂₅F₄N₃O₆S 583.14, found 584.3 (MH⁺).

Example 11.10: Preparation of 4-{6-[6-(2-Methoxy-ethanesulfonyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C24).

A solution of 4-{6-[6-(2-methoxy-ethylsulfanyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (8.8 mg, 0.0185 mmol) in 1mL
methylene chloride was cooled in an ice bath and 3-chloroperoxybenzoic acid (9.4 mg, 0.038 mmol)
was added. After stirring for one hour in an ice-bath, mixture was quenched with a aqueous
15 bicarbonate solution and purified by HPLC to give Compound C24 as a white solid (TFA salt, 7.6 mg, 66%). ¹H NMR (MeOH-d₄, 400 MHz) δ 1.22-1.23 (d, 6H), 1.70-1.80 (m, 2H), 1.95-2.02 (m, 2H), 2.20 (s, 3H), 2.43 (s, 3H), 3.14 (s, 3H), 3.35-4.45 (m, 2H), 3.64-3.66 (t, J = 5.9 Hz, 3H), 3.70-3.76 (m, 4H), 4.82-4.88 (m, 1H), 5.35-5.39 (m, 1H), 7.72-7.75 (d, J = 8.36 Hz, 1H), 8.13 (s, 1H).
Exact mass calculated for C₂₃H₃₂N₄O₆S 508.20, found 509.4 (MH⁺).

20

Example 11.11: Preparation of 4-(2-Fluoro-4-methanesulfonyl-phenoxy)-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine (Compound C27).

Step 1: Preparation of N-hydroxy-isobutyramidine.

A solution of isobutyronitrile (276 g, 4.0 mol) in EtOH (2.0 L) was combined with
25 hydroxylamine (50% aqueous solution, 1.1 L, 16 mol), and refluxed for 5 h. The solvent was then removed in vacuo, and the residual water was azeotropically removed with toluene. The residue was then taken up in CH₂Cl₂, dried over MgSO₄, and the solvent was removed to afford a white solid (402 g, 98% yield). ¹H NMR (CDCl₃) δ 7.94 (br s, 1 H), 4.55 (br s, 2 H), 2.47 (m, 1 H), 1.20 (d, 6 H, J = 7.1 Hz).

30

Step 2: Preparation of 1-cyano-4-hydroxypiperidine.

A 5-liter, 3-neck flask was equipped with mechanical stirring, a reflux condenser, and a powder addition funnel. Sodium bicarbonate (840 g, 10 mmol) was added via the powder funnel while stirring, then water (ca. 300-400 mL) was gradually added while vigorously stirring to form a thick, uniform slurry. The flask was then placed in an ice bath, and a solution of 4-hydroxypiperidine
35 (506 g, 5.00 mol) in CH₂Cl₂ (1.0 L) was added, and the contents were vigorously mixed while cooling. A solution of cyanogen bromide (640 g, 6.0 mol) in CH₂Cl₂ (600 mL) was added in a dropwise fashion over 2 h, and stirring was continued for an additional 30 min. The ice bath was

removed, and the mechanical stirrer was replaced by a magnetic stirrer, and the reaction mixture was stirred for 16 h. The flask was once again placed under mechanical stirring, and sodium carbonate (100 g) was added in order to ensure complete neutralization. MgSO₄ (500 g) was added, and vigorous stirring was continued for 15 min. The resulting suspension was filtered, rinsing with CH₂Cl₂ (2.0 L). A light amber, viscous oil was obtained upon solvent removal to give 1-cyano-4-hydroxypiperidine (574 g, 91% yield). ¹H NMR (CDCl₃) δ 3.80 (m, 1 H), 3.39 (m, 2 H), 3.05 (m, 2 H), 1.87 (m, 2 H), 1.70 (br s, 1 H), 1.62 (m, 2 H), MS *m/z* 212.1 (M⁺).

Step 3: Preparation of 1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-ol.

In a variation of the method described by Yarovenko *et al.*, (Bull. Acad. Sci. USSR, Div. Chem. Sci. 1991, 40, 1924) ZnCl₂ (1 N in ether, 120 mL, 120 mmol) was added in a dropwise fashion over 15 min to a magnetically stirred solution of *N*-hydroxy-isobutyramidine (12.2 g, 120 mmol) and 4-hydroxy-piperidine-1-carbonitrile (12.6 g, 100 mmol) in ethyl acetate (500 mL). Precipitate formed immediately upon addition, and at a point the stirring bar became immobilized in the matrix, requiring the reaction to be manually shaken for the remainder of addition. After standing for 15 min, the supernatant was decanted and filtered, and the residue was rinsed twice with ether, furnishing a hard white precipitate which was collected by filtration. This material was taken up in conc. HCl (50 mL), diluted to 4 N with EtOH (100 mL), and refluxed for 1 h. Upon cooling, a white precipitate was removed by filtration, then the filtrate was reduced to 50 mL and diluted with 100 mL water. Solid Na₂CO₃ was added until the mixture was basic, CH₂Cl₂ was added, and the resulting mixture was filtered, rinsing with CH₂Cl₂. The organic extract was separated, dried over MgSO₄, and the solvent was removed to afford a viscous, amber oil as 1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-ol (15.0 g, 71% yield): ¹H NMR (CDCl₃) δ 3.95 (m, 3 H), 3.37 (m, 2 H), 2.88 (m, 1 H), 2.34 (br s, 1 H), 1.93 (m, 2 H), 1.63 (m, 2 H), 1.28 (d, 6 H, *J* = 7.1 Hz), MS *m/z* 212.3 (M⁺).

Step 4: Preparation of 4-chloro-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine.

To a solution of 1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-ol (3.65 g, 17 mmol) and 4,6-dichloro-5-methyl pyrimidine (2.83 g, 17 mmol) in THF (70 mL), 1M potassium-*t*-butoxide in THF (16 mL, 16 mmol) was added dropwise. The mixture was stirred at room temperature for 10 min. The crude mixture was purified by column chromatography on silica gel with hexane/ethyl acetate (3:1 v/v) to provide 4-chloro-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine as a solid (4.15 g, 71%). Exact mass calculated for C₁₅H₂₀ClN₅O₂ 337.13, found 338.2 (MH⁺).

Step 5: Preparation of 4-(2-fluoro-4-methanesulfonyl-phenoxy)-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine (Compound C27).

A mixture of 4-chloro-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine (756 mg, 2.24 mmol), 2-fluoro-4-methanesulfonyl-phenol (635 mg, 3.33 mmol), and sodium hydride, 60% dispersion in mineral oil (232 mg, 5.81 mmol) in DMAA (30 mL) was divided

into two 20 mL microwave vials and heated under microwave irradiation at 150°C for 1 hour. The crude mixture was purified by HPLC to provide Compound C27 as a solid (TFA salt, 75.5 mg, 5.6%). ¹H NMR (CDCl₃, 400 MHz) δ 1.28-1.31 (d, J = 6.32 Hz, 6H), 1.96-2.02 (m, 2H), 2.08-2.15 (m, 2H), 2.20 (s, 3H), 2.91-2.98 (m, 1H), 3.09 (s, 3H), 3.65-3.71 (m, 2H), 3.82-3.89 (m, 2H), 5.42-5.46 (m, 1H), 7.41-7.44 (m, 1H), 7.77-7.81 (m, 2H) 8.21 (s, 1H). Exact mass calculated for C₂₂H₂₆FN₅O₅S 491.16, found 492.3 (MH⁺).

Example 11.12: Preparation of 4-(6-{2-Fluoro-4-[(2-hydroxy-ethylcarbamoyl)-methyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (Compound C31).

Compound C31 was obtained in a similar manner as described in Example 11.36 as a solid (39 mg, 24%). ¹H NMR (MeOH-d₄, 400 MHz) δ 1.20 (d, 6H), 1.63-1.72 (m, 2H), 1.89-1.99 (m, 2H), 2.11 (s, 3H), 3.19-3.26 (m, 3H), 3.33-3.43 (m, 2H), 3.44-3.56 (m, 4H), 3.64-3.73 (m, 2H), 4.80 (s, 1H), 5.30 (h, 1H), 7.02-7.16 (m, 3H), 8.03 (s, 1H). Exact mass calculated for C₂₄H₃₁FN₄O₆ 490.52, found 491.4 (MH⁺).

Example 11.13: Preparation of 4-[6-(5-Iodo-pyridin-2-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C34).

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (1.02 g, 3.25 mmol), 2-hydroxy-5-iodopyridine, and potassium carbonate (903 mg, 6.53 mmol) in 15 mL DMF were heated under microwave irradiation for 1 hour at 150°C. Mixture was purified by HPLC to give Compound C34 (TFA salt, 177 mg, 9%) as a white solid. ¹H NMR (CDCl₃, 400 MHz) δ 1.24-1.25 (d, 6H), 1.75-1.83 (m, 2H), 1.97-2.03 (m, 2H), 2.20 (s, 3H), 3.40-3.45 (m, 2H), 3.71-3.79 (m, 2H), 4.91-4.97 (m, 1H), 5.33-5.39 (m, 1H), 6.93-6.95 (d, 1H), 8.04-8.07 (dd, 1H), 8.31 (s, 1H), 8.51-8.52 (d, 1H). Exact mass calculated for C₁₉H₂₃IN₄O₄ 498.08, found 499.2 (MH⁺).

Example 11.14: Preparation of 4-(6-{2-Fluoro-4-[N-(2-isopropoxy-ethyl)-carbamimidoyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (Compound C36) and 4-[6-(4-Carbamoyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C139).

Step 1: Preparation of 4-[6-(2-fluoro-4-phenylsulfanylcarbonimidoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester.

A mixture of 4-[6-(4-cyano-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (TFA salt, 109 mg, 0.21 mmol) and thiophenol (27 μl, 0.21 mmol) in Et₂O (1 mL) was stirred in an ice-bath under HBr atmosphere for 30 min. The crude compound was

used for the next step without further purification. Exact mass calculated for $C_{27}H_{29}FN_4O_4S$ 524.19, found 525.3 (MH^+).

Step 2: Preparation of 4-(6-{2-Fluoro-4-[N-(2-isopropoxy-ethyl)-carbamimidoyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (Compound C36) and 4-[6-(4-Carbamoyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C139).

A mixture of 4-[6-(2-fluoro-4-phenylsulfanylcabonimidoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (108.4 mg, 0.21 mmol) and 2-aminoethyl isopropyl ether (101 μ l, 0.83 mmol) in MeOH (2 mL) was stirred at room temperature for 30 min. Additional 2-aminoethyl isopropyl ether (2 mL, 16.3 mmol) was added and the mixture was stirred under 70°C for 10 min. The crude mixture was purified by HPLC to provide Compound C36 as a solid (TFA salt, 19.6 mg, 15%). and Compound C139 in solid as a by-product (TFA salt, 23.7 mg, 21%). Compound C36: 1H NMR ($CDCl_3$, 400 MHz) δ 1.13-1.20 (m, 6H), 1.26-1.27 (d, 6H), 1.78-1.80 (m, 2H), 1.98-1.99 (m, 2H), 2.19-2.20 (d, 3H), 2.82 (s, 5H), 3.39-3.46 (m, 3H), 3.68-3.79 (m, 4H), 4.92-4.95 (m, 1H), 5.34-5.35 (m, 1H), 7.38-7.44 (m, 1H), 7.68-7.71 (t, 1H), 8.13-8.19 (d, 1H). Exact mass calculated for Compound C36, $C_{26}H_{36}FN_5O_5$ 517.27, found 518.5 (MH^+) and for Compound C139, exact mass calculated for $C_{21}H_{25}FN_4O_5$ 432.18, found 433.1 (MH^+).

Example 11.15: Preparation of 4-[6-(4-Carboxy-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C38).

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (300 mg, 0.96 mmol), 2 (150 mg, 0.96 mmol) and potassium carbonate (160 mg, 1.15 mmol) in DMSO was heated under microwave for 4 hrs at 160°C. The mixture was purified through HPLC to afford Compound C38 (200 mg, 48%) as a solid and Compound C9 as a by product. Compound C38: 1H NMR ($MeOH-d_4$, 400 MHz) δ 1.15 (d, 6H), 1.64-1.67 (m, 2H), 1.88-1.92 (m, 2H), 2.09 (s, 3H), 3.29-3.31 (m, 2H), 3.62-3.66 (m, 2H), 4.72-4.78 (m, 1H), 5.25-5.28 (m, 1H), 7.23 (t, 1H), 7.70 (d, 1H), 7.77 (d, 1H), 8.02 (s, 1H). Exact mass calculated for $C_{21}H_{24}FN_3O_6$ 433.2, found 434.3 (MH^+). Compound C9: Exact mass calculated for $C_{20}H_{24}FN_3O_4$ 389.2, found 390.3 (MH^+).

Example 11.16: Preparation of 4-(4-Bromo-2-fluoro-phenoxy)-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine (Compound C40).

A mixture of 4-chloro-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine (1.51 g, 4.46 mmol), potassium carbonate (1.25 g, 9.03 mmol), and 4-bromo-2-fluorophenol (1.11 g, 5.82 mmol) in 15 mL DMF was heated in microwave for 1 hour at 150°C. The mixture was purified by column chromatography on silica gel with hexane/ethyl acetate (3:1 v/v) to give Compound C40 as an oil (1.05 g, 48%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.20-1.22 (d, 6H), 1.83-1.91 (m, 2H), 1.98-2.05 (m, 2H), 2.11 (s, 3H), 2.77-2.87 (m, 1H), 3.53-3.59 (m, 2H), 3.74-3.80 (m,

2H), 5.31-5.36 (m, 1H), 6.99-7.03 (m, 1H), 7.22-7.29 (m, 2H), 8.13 (s, 1H). Exact mass calculated for $C_{21}H_{23}BrFN_5O_3$ 491.1, found 492.4 (MH^+).

Example 11.17: Preparation of 4-[6-(5-Methanesulfonyl-pyridin-2-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C42).

A mixture of 4-[6-(5-iodo-pyridin-2-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (TFA salt, 35.1 mg, 0.07 mmol), sodium methanesulfinate (21.4 mg, 0.21 mmol), copper (I) trifluoromethanesulfonate benzene complex (3.5 mg, 0.007 mmol), and *N,N*-dimethyl-ethane-1,2-diamine in 1.5 mL DMSO were heated under microwave irradiation for 30 minutes at 160°C. Mixture was purified by HPLC to give Compound C42 as a white solid (TFA salt, 11.5 mg, 30%). 1H NMR ($MeOH-d_4$, 400MHz) δ 1.21-1.22 (d, 6H), 1.71-1.79 (m, 2H), 1.96-2.07 (m, 5H), 3.16 (s, 3H), 3.35-4.42 (m, 2H), 3.80-3.87 (m, 2H), 4.80-4.86 (m, 1H), 5.38-5.42 (m, 1H), 7.32-7.34 (d, 1H), 8.29 (s, 1H), 8.35-8.37 (dd, 1H), 8.69 (s, 1H). Exact mass calculated for $C_{20}H_{26}N_4O_6S$ 450.16, found 451.4 (MH^+).

Example 11.18: Preparation of 4-{6-[6-(2-Hydroxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C43).

To a solution of 4-{6-[6-(2-methoxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (TFA salt, 87 mg, 0.152 mmol) in methylene chloride, trimethylsilyl iodide (300 μ l, 1.5 mmol) was added. After stirring for 3 hours at room temperature, mixture was quenched with methanol and purified by HPLC to give Compound C43 as a white solid (TFA salt, 40.6 mg, 48%). 1H NMR ($MeOH-d_4$, 400 MHz) δ 1.22-1.23 (d, J = 6.2 Hz, 6H), 1.69-1.77 (m, 2H), 1.94-2.02 (m, 2H), 2.17 (s, 3H), 2.31 (s, 3H), 3.32-3.40 (m, 2H), 3.52-3.55 (t, J = 5.1 Hz, 2H), 3.70-3.80 (m, 4H), 4.84-4.88 (m, 1H), 5.34-5.38 (m, 1H), 6.93-6.95 (d, J = 9.6 Hz, 1H), 7.70-7.73 (d, J = 9.6 Hz, 1H), 8.13 (s, 1H). Exact mass calculated for $C_{22}H_{31}N_5O_5$ 445.23, found 446.3 (MH^+).

Example 11.19: Preparation of 4-[5-Cyclopropyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C44).

Step 1: Preparation of 4-(6-chloro-5-cyclopropyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester.

To a solution of 4,6-dichloro-5-cyclopropyl-pyrimidine (700 mg, 3.70 mmol) and 4-hydroxy-piperidine-1-carboxylic acid isopropyl ester (636.6 mg, 3.70 mmol) in dry THF under nitrogen at 0°C, potassium tert-butoxide (1M solution in THF, 4.45 mL) was added dropwise. The reaction was stirred at rt for 30 min. The mixture was quenched with water and extracted with EtOAc (3x). The organic layer was washed with water, sat. NH_4Cl and brine, followed by drying over sodium sulfate and concentration under *vacuo*. The resulting oil was purified by flash chromatography (0-20%

EtOAc/ Hexanes) to afford 4-(6-chloro-5-cyclopropyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (0.927 g, 73.7%) as colorless oil. Exact mass calculated for $C_{16}H_{22}ClN_3O_3$ 339.13, found 340.3 (MH^+).

Step 2: Preparation of 4-[5-Cyclopropyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C44).

4-(6-Chloro-5-cyclopropyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (200 mg, 0.588 mmol), 2-fluoro-4-methanesulfonyl-phenol (168 mg, 0.883 mmol) and sodium hydride (60% dispersion in mineral oil, 53 mg, 1.325 mmol) were dissolved in DMSO (2 mL) and the mixture stirred under Nitrogen for 10 min at rt. The mixture was then heated under microwave irradiation for 1h at 150°C. The crude was quenched with water and extracted with EtOAc (3x). The organic layer was concentrated and the residue was purified by HPLC to afford Compound C44 (51 mg, 14.3%) as oil. 1H NMR ($CDCl_3$, 400 MHz) δ 0.90-0.97 (m, 2H), 1.01-1.06 (m, 2H), 1.23-1.27 (d, $J=6.06$ Hz, 6H), 1.76-1.91 (m, 2H), 1.92-2.02 (m, 2H), 3.08 (s, 3H), 3.46-3.55 (m, 2H), 3.63-3.72 (m, 2H), 4.87-4.98 (m, 1H), 5.32-5.39 (m, 1H), 7.36-7.42 (m, 1H), 7.73-7.80 (m, 2H), 8.17 (s, 1H). Exact mass calculated for $C_{23}H_{28}FN_3O_6S$ 493.17, found 494.5 (MH^+).

Example 11.20: Preparation of 4-{6-[6-(2-Methanesulfonyl-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C46).

Using a similar procedure as described in Example 11.4 for the preparation of Compound C10, Compound C46 was obtained as a tanned solid (TFA salt, 27.0 mg, 30%). 1H NMR ($MeOH-d_4$, 400 MHz) δ 1.05-1.06 (d, 6H), 1.64-1.72 (m, 2H), 1.88-1.97 (m, 2H), 2.12 (s, 3H), 2.27 (s, 3H), 2.98 (s, 3H), 3.29-3.37 (m, 2H), 3.43-3.46 (t, 2H), 3.65-3.71 (m, 2H), 3.86-3.89 (t, 2H), 4.78-4.82 (m, 1H), 5.29-5.33 (m, 1H), 6.90-6.92 (d, 1H), 7.72-7.74 (d, 1H), 8.07 (s, 1H). Exact mass calculated for $C_{23}H_{33}N_5O_6S$ 507.22, found 508.6 (MH^+).

Example 11.21: Preparation of 1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-5-methyl-hexan-1-one (Compound C121).

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-(piperidin-4-yloxy)-pyrimidine (42mg, 0.1mmol), 5-methyl-hexanoic acid (0.12mmol, 1.2eq), HATU (0.12mmol, 1.2eq), and triethylamine (0.2mmol, 2eq) were dissolved in DMF (1mL), and then stirred at room temperature for 1 hour. The crude was filtered and then purified via prep-LCMS 5-95% to provide Compound C121 as a white powder (28.2 mg, 57 %). Exact mass calculated for $C_{24}H_{32}FN_3O_5S$ 493.2, found LCMS (ESI) m/z 494.5 (MH^+).

Example 11.22: Preparation of 4-{6-[6-(2-Methoxy-ethylsulfanyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C56).

To a solution of compound **C12** (15mg, 0.0324 mmol) in 1 mL THF, sodium hydride dispersion (8 mg, 0.2 mmol) was added. After 10 minutes, methyl iodide (20 μ l, 0.32 mmol) was added and mixture was stirred at room temperature for 17 hours. Mixture was purified by column chromatography (AcOEt/hexane) to give Compound **C56** as white solid (10.1 mg, 65%). ^1H NMR (CDCl₃, 400 MHz) δ 1.25-1.27 (d, J = 6.3 Hz, 6H), 1.75-1.85 (m, 2H), 1.95-2.05 (m, 2H), 2.19 (s, 3H), 2.34 (s, 3H), 3.37-3.45 (m, 7H), 3.65-3.68 (t, J = 6.7 Hz, 2H), 3.75-3.81 (m, 2H), 4.91-4.97 (m, 1H), 5.32-5.36 (m, 1H), 7.07-7.09 (d, J = 8.4 Hz, 1H), 7.22-7.20 (d, J = 8.4 Hz, 1H), 8.19 (s, 1H). Exact mass calculated for C₂₃H₃₂N₄O₅S 476.21, found 477.4 (MH⁺).

Example 11.23: Preparation of 1-(2,5-Dimethoxy-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone (Compound C60).

Compound **C60** was prepared using a similar procedure as described in Example 11.7 and was obtained as an oil (35.9 mg, 64%). Exact mass calculated for C₂₇H₃₀FN₃O₇S 559.2, found LCMS (ESI) m/z 560.4 (MH⁺).

Example 11.24: Preparation of 4-[6-(6-Chloro-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C65).

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (1.03 g, 3.27 mmol), 6-chloro-2-methyl-pyridin-3-ol (470 mg, 3.27 mmol), and potassium carbonate (903 mg, 6.53 mmol) in 15 mL DMF were heated in microwave for 1 hour at 150°C. The mixture was purified to give Compound **C65** as a white solid (0.975 g, 71%). ^1H NMR (CDCl₃, 400 MHz) δ 0.92-.94 (d, 6H), 1.74-1.82 (m, 2H), 1.95-2.02 (m, 2H), 2.19 (s, 3H), 2.47 (s, 3H), 3.39-3.45 (m, 2H), 3.74-3.79 (m, 2H), 4.91-4.97 (m, 1H), 5.33-5.36 (m, 1H), 7.21-7.23 (d, 1H), 7.36-7.38 (d, 1H), 8.19 (s, 1H). Exact mass calculated for C₂₀H₂₅ClN₄O₄ 420.16, found 421.3 (MH⁺).

Example 11.25: Preparation of 1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3,3-dimethyl-butan-2-one (Compound C78).

Compound **C78** was prepared using a similar procedure as described in Example 11.7 and was obtained as an oil (26 mg, 54%). Exact mass calculated for C₂₃H₃₀FN₃O₅S 479.2, found LCMS (ESI) m/z 480.4 (MH⁺).

Example 11.26: Preparation of 2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-pyridin-2-yl-ethanone (Compound C22).

Compound **C22** was prepared using a similar procedure as described in Example 11.7 and was purified by preparative HPLC. ^1H NMR (CDCl₃, 400MHz) δ 2.24 (s, 3H), 2.29 (m, 1H), 2.42 (m, 1H), 2.57 (m, 2H), 3.11 (s, 3H), 3.51 (m, 2H), 3.77 (m, 2H), 4.98 (s, 2H), 5.60 (m, 1H), 7.43 (t,

1H), 7.61 (m, 1H), 7.82 (m, 2H), 7.91 (m, 1H), 8.23 (m, 1H), 8.67 (m, 1H). Exact mass calculated for $C_{24}H_{25}FN_4O_5S$ 500.15, found 501.3 (MH^+).

Example 11.27: Preparation of 2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(3-fluoro-phenyl)-ethanone (Compound C16).

Compound C16 was prepared using a similar procedure as described in Example 11.7 and purified by preparative HPLC. 1H NMR ($CDCl_3$, 400MHz) δ 2.31 (s, 3H), 2.34 (m, 2H), 2.39 (m, 2H), 3.11 (s, 3H), 3.68 (m, 4H), 4.78 (s, 2H), 5.59 (m, 1H), 7.36 (m, 1H), 7.46 (m, 1H), 7.52 (m, 1H), 7.63 (m, 1H), 7.71 (m, 1H), 7.82 (m, 2H), 8.22 (s, 1H). Exact mass calculated for $C_{25}H_{25}F_2N_3O_5S$ 517.15, found 518.3 (MH^+).

Example 11.28: Preparation of 4-(6-{2-Fluoro-4-[(2-isopropoxy-ethylcarbamoyl)-methyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (Compound C101).

Compound C101 was obtained in a similar manner as described in Example 11.36 as a solid (30 mg, 20%). 1H NMR ($DMSO-d_6$, 400 MHz) δ 1.08 (d, 6H), 1.21 (d, 6H), 1.62-1.73 (m, 2H), 1.92-2.01 (m, 2H), 2.16 (s, 3H), 3.21 (q, 2H), 3.31-3.41 (m, 4H), 3.48 (s, 2H), 3.50-3.60 (m, 1H), 3.61-3.71 (m, 2H), 4.80 (h, 1H), 5.33 (h, 1H), 7.13 (d, 1H), 7.22-7.30 (m, 2H), 8.17 (t, 1H), 8.26 (s, 1H). Exact mass calculated for $C_{27}H_{37}FN_4O_6$ 532.60, found 533.4 (MH^+).

Example 11.29: Preparation of 4-{6-[2-Fluoro-4-(2-isopropoxy-ethylcarbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C113).

Compound C113 was obtained in a similar manner as described in Example 11.36 as a solid (25 mg, 83%). 1H NMR ($MeOH-d_4$, 400 MHz) δ 1.03 (d, 6H), 1.12 (d, 6H), 1.62-1.65 (m, 2H), 1.86-1.91 (m, 2H), 2.07 (s, 3H), 3.25-3.31 (m, 2H), 3.38-3.44 (m, 2H), 3.48-3.53 (m, 3H), 3.60-3.68 (m, 2H), 4.74-4.76 (m, 1H), 5.24-5.27 (m, 1H), 7.21 (t, 1H), 4.56-7.59 (m, 2H), 8.00 (s, 1H), 8.43 (t, 1H). Exact mass calculated for $C_{26}H_{35}FN_4O_6$ 518.2, found 519.5 (MH^+).

Example 11.30: Preparation of 1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butan-1-one (Compound C115).

Compound C115 was prepared in a similar manner as described in Example 11.21 and was obtained as a white powder (27.1 mg, 60%). Exact mass calculated for $C_{21}H_{26}FN_3O_5S$ 451.2, found 452.2 (MH^+).

Example 11.31: Preparation of 1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-pentan-1-one (Compound C116).

Compound **C116** was prepared in a similar manner as described in Example 11.21 and was obtained as a white powder (29.9 mg, 64%). Exact mass calculated for $C_{22}H_{28}FN_3O_5S$ 465.2, found 466.4 (MH^+).

5 Example 11.32: Preparation of 4-[6-(2,4-Difluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C117).

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (1.0 g, 3.19 mmol), 2,4-difluoro-phenol (585 mg, 4.5 mmol), and potassium carbonate (882 mg, 6.38 mmol) in DMF (11 mL) was heated under microwave irradiation for 80 min at 150°C. The crude
10 mixture was concentrated in vacuo and purified by HPLC to provide Compound **C117** as a beige solid (890 mg, 69%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.19 (d, 6H), 1.66-1.76 (m, 2H), 1.86-1.95 (m, 2H), 2.11 (s, 3H), 3.30-3.40 (m, 2H), 3.63-3.73 (m, 2H), 4.86 (m, 1H), 5.26 (m, 1H), 6.80-6.91 (m, 2H), 7.09 (q, 1H), 7.19 (s, 1H). Exact mass calculated for $C_{20}H_{23}F_2N_3O_4$ 407.41, found 408.3 (MH^+).

15 Example 11.33: Preparation of 1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-butan-1-one (Compound C119).

Compound **C119** was obtained in a similar manner as described in Example 11.21 as a white powder (28.5 mg, 61%). Exact mass calculated for $C_{22}H_{28}FN_3O_5S$ 465.2, found 466.4 (MH^+).

20 Example 11.34: Preparation of 1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-methyl-pentan-1-one (Compound C120).

Compound **C120** was obtained in a similar manner as described in Example 11.21 as an oil (33.3 mg, 69%). Exact mass calculated for $C_{23}H_{30}FN_3O_5S$ 479.2, found LCMS (ESI) m/z 480.4 (MH^+).

25 Example 11.35: Preparation of 4-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-oxo-butyric acid (Compound C51).

Compound **C51** was obtained in a similar manner as described in Example 11.21 as a white powder (11.9 mg, 25%). Exact mass calculated for $C_{21}H_{24}FN_3O_7S$ 481.1, found LCMS (ESI) m/z
30 482.2 (MH^+).

Example 11.36: Preparation of 4-{6-[2-Fluoro-4-(2-methoxy-ethylcarbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C122).

A mixture of 4-[6-(4-Carboxy-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (150 mg, 0.346 mmol), 2-Methoxy-ethylamine (31 mg, 0.41 mmol), HATU (157 mg, 0.42 mmol) and triethyl amine (70 mg, 0.7 mmol) in DMF (5 mL) was stirred at
35 room temperature for 2 h. The mixture was purified through HPLC to provide Compound **C122** (115

mg, 68%) as a solid. ^1H NMR (MeOH-d_4 , 400 MHz) δ 1.22 (d, $J = 6.82$ Hz, 6H), 1.71-1.74 (m, 2H), 1.97-2.00 (m, 2H), 2.16 (s, 3H), 3.34 (s, 3H), 3.35-3.39 (m, 2H), 3.53 (s, 4H), 3.71-3.74 (m, 2H), 4.81-4.84 (m, 1H), 5.33-5.36 (m, 1H), 7.30 (t, $J = 8.1$ Hz, 1H), 7.66-7.69 (m, 2H), 8.09 (s, 1H). Exact mass calculated for $\text{C}_{24}\text{H}_{31}\text{FN}_4\text{O}_6$ 490.2, found 491.4 (MH^+).

5

Example 11.37: Preparation of 4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C127).

Step 1: Preparation of 4-[6-(4-bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester.

10 A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (1.0098 g, 3.2 mmol), potassium carbonate (889.5 mg, 6.43 mmol), and 4-bromo-2-fluorophenol (458 μL , 4.18 mmol) in 15 mL DMF was heated in microwave for 1 hour at 150°C . The mixture was purified by HPLC to give 4-[6-(4-bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester as a tanned solid as the TFA salt (Compound C130, 741 mg, 39%).

15 Exact mass calculated for $\text{C}_{20}\text{H}_{23}\text{BrFN}_3\text{O}_4$ 467.09, found 468.3 (MH^+).

Step 2: Preparation of 4-[6-(4-Bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C127).

A mixture of 4-[6-(4-bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (741 mg, 1.27 mmol), sodium methane sulfinate (288.4 mg, 2.82 mmol), and N,N' -dimethyl-ethylene diamine (28 mg, 0.317 mmol) and copper (I) trifluoromethane sulfonate benzene complex (95.6 mg, 0.190 mmol) in 10 mL DMSO was heated in microwave for 30 min at 160°C . The mixture was purified by HPLC to give Compound C127 as a white solid (TFA salt, 327.1 mg, 44%). ^1H NMR ($\text{CD}_3\text{CN-d}_3$, 400 MHz) δ 1.21-1.23 (d, $J = 6.32$ Hz, 6H), 1.69-1.77 (m, 2H), 1.93-1.95 (m, 2H), 2.184 (s, 3H), 3.125 (s, 3H), 3.359-3.441 (m, 2H), 3.650-3.734 (m, 2H), 20 4.84 (hept, $J = 6.32$ Hz, 1H), 5.313-5.380 (m, 1H), 7.470-7.526 (m, 1H), 7.781-7.846 (m, 2H) 8.158 (s, 1H). Exact mass calculated for $\text{C}_{21}\text{H}_{26}\text{FN}_3\text{O}_6\text{S}$ 467.15, found 468.4 (MH^+).

25

Example 11.38: Preparation of 4-[6-[2-Fluoro-4-(methoxy-methyl-carbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C132).

30 Compound C132 was obtained in a similar manner as described in Example 11.36 as an oil (40 mg, 85%). ^1H NMR (CDCl_3 , 400 MHz) δ 1.26 (d, $J = 6.32$ Hz, 6H), 1.75-1.83 (m, 2H), 1.96-2.02 (m, 2H), 2.18 (s, 3H), 3.38 (s, 3H), 3.39-3.46 (m, 2H), 3.60 (s, 3H), 3.71-3.77 (m, 2H), 4.93 (hept, $J = 6.32$ Hz, 1H), 5.31-5.36 (m, 1H), 7.21-7.26 (m, 1H), 7.58-7.62 (m, 2H), 8.20 (s, 1H). Exact mass calculated for $\text{C}_{23}\text{H}_{29}\text{FN}_4\text{O}_6$ 476.2, found 477.3 (MH^+).

35

Example 11.39: Preparation of 1-[4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl]-3-methoxy-propan-1-one (Compound C133).

Compound C132 was prepared in a similar manner as described in Example 11.21 and was obtained as a white powder (30.5 mg, 65%). Exact mass calculated for $C_{21}H_{26}FN_3O_6S$ 467.1, found LCMS (ESI) m/z 468.2 (MH^+).

5 **Example 11.40: Preparation of 4-[6-(4-Cyano-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C134).**

A mixture of 4-[6-(4-bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C130, 1.14 g, 2.4 mmol), zinc cyanide (290 mg, 2.42 mmol), and tetrakis(triphenylphosphine)palladium (0) (281 mg, 0.24 mmol) in DMF (15 mL) was
10 purged with Argon and heated under microwave irradiation at 180°C for 8 min. The crude mixture was purified by HPLC to provide Compound C134 as a solid (TFA salt, 318 mg, 27%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.27-1.28 (d, J = 6.32 Hz, 6H), 1.81-1.84 (m, 2H), 1.99-2.02 (m, 2H), 2.20 (s, 3H), 3.43-3.49 (m, 2H), 3.75-3.77 (m, 2H), 4.94-4.97 (m, J = 6.32 Hz, 1H), 5.35-5.36 (m, J = 3.79 Hz, 1H), 7.33-7.37 (m, 1H), 7.49-7.54 (m, 2H), 8.21 (s, 1H). Exact mass calculated for $C_{21}H_{23}FN_4O_4$
15 414.17, found 415.4 (MH^+).

Example 11.41: Preparation of 4-[5-(5-Aminomethyl-4,5-dihydro-oxazol-2-yl)-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C135).

In a 100mL round-bottomed flask fitted with a condenser and N_2 inlet was placed a stir bar, 4-[5-cyano-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (1g, 2mmol), $ZnCl_2$ (30mg, 0.2mmol), 1,3-diamino-propan-2-ol (180mg, 2mmol), and chlorobenzene (20mL). The reaction mixture was heated under reflux overnight. After
20 it was cooled down to room temperature, the reaction was quenched with H_2O . The resulting suspension was extracted with EtOAc. The organic extracts was dried and concentrated under vacuum. The crude residue was purified by preparative HPLC to give Compound C135. 1H NMR ($CDCl_3$, 400MHz) δ 1.25 (d, 6H), 1.73 (m, 2H), 1.94 (m, 2H), 2.98 (s, 3H), 3.31 (m, 2H), 3.46 (m, 2H), 3.69 (m, 4H), 4.48 (m, 1H), 4.83 (m, 1H), 5.36 (m, 1H), 7.40 (d, 1H), 7.49 (d, 1H), 8.36 (t, 1H), 8.44 (s, 1H), 9.52 (m, 2H). Exact mass calculated for $C_{24}H_{31}FN_6O_6S$ 550.20, found 551.3 (MH^+).

30 **Example 11.42: Preparation of 4-{6-[6-(2-Methoxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C136).**

Using a similar procedure as described in Example 11.4 for the preparation of Compound C10, Compound C136 was obtained as a tanned solid (TFA salt, 167.8 mg, 65%). 1H NMR ($MeOH-d_4$, 400 MHz) δ 1.48-1.49 (d, 6H), 1.95-2.02 (m, 2H), 2.21-2.28 (m, 2H), 2.43 (s, 3H), 2.58 (s, 3H),
35 3.61-3.67 (m, 5H), 3.85-3.89 (m, 4H), 3.95-4.03 (m, 2H), 5.10-5.13 (m, 1H), 7.20-7.22 (d, 1H), 7.98-

8.00 (d, 1H), 7.98-8.00 (d, 1H), 8.40 (s, 1H). Exact mass calculated for $C_{23}H_{33}N_5O_5$ 459.25, found 460.3 (MH^+).

Example 11.43: Preparation of 4-{6-[6-(3-Methanesulfonyl-pyrrolidin-1-yl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C137).

Using a similar procedure as described in Example 11.4 for the preparation of Compound C10, Compound C137 was obtained as an oil (TFA salt, 54.4 mg, 58%). 1H NMR (MeOH- d_4 , 400 MHz) δ 1.22-1.23 (d, 6H), 1.69-1.77 (m, 2H), 1.95-2.02 (m, 2H), 2.18 (s, 3H), 2.38 (s, 3H), 2.52-2.70 (m, 2H), 3.07 (s, 3H), 3.32-3.42 (m, 2H), 3.71-4.19 (m, 8H), 5.35-5.38 (m, 1H), 6.97-6.99 (d, 1H), 7.80-7.83 (d, 1H), 8.12 (s, 1H). Exact mass calculated for $C_{25}H_{35}N_5O_6S$ 533.23, found 534.5 (MH^+).

Example 11.44: Preparation of 4-[6-(6-Benzylamino-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C138).

Using a similar procedure as described in Example 11.4 for the preparation of Compound C10, Compound C138 was obtained as an oil (TFA salt, 80.8 mg, 61%). 1H NMR (MeOH- d_4 , 400 MHz) δ 1.07-1.10 (d, 6H), 1.68-1.77 (m, 2H), 1.93-1.91 (m, 2H), 2.17 (s, 3H), 2.34 (s, 3H), 3.31-3.41 (m, 2H), 3.69-3.78 (m, 2H), 3.94 (s, 1H), 4.61 (s, 2H), 5.31-5.36 (m, 1H), 6.89-6.91 (d, 1H), 7.30-7.39 (m, 5H), 7.73-7.75 (d, 1H), 8.13 (s, 1H). Exact mass calculated for $C_{27}H_{33}N_5O_4$ 491.25, found 492.5 (MH^+).

Example 11.45: Preparation of 2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-pyridin-2-yl-ethanone (Compound C61).

Compound C61 was prepared in a similar manner as described in Example 11.7. 1H NMR ($CDCl_3$, 400MHz) δ 2.18 (s, 3H), 2.24 (m, 1H), 2.29 (m, 1H), 2.57 (m, 2H), 3.11 (s, 3H), 3.52 (m, 2H), 3.77 (m, 2H), 4.98 (s, 2H), 5.60 (m, 1H), 7.45 (t, 1H), 7.60 (m, 1H), 7.82 (m, 2H), 7.91 (m, 1H), 8.10 (m, 1H), 8.23 (m, 1H), 8.67 (m, 1H). Exact mass calculated for $C_{24}H_{25}FN_4O_5S$ 500.15, found 501.1 (MH^+).

Example 11.46: Preparation of 4-{6-[2-Fluoro-4-(2-isopropoxy-ethylamino)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C140).

Compound C140 was obtained in a similar manner as described in Example 9.71 (Compound A95) as a orange oil (54 mg, 55%). 1H NMR ($CDCl_3$, 400 MHz) δ 1.19 (d, $J = 6.1$ Hz, 6H), 1.27 (d, $J = 6.1$ Hz, 6H), 1.79-1.83 (m, 2H), 1.97-2.02 (m, 2H), 2.20 (s, 3H), 3.41-3.48 (m, 2H), 3.65-3.69 (m, 3H), 3.73-3.78 (m, 2H), 4.95 (sept, $J = 6.3$ Hz, 1H), 5.33-5.36 (m, 1H), 6.91-6.97 (m, 2H), 7.18 (t, $J = 8.3$ Hz, 1H), 8.23 (s, 1H). Exact mass calculated for $C_{25}H_{35}FN_4O_5$ 490.3, found 491.4 (MH^+).

Example 11.47: Preparation of 4-(6-{2-Fluoro-4-[(tetrahydro-furan-2-ylmethyl)-amino]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (Compound C141).

5 Compound **C141** was obtained in a similar manner as described in Example 9.71 as a tan solid (62 mg, 63%). ¹H NMR (CDCl₃, 400 MHz) δ 1.26 (d, J = 6.3 Hz, 6H), 1.61-1.70 (m, 1H), 1.76-1.84 (m, 2H), 1.94-1.99 (m, 4H), 2.04-2.11 (m, 1H), 2.18 (s, 3H), 3.11 (dd, J = 12.4 Hz, 8.3Hz 1H), 3.29 (dd, J = 12.4 Hz, 3.8Hz, 1H), 3.41-3.47 (m, 2H), 3.72-3.79 (m, 2H), 3.80-3.84 (m, 2H), 3.88-3.94 (m, 2H), 4.17 (qd, J = 8.5 Hz, 3.5Hz, 1H), 4.94 (sept, J = 6.3 Hz, 1H), 5.30-5.36 (m, 1H), 6.64 (dd, J = 12.1 Hz, 2.5Hz, 1H), 6.66 (dd, J = 14.9 Hz, 2.5Hz, 1H), 7.05 (t, J = 8.3 Hz, 1H), 8.23 (s, 1H). Exact mass calculated for C₂₅H₃₃FN₄O₅ 488.2, found 489.4 (MH⁺).

Example 11.48: Preparation of 4-(6-{[2-Methanesulfonyl-ethyl)-methyl-amino]-2-methyl-pyridin-3-yloxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (Compound C142).

15 Using a similar procedure as described in Example 11.4 for the preparation of Compound **C10**, Compound **C142** was obtained as an oil (TFA salt, 54.1 mg, 50%). ¹H NMR (MeOH-d₄, 400 MHz) δ 1.00-1.02 (d, 6H), 1.70-1.77 (m, 2H), 1.95-2.02 (m, 2H), 2.17 (s, 3H), 2.33 (s, 3H), 3.00 (s, 3H), 3.19 (s, 3H), 3.34-3.41 (m, 2H), 3.49-3.53 (t, 2H), 3.70-3.76 (m, 2H), 3.94 (s, 1H), 4.09-4.12 (t, 2H), 5.33-5.36 (m, 1H), 6.84-6.86 (d, 1H), 7.53-7.55 (d, 1H), 8.10 (s, 1H). Exact mass calculated for C₂₄H₃₅N₅O₆S 521.23, found 522.5 (MH⁺).

Example 11.49: Preparation of 4-[6-(2-Fluoro-4-hydroxycarbamoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C143).

25 Compound **C143** was obtained in a similar manner as described in Example 11.36 as an oil (66 mg, 80%). Exact mass calculated for C₂₁H₂₅FN₄O₆ 448.2, found 449.3 (MH⁺).

Example 11.50: Preparation of 4-{6-[2-Fluoro-4-(2-pyrrolidin-1-yl-ethylcarbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C144).

30 Compound **C144** was obtained in a similar manner as described in Example 11.36 as a solid (30 mg, 58%). ¹H NMR (CDCl₃, 400 MHz) δ 1.13 (d, 6H), 1.65-1.69 (m, 2H), 1.83-1.87 (m, 2H), 2.06 (s, 3H), 2.13-2.22 (m, 4H), 2.90-2.93 (m, 2H), 3.28-3.37 (m, 4H), 3.58-3.65 (m, 2H), 3.70-3.79 (m, 4H), 4.81 (hept, 1H), 5.18-5.23 (m, 1H), 7.15-7.18 (m, 1H), 7.59-7.65 (m, 2H), 7.82 (t, 1H), 8.05 (s, 1H), 10.0 (s, 1H). Exact mass calculated for C₂₇H₃₆FN₅O₅ 529.3, found 530.3 (MH⁺).

Example 11.51: Preparation of 4-{6-[2-Fluoro-4-(4-isopropyl-piperazine-1-carbonyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C145).

Compound **C145** was obtained in a similar manner as described in Example 11.36 as a solid (29 mg, 53%). ¹H NMR (CDCl₃, 400 MHz) δ 1.17 (d, 6H), 1.33 (d, 6H), 1.62-1.77 (m, 2H), 1.85-1.95 (m, 6H), 2.10 (s, 3H), 2.70-2.80 (m, 1H), 3.31-3.51 (m, 6H), 3.58-3.69 (m, 2H), 4.84 (hept, 1H), 5.23-5.28 (m, 1H), 7.21-7.26 (m, 3H), 8.09 (s, 1H), 10.2 (s, 1H). Exact mass calculated for C₂₈H₃₈FN₅O₅ 543.3, found 544.5 (MH⁺).

Example 11.52: Preparation of 4-{6-[2-Fluoro-4-(2-morpholin-4-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C146).

Step 1: Preparation of (3-fluoro-4-hydroxy-phenyl)-acetic acid methyl ester.

To a solution of (3-fluoro-4-hydroxy-phenyl)-acetic acid (20 g, 117.5 mmol) in MeOH (150 mL), was H₂SO₄ (3 drops) added. The reaction mixture was heated to reflux and maintained for 2 hours. The reaction was cooled to room temperature and 5 g of NaHCO₃ was added portionwise. The reaction was concentrated under vacuum and dissolved in ether (200 mL). The ether layer was washed with sat. NaHCO₃. The ether layer was dried over MgSO₄, and concentrated under vacuum to afford the desired compound (19.9 g, 92 %) as an oil. The crude compound was used for the next step without further purification. ¹H NMR (400Mz, DMSO-d₆) δ 9.92 (s, 1H), 7.01-7.20 (m, 2H), 3.62 (s, 2H), 3.61 (s, 3H). LCMS 185.1 [MH⁺].

Step 2: Preparation of (4-benzyloxy-3-fluoro-phenyl)-acetic acid methyl ester.

To a solution of (3-fluoro-4-hydroxy-phenyl)-acetic acid methyl ester (15 g, 54.3 mmol) and benzyl bromide (9.28 g, 54.3 mmol) in DMF (50 mL), was K₂CO₃ (7.24 g, 54.3 mmol) added at an ambient temperature. The reaction mixture was heated to 60°C and maintained for 2 hours. The reaction was cooled to room temperature and poured into H₂O (150 mL). The organic compound was extracted with ether (150 mL) and washed with sat NaHCO₃ (100 mL). The ether layer was dried over MgSO₄, and concentrated under vacuum to afford the desired compound (12.9 g, 87.9 %) as a white crystal. The crude compound was used for the next step without further purification. ¹H NMR (400Mz, DMSO-d₆) δ 7.41-7.49 (m, 5H), 7.15-7.23 (m, 2H), 7.02-7.04 (m, 1H), 5.19 (s, 2H), 3.65 (s, 2H), 3.63 (s, 3H). LCMS 273.4 [MH⁺].

Step 3: Preparation of 2-(4-benzyloxy-3-fluoro-phenyl)-ethanol.

To a solution of (4-benzyloxy-3-fluoro-phenyl)-acetic acid methyl ester (7.1 g, 25.7 mmol) in ether (150 mL), was LAH (1.07 g, 28.3 mmol) added portionwise at 0°C. The reaction mixture stirred for 2 hours at the same temperature. The reaction was quenched with H₂O (5 mL) at 0°C. The solid material was filtrated off and washed with ether (50 mL). The ether was dried over MgSO₄ and concentrated under vacuum to afford the desired compound (5.2 g, 82 %) as a white solid. The crude compound was used for the next step without further purification. ¹H NMR (400Mz, DMSO-d₆) δ 7.34-7.48 (m, 5H), 7.09-7.17 (m, 2H), 6.96-6.98 (m, 1H), 5.17 (s, 2H), 4.66 (s, 1H), 3.60 (b, 2H), 2.68 (m, 2H). LCMS 246.3 [MH⁺].

Step 4: Preparation of 1-benzyloxy-4-(2-bromo-ethyl)-2-fluoro-benzene

To a solution of 2-(4-benzyloxy-3-fluoro-phenyl)-ethanol (1.0 g, 4.0 mmol) and CBr₄ (1.5 g, 4.5 mmol) in CH₂Cl₂ (10 mL), was PPh₃ (1.2 g, 4.5 mmol) added portionwise at 0°C. The reaction mixture stirred for 2 hours at the same temperature. The reaction was concentrated under vacuum and the residue was stirred in ether (10 mmol). The solid, mainly triphenylphosphine oxide, was filtrated off and the filtrate was concentrated under vacuum. The residue was purified over SiO₂ to afford the desired compound (1.15 g, 93.5 %) as a white crystal. ¹H NMR (400Mz, DMSO-d₆) δ 7.36-7.49 (m, 5H), 7.17-7.23 (m, 2H), 7.03-7.05 (m, 1H), 5.18 (s, 2H), 3.72 (t, 2H), 3.08 (t, 2H). LCMS 273.4 [MH⁺].

Step 5: Preparation of 2-fluoro-4-(2-morpholin-4-yl-ethyl)-phenol.

To a solution of 1-benzyloxy-4-(2-bromo-ethyl)-2-fluoro-benzene (1.0 g, 3.2 mmol) and morpholine (278 mg, 3.2 mmol) in DMF (5 mL), was K₂CO₃ (432 mg, 3.2 mmol) added. The reaction mixture was heated to 60°C and maintained for 6 hours. The reaction was cooled to room temperature and poured into H₂O (50 mL). The organic compound was extracted with ethyl acetate (50 mL) and dried over MgSO₄. The ethyl acetate layer was concentrated under vacuum and dissolved in methanol (50 mL). The solution was treated with Pd/C (20 mg) and stirred under H₂ (1atm) for 3 hours. The solid material was filtrated off and the filtrate was concentrated under vacuum to afford the desired compound (790 mg, 93 %) as a white oil. The crude compound was used for the next step without further purification. ¹H NMR (400Mz, DMSO-d₆) δ 7.33-7.45 (m, 5H), 7.09-7.15 (m, 2H), 6.97-7.01 (m, 1H), 5.13 (s, 2H), 3.56 (m, 4H), 2.64-2.68 (m, 2H), 2.44-2.50 (m, 2H), 2.39 (b, 2H). LCMS 310.5 [MH⁺].

Step 6: Preparation of 4-{6-[2-Fluoro-4-(2-morpholin-4-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C146).

To a solution of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (270 mg, 0.84 mmol) and 2-fluoro-4-(2-morpholin-4-yl-ethyl)-phenol (225 mg, 0.84 mmol) in DMF (5 mL), was K₂CO₃ (137 mg, 0.84 mmol) added. The reaction mixture was irradiated under microwave for 1 hour at 150°C. The reaction was cooled to room temperature and poured into H₂O (50 mL) and extracted with ethyl acetate (50 mL). The ethyl acetate was dried over MgSO₄, and concentrated under vacuum and purified over SiO₂ to afford Compound C146 (380 mg, 90 %) as a white solid. ¹H NMR (400Mz, DMSO-d₆) δ 8.06 (s, 1H), 7.09 ~ 7.02 (m, 1H), 6.93 ~ 6.91 (m, 2H), 5.12 (m, 1H), 4.61 (m, 1H), 3.46 ~ 3.37 (m, 6H), 3.18 ~ 3.13 (m, 2H), 2.61 ~ 2.57 (m, 2H), 2.38 ~ 2.26 (m, 2H), 2.25 (b, 2H), 1.96 (s, 3H), 1.79 ~ 1.74 (m, 2H), 1.49 ~ 1.45 (m, 2H), 1.03 (d, 6H). LCMS 503.5 [M+1].

Example 11.53: Preparation of 4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C147).

Step 1: Preparation of 1-benzyloxy-2-fluoro-4-(2-methanesulfonyl-ethyl)-benzene

To a solution of 1-benzyloxy-4-(2-bromo-ethyl)-2-fluoro-benzene (1.9 g, 6.25 mmol) in MeOH (150 mL), was NaSCH₃ (439 mg, 6.25 mmol) added at 0°C. The reaction mixture was warmed to room temperature. After stirring for 5 hours, the reaction was concentrated under vacuum. The residue was dissolved in CH₂Cl₂ (10 mL) and mCPBA (2.7 g, 15.6 mmol) was added portionwise at 0°C. The reaction was warmed to room temperature and stirred for 3 hours. The reaction was dissolved with ether (20 mL) and washed with sat. NaHCO₃. The ether layer was dried over MgSO₄, and concentrated under vacuum. The residue was purified over SiO₂ to afford the desired compound (1.92 g, 89.6 %) as a yellowish crystal. ¹H NMR (400Mz, DMSO-d₆) δ 7.49 ~ 7.33 (m, 5H), 7.23 ~ 7.15 (m, 2H), 7.05 ~ 7.03 (m, 1H), 5.15 (s, 2H), 3.42 ~ 3.36 (m, 2H), 2.96 ~ 2.93 (m, 2H), 2.94 (s, 3H). LCMS 309.5 [MH⁺].

Step 2: Preparation of 2-fluoro-4-(2-methanesulfonyl-ethyl)-phenol.

To a solution of 1-benzyloxy-2-fluoro-4-(2-methylsulfonyl-ethyl)-benzene (1.5 g, 4.87 mmol) in MeOH (25 mL) was added Pd/C (50 mg). The reaction was stirred under H₂ (1 atm) for 3 hours. The solid material was filtrated off and the filtrate was concentrated under vacuum to afford the desired compound (981 mg, 92.4 %) as a yellowish solid. The crude compound was used for the next step without further purification. ¹H NMR (400Mz, DMSO-d₆) δ 9.73 (s, 1H), 7.14 ~ 7.11 (m, 1H), 6.94 ~ 6.87 (m, 2H), 3.43 ~ 3.37 (m, 2H), 2.98 (s, 3H), 2.95 ~ 2.91 (m, 2H). LCMS 228.2 [MH⁺].

Step 3: Preparation of 4-{6-[2-fluoro-4-(2-methanesulfonyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C147).

To a solution of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (270 mg, 0.84 mmol) and 2-fluoro-4-(2-methylsulfonyl-ethyl)-phenol (218 mg, 0.84 mmol) in DMF (5 mL), was K₂CO₃ (137 mg, 0.84 mmol) added. The reaction mixture was irradiated under microwave for 1 hour at 150°C. The reaction was cooled to room temperature and poured into H₂O (50 mL) and extracted with ethyl acetate (50 mL). The ethyl acetate was dried over MgSO₄, and concentrated under vacuum and purified over SiO₂ to afford Compound C147 (286 mg, 68 %) as a white solid. ¹H NMR (400Mz, DMSO-d₆) δ 8.25 (s, 1H), 7.39 ~ 7.35 (m, 1H), 7.30 ~ 7.26 (m, 1H), 7.21 ~ 7.18 (m, 1H), 5.31 (m, 1H), 4.79 (m, 1H), 3.67 ~ 3.65 (m, 2H), 3.64 ~ 3.47 (m, 2H), 3.37 ~ 3.31 (m, 2H), 3.08 (m, 2H), 2.97 (s, 3H), 2.15 (s, 3H), 2.00 ~ 1.93 (m, 2H), 1.68 ~ 1.64 (m, 2H), 1.23 (d, 6H). LCMS 496.5 [MH⁺].

Example 11.54: Preparation of 4-{6-[2-Fluoro-4-(2-hydroxy-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C148).

Step 1: Preparation of 4-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-2-fluoro-phenol.

To a solution of 2-(4-benzyloxy-3-fluoro-phenyl)-ethanol (9.2 g, 37.4 mmol) in CH₂Cl₂ (150 mL) and TBDMS-Cl (5.6 g, 37.4 mmol), was added Et₃N (5.2 mmol, 37.4 mmol) portionwise at 0°C. The reaction mixture was warmed to room temperature and stirred for 2 hours at the same temperature. The reaction was washed with H₂O (150 mL). The CH₂Cl₂ was dried over MgSO₄ and

concentrated under vacuum. To a solution of the residue in MeOH (100 mL), was Pd/C (150 mg) added. The reaction was stirred under H₂ (1atm) for 5 hours. The solid material was filtrated off and the filtrate was concentrated under vacuum to afford the desired compound (8.9 g, 88.3 %) as a grayish solid. The crude compound was used for the next step without further purification. ¹H NMR (400Mz, DMSO-d₆) δ 9.53 (s, 1H), 6.99 ~ 6.96 (m, 1H), 6.83 ~ 6.81 (m, 2H), 3.70 (t, 2H), 2.63 (t, 2H), 0.83 (s, 9H), 0.01 (s, 6H).

Step 2: Preparation of 4-{6-[2-fluoro-4-(2-hydroxy-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C148).

To a solution of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (270 mg, 0.84 mmol) and 4-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-2-fluoro-phenol (279 mg, 0.84 mmol) in DMF (5 mL), was K₂CO₃ (137 mg, 0.84 mmol) added. The reaction mixture was irradiated under microwave for 1 hour at 150°C. The reaction was cooled to room temperature and treated with 1.0 M TBAF in THF (0.9 mL). After stirring for 2 hours, the reaction was poured into H₂O (50 mL) and extracted with ethyl acetate (50 mL). The ethyl acetate was dried over MgSO₄, and concentrated under vacuum and purified over SiO₂ to afford the desired compound (321 mg, 89 %) as a white solid. ¹H NMR (400Mz, DMSO-d₆) δ 8.21 (s, 1H), 7.38 ~ 7.37 (m, 1H), 7.29 ~ 7.28 (m, 1H), 7.21 ~ 7.17 (m, 1H), 5.30 (m, 1H), 4.75 (m, 1H), 3.67 ~ 3.65 (m, 2H), 3.64 ~ 3.41 (m, 2H), 3.33 ~ 3.30 (m, 2H), 3.02 (m, 2H), 2.91 (s, 3H), 1.99 ~ 1.93 (m, 2H), 1.66 ~ 1.64 (m, 2H), 1.22 (d, 6H). LCMS 432.6 [MH⁺].

Example 11.55: Preparation of 4-[6-(4-Carboxymethyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C149).

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (1.6 g, 5.32 mmol), (3-fluoro-4-hydroxy-phenyl)-acetic acid (1.8 g, 10.64 mmol), and sodium hydride (638 mg, 26.61 mmol) in dimethylacetamide (18 mL) was heated under microwave irradiation for 1 hr at 150°C. The reaction was quenched with water and the product extracted in ethyl acetate. The organic layer was concentrated in *vacuo* and purified by flash chromatography to provide compound C149 as a white solid (3.4 g, 48%). ¹H NMR (DMSO-d₆, 400 MHz) δ 1.12 (d, 6H), 1.53-1.63 (m, 2H), 1.82-1.92 (m, 2H), 2.07 (s, 3H), 3.22-3.31 (m, 2H), 3.37-3.52 (m, 2H), 3.52-3.61 (m, 2H), 4.71 (h, 1H), 5.19-5.28 (h, 1H), 7.06 (d, 1H), 7.16-7.23 (m, 2H), 8.17 (s, 1H). Exact mass calculated for C₂₂H₂₆FN₃O₆ 447.46, found 448.3 (MH⁺).

Example 11.56: Preparation of 4-[6-(4-Dimethylcarbamoylmethyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C150).

A mixture of 4-[6-(4-carboxymethyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (150 mg, 0.335 mmol) and HATU (178 mg, 0.469 mmol) in DMF (4 mL) was stirred for 30 minutes at room temperature. Then, dimethylamine (235 μL, 0.47

mmol) was added and the reaction was stirred for 24 hours at room temperature. The reaction was quenched with water and the product extracted in ethyl acetate. The organic layer was concentrated in *vacuo* and purified by HPLC to provide Compound C150 as a white solid (40 mg, 25%). ¹H NMR (DMSO-d₆, 400 MHz) δ 1.21 (d, 6H), 1.62-1.72 (m, 2H), 1.92-2.01 (m, 2H), 2.16 (s, 3H), 2.87 (s, 3H), 3.06 (s, 3H), 3.32-3.41 (m, 2H), 3.61-3.69 (m, 2H), 3.75 (s, 2H), 4.08 (h, 1H), 5.33 (h, 1H), 7.08 (d, 1H), 7.23-7.33 (m, 2H), 8.27 (s, 1H). Exact mass calculated for C₂₄H₃₁FN₄O₅, 474.53, found 475.5 (MH⁺).

Example 11.57: Preparation of 4-[6-(2-Fluoro-4-sulfamoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C151).

A solution of 4-[6-(4-bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (475 mg, 1.01 mmol) in 5 mL tetrahydrofuran was cooled to -78°C and n-butyl lithium was added. After stirring for 30 minutes at -78°C, sulfur dioxide was bubbled vigorously through the solution for 10 minutes. Solution was allowed to warm to room temperature and was concentrated on a rotary evaporator. Residue was dissolved in 2 mL tetrahydrofuran and hexane was added until a white solid precipitated. Solid was filtered and dried under high vacuum (white solid, 265 mg). White solid was dissolved in 15 mL methylene chloride and sulfuryl chloride (100 µl, 1.2 mmol) was added. After stirring for 10 minutes at room temperature, mixture was concentrated and residue was dried under high vacuum. Residue was dissolved in 1 mL dioxane, cooled in an ice bath, and 5 mL ammonium hydroxide (28-30% NH₃) was added. After stirring for 5 minutes, mixture was concentrated and purified by HPLC to give Compound C151 as a white solid (46.7 mg, 10%). ¹H NMR (CDCl₃, 400 MHz) δ 1.26-1.27 (d, J = 6.3 Hz, 6H), 1.76-1.84 (m, 2H), 1.97-2.02 (m, 2H), 2.20 (s, 3H), 3.39-3.46 (m, 2H), 3.74-3.80 (m, 2H), 4.89 (s, 2H), 4.89-4.97 (m, 1H), 5.32-5.38 (m, 1H), 7.35-7.39 (m, 1H), 7.75-7.79 (m, 2H), 8.19 (s, 1H). Exact mass calculated for C₂₀H₂₅FN₄O₆S 468.15, found 469.4 (MH⁺).

Example 11.58: Preparation of 4-[6-(2-Fluoro-4-propionylsulfamoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C152).

To a solution of Compound C151 (33.5 mg, 0.0715 mmol) in 3 mL methylene chloride, triethylamine (30 µl, 0.215 mmol) and propionic anhydride (27.7 µl, 0.215 mmol) were added. After stirring at room temperature for 22 hours, solution was concentrated and purified by HPLC. Fractions which contained intermediate were concentrated and dried under high vacuum. Residue was dissolved in 2 mL methanol and sodium bicarbonate (6.9 mg, 0.082 mmol) was added. After stirring for 18 hours at room temperature, solution was concentrated, dissolved in water/acetonitrile and lyophilized to give Compound C152 as a white solid (34.0 mg, 87%). ¹H NMR (DMSO-d₆, 400 MHz) δ 0.85-0.88 (t, J = 7.6, 3H), 1.19-1.20 (d, J = 6.2, 6H), 1.61-1.69 (m, 2H), 1.92-1.99 (m, 4H), 2.14 (s, 3H), 3.33-3.38 (m, 2H), 3.61-3.67 (m, 2H), 4.75-4.81 (m, 1H), 5.29-5.34 (m, 1H), 7.32-7.36

(m, 1H), 7.57-7.67 (m, 2H), 8.27 (s, 1H). Exact mass calculated for $C_{23}H_{29}FN_4O_7S$ 524.17, found 525.2 (MH^+).

Example 11.59: Preparation of 4-[5-Ethynyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C153).

4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-trimethylsilanylethynyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (10 mg, 0.018 mmol) was dissolved in THF (0.60 mL) and MeOH (0.30 mL), 1.0 N NaOH (0.036 mL, 0.036 mmol) was added to the reaction mixture, and it was stirred at room temperature for 1h. Glacial AcOH (0.0041 mL, 0.072 mmol) was then added to give a pH of 5, then the solvents were evaporated in vacuo to give Compound C153 as a crude solid (11 mg) which contained 2 mol eq of AcONa and was of 80% purity (LCMS). LRMS calculated for $C_{22}H_{24}FN_3O_6S$: 477.14. Found: 478.3 ($M+H$)⁺.

Example 11.60: Preparation of 4-{6-[2-Fluoro-4-(2-phosphonooxy-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C154).

A mixture of 4-{6-[2-fluoro-4-(2-hydroxy-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (100 mg, 0.23 mmol), and phosphorus oxychloride (0.105 mL, 1.15 mmol) in dichloroethane (5 mL) was stirred for 5 hours at room temperature. Upon evaporation of the organic solvent, the reaction was quenched with water and the product extracted in ethyl acetate. The organic layer was concentrated in vacuo and purified by HPLC to provide Compound C154 as a white solid (40 mg, 33%). ¹H NMR ($CDCl_3$, 400 MHz) δ 1.23 (d, 6H), 1.71-1.81 (m, 2H), 1.90-2.01 (m, 2H), 2.15 (s, 3H), 2.89 (t, 2H), 3.34-3.44 (m, 2H), 3.66-3.77 (m, 2H), 4.16 (q, 2H), 4.90 (h, 1H), 5.31 (h, 1H), 6.96-7.09 (m, 3H), 7.38-7.58 (s broad, 2H), 8.18 (s, 1H). Exact mass calculated for $C_{22}H_{29}FN_3O_8P$ 513.45, found 514.3 (MH^+).

Example 11.61: Preparation of 4-(6-{2-Fluoro-4-[2-(2-methanesulfonyl-pyrrolidin-1-yl)-2-oxo-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (Compound C156).

Compound C156 was prepared in a similar manner as described in Example 11.56 as a solid (200 mg, 77%). Exact mass calculated for $C_{27}H_{35}FN_4O_7S$ 578.65, found 579.4 (MH^+).

Example 11.62: Preparation of 4-[6-(4-Carbamoylmethyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C157).

Compound C157 was prepared in a similar manner as described in Example 11.56 as a solid (80 mg, 40%). Exact mass calculated for $C_{22}H_{27}FN_4O_5$ 446.47, found 447.6 (MH^+).

Example 11.63: Preparation of 4-[6-(2-Fluoro-4-[(tetrahydro-furan-2-ylmethyl)-carbamoyl]-methyl]-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C158).

Compound C158 was prepared in a similar manner as described in Example 11.56 as a solid (230 mg, 97%). Exact mass calculated for $C_{27}H_{35}FN_4O_6$ 530.59, found 531.5 (MH⁺).

Example 11.64: Preparation of 4-(6-{2-Fluoro-4-[2-(3-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (Compound C166).

Compound C166 was prepared in a similar manner as described in Example 11.56 as a solid (150 mg, 62%). Exact mass calculated for $C_{27}H_{35}FN_4O_5$ 530.59, found 531.3 (MH⁺).

Example 11.65: Preparation of 4-{6-[2-Fluoro-4-(2-morpholin-4-yl-2-oxo-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C167).

Compound C167 was prepared in a similar manner as described in Example 11.56 as a solid (90 mg, 38%). Exact mass calculated for $C_{26}H_{33}FN_4O_6$ 516.56, found 517.4 (MH⁺).

Example 11.66: Preparation of 4-{6-[2-Fluoro-4-(2-imidazol-1-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C168).

General Protocol of Alkylation

A mixture of 4-{6-[4-(2-bromo-ethyl)-2-fluoro-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (30.0 mg, 0.44 mmol) and sodium hydride (11.0 mg, 0.44 mmol) in DMF (4 mL) was stirred for 30 minutes at room temperature. Then, imidazole (200 μ L, 0.40 mmol) was added and the reaction was stirred for 2 hours at room temperature. The reaction was quenched with water and the product extracted in ethyl acetate. The organic layer was concentrated in *vacuo* and purified by HPLC to provide compound C168 as a white solid (28 mg, 15%). ¹H NMR (CDCl₃, 400 MHz) δ 1.17 (d, 6H), 1.66-1.76 (m, 2H), 1.85-1.96 (m, 2H), 2.11 (s, 3H), 3.06-3.14 (m, 2H), 3.30-3.37 (m, 2H), 3.62-3.72 (m, 2H), 3.33-3.40 (m, 2H), 4.86 (h, 1H), 5.24 (h, 1H), 6.80-6.95 (m, 3H), 7.09 (t, 1H), 7.32 (s, 1H), 8.11 (s, 1H), 8.72 (s, 1H). Exact mass calculated for $C_{25}H_{30}FN_5O_4$ 483.54, found 484.4 (MH⁺).

Example 11.67: Preparation of 4-{6-[2-Fluoro-4-(2-[1,2,3]triazol-1-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C169).

Compound C169 was prepared in a similar manner as described in Example 11.66 as a solid (105 mg, 54%). Exact mass calculated for $C_{24}H_{29}FN_6O_4$ 484.52 found 485.4 (MH⁺).

Example 11.68: Preparation of 4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid (R)-tetrahydro-furan-3-yl ester (Compound C177).

To a solution of 1,1'-carbonyldiimidazole (54.8 mg, 0.338 mmol) in 1 mL THF was added (R)-(+)-3-hydroxytetrahydrofuran (32 μ l, 0.38 mmol). After stirring for 30 minutes at room temperature, 1 mL triethylamine, 1 mL THF, and 4-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-(piperidin-4-yloxy)-pyrimidine (70 mg, 0.166 mmol) were added. The resulting mixture was stirred at 60°C for 48 hours and purified by HPLC to give Compound C177 as a white solid (35.3 mg, 42%). ¹H NMR (CDCl₃, 400 MHz) δ 1.81-1.88 (m, 2H), 1.96-2.09 (m, 3H), 2.12-2.20 (m, 4H), 3.10 (s, 3H), 3.42-3.48 (m, 2H), 3.75-3.81 (m, 2H), 3.84-3.97 (m, 4H), 5.27-5.31 (m, 1H), 5.34-5.38 (m, 1H), 7.41-7.45 (m, 1H), 7.77-7.82 (m, 2H), 8.20 (s, 1H). Exact mass calculated for C₂₂H₂₆FN₃O₇S 495.15, found 496.3 (MH⁺).

Example 11.69: Preparation of 4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid (S)-tetrahydro-furan-3-yl ester (Compound C176).

Compound C176 was prepared in a simpler manner as described in Example 11.68 as a white solid (40.8 mg, 46%). ¹H NMR (CDCl₃, 400 MHz) δ 1.81-1.88 (m, 2H), 1.96-2.09 (m, 3H), 2.12-2.20 (m, 4H), 3.10 (s, 3H), 3.42-3.48 (m, 2H), 3.75-3.81 (m, 2H), 3.84-3.97 (m, 4H), 5.27-5.31 (m, 1H), 5.34-5.38 (m, 1H), 7.41-7.45 (m, 1H), 7.77-7.82 (m, 2H), 8.20 (s, 1H). Exact mass calculated for C₂₂H₂₆FN₃O₇S 495.15, found 496.3 (MH⁺).

Example 11.70: Preparation of 4-{6-[2-Fluoro-4-(6-methoxy-pyridin-3-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C181).

A mixture of 4-[6-(4-bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (184 mg, 0.393 mmol), 6-methoxypyridine-3-boronic acid (67.3, 0.396 mmol), potassium carbonate (164 mg, 1.24 mmol), and tetrakis (triphenylphosphine) palladium (20 mg, 0.046 mmol) in 4 mL THF and 0.4 mL H₂O was heated under microwave irradiation for 1 hour at 120°C. The mixture was purified by HPLC to give Compound C181 as a colorless oil (TFA salt, 186 mg, 78%). ¹H NMR (CDCl₃, 400 MHz) δ 1.27-1.29 (d, J = 6.3 Hz, 6H), 1.80-1.88 (m, 2H), 1.98-2.05 (m, 2H), 2.23 (s, 3H), 3.44-3.50 (m, 2H), 3.74-3.80 (m, 2H), 4.12 (s, 3H), 4.93-4.99 (m, 1H), 5.34-5.39 (m, 1H), 5.34-5.39 (m, 1H), 7.06-7.08 (m, 1H), 7.33-7.39 (m, 3H), 8.07-8.10 (m, 1H), 8.27 (s, 1H), 8.57-8.58 (d, J = 2.4 Hz, 1H). Exact mass calculated for C₂₆H₂₉FN₄O₅ 496.21, found 497.4 (MH⁺).

Example 11.71: Preparation of 4-[5-Bromo-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C155).

4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (1.02 g, 2.25 mmol) and NBS (601 mg, 3.38 mmol) in acetic acid (10 mL) was stirred at 40°C for three days. The mixture was purified through HPLC to provide compound **C155** as a solid (685 mg, 50%). ¹H NMR (CDCl₃, 400 MHz) δ 1.24 (d, 6H), 1.82-1.86 (m, 2H), 1.94-1.98 (m, 2H), 3.08 (s, 3H), 3.46-3.52 (m, 2H), 3.68-3.74 (m, 2H), 4.92 (hept, 1H), 5.38-5.42 (m, 1H), 7.42-7.46 (m, 1H), 7.77-7.81 (m, 2H), 8.18 (s, 1H). Exact mass calculated for C₂₀H₂₃FBrN₃O₆S 531.1, found 532.4/534.4 (MH⁺).

Example 11.72: Preparation of 4-[6-(4-Methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C179).

Compound **C179** was prepared in a similar manner as described in Example 11.15 as a solid (60 mg, 67%). ¹H NMR (CDCl₃, 400 MHz) δ 1.26 (d, 6H), 1.77-1.85 (m, 2H), 1.98-2.03 (m, 2H), 2.18 (s, 3H), 3.08 (s, 3H), 3.41-3.47 (m, 2H), 3.74-3.80 (m, 2H), 4.95 (hept, 1H), 5.35-5.39 (m, 1H), 7.30-7.33 (m, 2H), 7.98-8.01 (m, 2H), 8.28 (s, 1H). Exact mass calculated for C₂₁H₂₇N₃O₆S 449.2, found 450.3 (MH⁺).

Example 11.73: Preparation of 4-[6-(2-Amino-4-ethanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C178).

Compound **C178** was prepared in a similar manner as described in Example 11.15 as a solid (66 mg, 69%). ¹H NMR (CDCl₃, 400 MHz) δ 1.11 (t, 3H), 1.20 (d, 6H), 1.59-1.66 (m, 2H), 1.90-1.95 (m, 2H), 2.08 (s, 3H), 3.19 (quart, 2H), 3.30-3.35 (m, 2H), 3.60-3.67 (m, 2H), 4.78 (hept, 1H), 5.24-5.30 (m, 1H), 7.10 (d, 1H), 7.48 (dd, 1H), 7.99 (s, 2H), 8.27 (s, 1H), 8.33 (d, 1H). Exact mass calculated for C₂₂H₃₀N₄O₆S 478.2, found 479.2 (MH⁺).

Example 11.74: Preparation of 4-[5-Methyl-6-(4-sulfo-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C185).

Compound **C185** was prepared in a similar manner as described in Example 11.15 as a solid (21 mg, 23%). ¹H NMR (CDCl₃, 400 MHz) δ 1.20 (d, 6H), 1.63-1.67 (m, 2H), 1.91-1.97 (m, 2H), 2.12 (s, 3H), 3.32-3.36 (m, 2H), 3.62-3.67 (m, 2H), 4.78 (quint, 1H), 5.30-5.32 (m, 1H), 7.06-7.09 (m, 2H), 7.61-7.64 (m, 2H), 8.24 (s, 1H). Exact mass calculated for C₂₀H₂₅N₃O₇S 451.1, found 452.3 (MH⁺).

Example 11.75: Preparation of 4-{6-[2-Fluoro-4-(2-isopropoxy-ethoxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester (Compound C184).

Mixture of 4-(6-Chloro-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester (147 mg, 0.47 mmole), 2-fluoro-4-(2-isopropoxy-ethoxy)-phenol (0.47 mmole, 1 eq) and K₂CO₃ (0.75 mmole, 1.5 eq) in 2mL DMSO was heated under microwave irradiation at 150 °C for 40

minutes. Mixture was purified by HPLC to give compound **C184** as a yellow oil (273 mg, 96%). ¹H NMR (DMSO-d₆, 400 MHz) δ 1.23 (d, 6H), 1.27 (d, 6H), 1.79-1.86 (m, 2H), 1.97-2.04 (m, 1H), 2.20 (s, 3H), 3.43-3.50 (m, 2H), 3.73-3.79 (m, 3H), 3.82 (t, 2H), 4.09 (t, 2H), 4.95 (sep, 1H), 5.33-5.36 (m, 1H), 6.70-6.78 (m, 2H), 7.08 (t, 1H), 8.27 (s, 1H). Exact mass calculated for C₂₅H₃₄FN₃O₆ 491.2, found 492.4 (MH⁺).

Example 11.76: Preparation of 3-tert-Butoxy-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-propan-1-one (Compound C161).

Mixture of 4-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-(piperidin-4-yloxy)-pyrimidine (76 mg, 0.2 mmole), 3-tert-butoxy-propionic acid (0.26 mmole, 1.3 eq), HATU (0.26 mmole, 1.3 eq), and triethylamine (0.4 mmole, 2 eq) in 2 mL DMF was stirred at room temp for 2 hours. Mixture was purified by HPLC to give compound **C161** as a white solid (88 mg, 86 %). ¹H NMR (CDCl₃, 400 MHz) δ 1.20 (s, 9H), 1.91-1.96 (m, 2H), 2.02-2.10 (m, 2H), 2.22 (s, 3H), 2.72 (t, 2H), 3.11 (s, 3H), 3.62-3.64 (m, 2H), 3.72 (t, 2H), 3.86-3.91 (m, 2H), 5.41-5.46 (m, 1H), 7.44 (t, 1H), 7.78-7.82 (m, 2H), 8.23 (s, 1H). Exact mass calculated for C₂₄H₃₂FN₃O₆S 509.2, found 510.6 (MH⁺).

Example 11.77: Preparation of 2-Ethoxy-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone (Compound C163).

Mixture of 4-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-(piperidin-4-yloxy)-pyrimidine (76 mg, 0.2 mmole), 2-ethoxyacetic acid (0.26 mmole, 1.3 eq), HATU (0.26 mmole, 1.3 eq), and TEA (0.4 mmole, 2 eq) in 2 mL THF was heated under microwave irradiation at 120 °C for 30 minutes. Mixture was purified by HPLC to give compound **C163** as a white solid (55 mg, 59%). ¹H NMR (CDCl₃, 400 MHz) δ 1.25 (t, 3H), 1.86-1.90 (m, 2H), 2.01-2.09 (m, 2H), 2.22 (s, 3H), 3.10 (s, 3H), 3.51-3.69 (m, 2H), 3.59 (q, 2H), 3.76-3.87 (m, 2H), 4.21 (s, 2H), 5.41-5.44 (m, 1H), 7.43 (t, 1H), 7.78-7.82 (m, 2H), 8.21 (s, 1H). Exact mass calculated for C₂₁H₂₆FN₃O₆S 467.2, found 468.5 (MH⁺).

Example 11.78: Preparation of {4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-(tetrahydro-furan-2-yl)-methanone (Compound C164).

Mixture of 4-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-(piperidin-4-yloxy)-pyrimidine (76 mg, 0.2 mmole), tetrahydro-furan-2-carboxylic acid (0.26 mmole, 1.3 eq), HATU (0.26 mmole, 1.3 eq), and TEA (0.4 mmole, 2 eq) in 2 mL THF was heated under microwave irradiation at 120 °C for 30 minutes. Mixture was purified by HPLC to give compound **C164** as a yellow solid (78 mg, 81%). ¹H NMR (CDCl₃, 400 MHz) δ 1.85-2.17 (m, 8H), 2.21 (s, 3H), 3.10 (s, 3H), 3.75-3.80 (m, 2H), 3.86-3.91 (m, 2H), 3.96-4.01 (m, 2H), 4.68 (t, 1H), 5.40-5.45 (m, 1H), 7.44 (t, 1H), 7.78-7.82 (m, 2H), 8.21 (s, 1H). Exact mass calculated for C₂₂H₂₆FN₃O₆S 479.2, found 480.3 (MH⁺).

Example 11.79: Preparation of (S)-1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-2-methylamino-butan-1-one (Compound C165).

Compound C165 was obtained in a similar manner as described in Example 11.78 as a yellow solid (7 mg, 7%). Exact mass calculated for $C_{23}H_{31}FN_4O_5S$ 494.2, found 495.5 (MH^+).

Example 11.80: Preparation of (S)-1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-hydroxy-butan-1-one (Compound C171).

Compound C171 was obtained in a similar manner as described in Example 11.78 as a white solid (31 mg, 33 %). Exact mass calculated for $C_{21}H_{26}FN_3O_6S$ 467.2, found 468.6 (MH^+).

Example 11.81: Preparation of (R)-1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-2-methylamino-butan-1-one (Compound C170).

Compound C170 was obtained in a similar manner as described in Example 11.78 as a yellow solid (16 mg, 16%). Exact mass calculated for $C_{23}H_{31}FN_4O_5S$ 494.2, found 495.5 (MH^+).

Example 11.82: Preparation of (R)-N-(1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carbonyl}-2-methyl-propyl)-acetamide (Compound C172).

Compound C172 was obtained in a similar manner as described in Example 11.78 as a white solid (83 mg, 80%). 1H NMR (CD_3OD , 400 MHz) δ 0.95-0.99 (m, 6H), 1.78-1.91 (m, 2H), 2.00 (s, 3H), 2.04-2.05 (m, 2H), 2.23 (d, 3H), 3.19 (s, 3H), 3.58-3.64 (m, 2H), 3.77 (m, 2H), 4.02 (m, 2H), 4.70 (d, 1H), 5.45-5.47 (m, 1H), 7.54 (t, 1H), 7.84-7.88 (m, 2H), 8.16 (s, 1H). Exact mass calculated for $C_{24}H_{31}FN_4O_6S$ 522.2, found 523.5 (MH^+).

Example 11.83: Preparation of (S)-N-(1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carbonyl}-2-methyl-propyl)-acetamide (Compound C173).

Compound C173 was obtained in a similar manner as described in Example 11.78 as a white solid (89 mg, 80%). 1H NMR (CD_3OD , 400 MHz) δ 0.96-0.99 (m, 6H), 1.77-1.82 (m, 2H), 2.00 (s, 3H), 2.04-2.10 (m, 2H), 2.23 (d, 3H), 3.19 (s, 3H), 3.75-3.77 (m, 2H), 3.86-3.88 (m, 2H), 4.02 (m, 2H), 4.70 (d, 1H), 5.45-5.47 (m, 1H), 7.55 (t, 1H), 7.85-7.89 (m, 2H), 8.17 (s, 1H). Exact mass calculated for $C_{24}H_{31}FN_4O_6S$ 522.2, found 523.5 (MH^+).

Example 11.84: Preparation of (R)-N-(2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-methyl-2-oxo-ethyl)-acetamide (Compound C174).

Compound C174 was obtained in a similar manner as described in Example 11.78 as a white solid (84 mg, 85%). 1H NMR (CD_3OD , 400 MHz) δ 1.32 (d, 3H), 1.79-1.91 (m, 2H), 1.98 (s, 3H), 2.04-2.10 (m, 2H), 2.23 (d, 3H), 3.20 (s, 3H), 3.75-3.99 (m, 2H), 3.86-3.88 (m, 1H), 4.02 (m, 2H),

5.46-5.48 (m, 1H), 7.55 (t, 1H), 7.85-7.89 (m, 2H), 8.17 (s, 1H). Exact mass calculated for $C_{22}H_{27}FN_4O_6S$ 494.2, found 495.5 (MH^+).

Example 11.85: Preparation of (S)-N-(2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-methyl-2-oxo-ethyl)-acetamide (Compound C175).

Compound C175 was obtained in a similar manner as described in Example 11.78 as a white solid (81 mg, 82%). 1H NMR (CD_3OD , 400 MHz) δ 1.32 (d, 3H), 1.82-1.99 (m, 2H), 2.01 (s, 3H), 2.05-2.15 (m, 2H), 2.24 (d, 3H), 3.20 (s, 3H), 3.46-3.55 (m, 2H), 3.74-3.81 (m, 1H), 3.92-4.02 (m, 2H), 5.45-5.48 (m, 1H), 7.55 (t, 1H), 7.85-7.90 (m, 2H), 8.17 (s, 1H). Exact mass calculated for $C_{22}H_{27}FN_4O_6S$ 494.2, found 495.5 (MH^+).

Example 11.86: Preparation of 3-Amino-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-methyl-pentan-1-one (Compound C182).

Compound C182 was obtained in a similar manner as described in Example 11.78 as a white solid (3 mg, 3%). Exact mass calculated for $C_{23}H_{31}FN_4O_5S$ 494.2, found 495.5 (MH^+).

Example 11.87: Preparation of (1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carbonyl}-2-methyl-propyl)-carbamic acid tert-butyl ester (Compound C180).

Compound C180 was obtained in a similar manner as described in Example 11.78 as a yellow solid (143 mg, 88%). Exact mass calculated for $C_{27}H_{37}FN_4O_7S$ 580.2, found 581.4 (MH^+).

Example 11.88: Preparation of 2-Amino-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-butan-1-one (Compound C183).

Mixture of compound C180 (68 mg, 0.12 mmole) in 4M HCl in Dioxane (1.5 mL) and dioxane (2 mL) was stirred at room temperature for 40 minutes. Mixture was purified by HPLC to give Compound C183 as a white solid (50 mg, 86%). 1H NMR (CD_3OD , 400 MHz) δ 1.04 (d, 3H), 1.13 (d, 3H), 1.87-1.92 (m, 2H), 2.04-2.14 (m, 3H), 2.24 (d, NH_2), 2.82 (s, 3H), 3.20 (s, 3H), 3.52-3.65 (m, 2H), 3.81-3.83 (m, 2H), 4.36 (m, 1H), 5.49-5.51 (m, 1H), 7.55 (t, 1H), 7.86-7.90 (m, 2H), 8.18 (s, 1H). Exact mass calculated for $C_{22}H_{29}FN_4O_5S$ 480.2, found 481.3 (MH^+).

Example 11.89: Preparation of 4-[6-(3-Fluoro-biphenyl-4-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C236).

Compound C236 was made in a similar manner as described in Example 11.70 as a white solid (51 mg, 55 %). 1HNMR ($CDCl_3$, 400 MHz) δ 1.28 (d, 6H), 1.81-1.86 (m, 2H), 1.99-2.04 (m, 2H), 2.23 (s, 3H), 3.44-3.50 (m, 2H), 3.74-3.80 (m, 2H), 4.96 (sep, 1H), 5.34-5.38 (m, 1H), 7.27 (t,

1H), 7.36-7.47 (m, 5H), 7.57 (d, 2H), 8.28 (s, 1H). Exact mass calculated for $C_{26}H_{28}FN_3O_4$ 465.2, found 466.5.

5 **Example 11.90: Preparation of 4-[6-(2-Fluoro-4-pyridin-3-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C237).**

Compound C237 was made in a similar manner as described in Example 11.70 as a yellow solid (11 mg, 12 %). 1H NMR ($CDCl_3$, 400 MHz) δ 1.27 (d, 6H), 1.80-1.83 (m, 2H), 1.98-2.02 (m, 2H), 2.23 (s, 3H), 3.41-3.47 (m, 2H), 3.74-3.79 (m, 2H), 4.94 (sep, 1H), 5.34-5.38 (m, 1H), 7.43 (t, 1H), 7.48 (d, 2H), 7.90-7.94 (m, 1H), 8.23 (s, 1H), 8.43-8.46 (m, 1H), 8.83 (d, 1H), 9.12 (s, 1H).
10 Exact mass calculated for $C_{25}H_{27}FN_4O_4$ 466.2, found 467.6.

Example 11.91: Preparation of 4-[6-(2-Fluoro-4-pyridin-4-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C235).

Compound C235 was made in a similar manner as described in Example 11.70 as a yellow
15 solid (13 mg, 14 %). 1H NMR ($CDCl_3$, 400 MHz) δ 1.27 (d, 6H), 1.79-1.83 (m, 2H), 1.98-2.04 (m, 2H), 2.23 (s, 3H), 3.40-3.47 (m, 2H), 3.75-3.79 (m, 2H), 4.95 (sep, 1H), 5.34-5.38 (m, 1H), 7.48 (t, 1H), 7.58-7.62 (m, 2H), 8.01 (d, 2H), 8.23 (s, 1H), 8.90 (d, 2H). Exact mass calculated for $C_{25}H_{27}FN_4O_4$ 466.2, found 467.6.

20 **Example 11.92: Preparation of 4-[6-(2-Fluoro-4-thiophen-3-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C239).**

Compound C239 was made in a similar manner as described in Example 11.70 as a white solid (29 mg, 31 %). 1H NMR ($CDCl_3$, 400 MHz) δ 1.27 (d, 6H), 1.78-1.83 (m, 2H), 1.97-2.01 (m, 2H), 2.21 (s, 3H), 3.40-3.47 (m, 2H), 3.73-3.79 (m, 2H), 4.94 (sep, 1H), 5.34-5.36 (m, 1H), 7.22 (t, 25 1H), 7.34 (d, 1H), 7.39-7.45 (m, 4H), 8.24 (s, 1H). Exact mass calculated for $C_{24}H_{26}FN_3O_4S$ 471.2, found 472.4.

Example 11.93: Preparation of 4-[6-(2-Fluoro-4-pyrimidin-5-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C238).

30 Compound C238 was made in a similar manner as described in Example 11.70 as a white solid (10 mg, 11 %). Exact mass calculated for $C_{24}H_{26}FN_5O_4$ 467.2, found 468.6.

Example 11.94: Preparation of 4-[6-[2-Fluoro-4-(5-methoxy-pyridin-3-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C234).

35 Compound C234 was made in a similar manner as described in Example 11.70 as a white solid (TFA salt, 192 mg, 71%). 1H NMR ($CDCl_3$, 400 MHz) 1.26-1.28 (d, J = 6.3 Hz, 6H), 1.77-1.83 (m, 2H), 1.98-2.06 (m, 2H), 2.23 (s, 3H), 3.40-3.46 (m, 2H), 3.76-3.85 (m, 2H), 4.04 (s, 3H), 4.92-

4.98 (m, 1H), 5.33-5.38 (m, 1H), 7.37-7.45 (m, 3H), 7.77-7.78 (m, 1H), 8.23-8.24 (d, J = 4.67, 1H), 8.45-8.46 (d, J = 2.3 Hz, 1H), 8.61 (s, 1H). Exact mass calculated for $C_{26}H_{29}FN_4O_5$ 496.21, found 497.4.

5 **Example 11.95: Preparation of 4-[6-(4-Ethynyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (Compound C240).**

Pd(II)(PhCN)₂Cl₂ (15 mg, 0.039 mmol) and CuI (9 mg, 0.047 mmol) were dissolved in anhydrous dioxane (3 mL), 10 wt % P(t-Bu)₃ in hexanes (0.200 mL, 13.6 mg, 0.067 mmol), NH(i-Pr)₂ (0.085 mL, 0.60 mmol), 4-[6-(4-bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester (243 mg, 0.50 mmol), and TMS-acetylene (0.083 mL, 0.60 mmol),
10 were added and the reaction mixture was sealed, purged with nitrogen, and stirred at room temperature for 12 h. The reaction mixture was diluted with EtOAc (15 mL), the mixture was filtered through a silica pad, the solvent was evaporated in vacuo at 23 C to give a dark oil which was dissolved in THF (3.5 mL) and MeOH (1.5 mL). 1.0 N NaOH (0.75 mL, 0.75 mmol) was added and
15 after 5 min glacial AcOH (0.086 mL, 1.5 mmol) was added. After addition of silica (2.4 g) the solvent was evaporated in vacuo at 25 C to give a solid which was pulverized with mortar and pestle. This crude product was adsorbed onto silica and purified by flash chromatography using hexanes-EtOAc, 82:18, then hexanes-EtOAc, 75:25, v/v, to give the title Compound C240 as a resin (67 mg, 32% over 2 steps). ¹H NMR (400 HMz, CDCl₃) δ 1.25 (d, J = 6 Hz, 6H), 1.78 (m, 2H), 1.98 (m, 2H),
20 2.17 (s, 3H), 3.07 (s, 1H), 3.40 (m, 2H), 3.74 (m, 2H), 4.92 (m, 1H), 5.32 (m, 1H), 7.14 (m, 1H), 7.29 (m, 2H), 8.18 (s, 1H). LRMS calculated for $C_{22}H_{24}FN_3O_4$: 413.18. Found: 414.5 (M+H).

EXAMPLE 12

25 **SYNTHESES OF COMPOUNDS OF THE PRESENT INVENTION**

Example 12.1: Preparation of 4-({Cyclopropyl-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-amino}-methyl)-piperidine-1-carboxylic acid tert-butyl ester (Compound D1).

A mixture of 4-{{ (6-chloro-5-methyl-pyrimidin-4-yl)-cyclopropyl-amino}-methyl}-piperidine-1-carboxylic acid *tert*-butyl ester (200mg, 0.5mmole), potassium carbonate (208mg, 1.5mmole), and 2-fluoro-4-methanesulfonyl-phenol (95mg, 0.5mmole) in 2mL DMF was heated under microwave irradiation at 160° C for 4 hours. Mixture was purified by HPLC to give compound D1 as a yellow solid (93 mg, 35%). ¹H NMR (CDCl₃, 400 MHz) δ 0.69-0.73 (m, 2H), 0.97-1.02 (m, 2H), 1.10-1.17 (m, 2H), 1.45 (s, 9H), 1.71 (d, J = 12.4 Hz, 2H), 1.98-2.07 (m, 1H), 2.38 (s, 3H), 2.72 (t, J = 12.4 Hz, 2H), 3.07-3.11 (m, 1H), 3.12 (s, 3H), 3.52 (d, J = 7.3 Hz, 2H), 4.12 (d, J = 12.9 Hz, 2H), 7.45 (t, J = 8.3 Hz, 1H), 7.80-7.84 (m, 2H), 8.34 (s, 1H). Exact mass calculated for $C_{26}H_{35}FN_4O_5S$ 534.2, found 535.4 (MH⁺).

Example 12.2: Preparation of 4-({Cyclopropyl-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-amino}-methyl)-piperidine-1-carboxylic acid isopropyl ester (Compound D2)

5 A mixture of cyclopropyl-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-piperidin-4-ylmethyl-amine (65mg, 0.15mmole) and triethylamine (23mg, 0.225mmole) in 2 mL THF was stirred at room temperature for 10 minutes. Into this mixture was added drop-wise isopropyl chloroformate (0.225mmole). The mixture was quenched with water, extracted with ethyl acetate, and dried in vacuum to give Compound **D2** as a yellow solid (65 mg, 83%). ¹H NMR (CDCl₃, 400 MHz) δ 0.69-0.64 (m, 2H), 0.83-0.88 (m, 2H), 1.12-1.15 (m, 2H), 1.23 (d, J = 6.3 Hz, 6H), 1.72 (d, J = 13.1 Hz, 2H), 1.94-2.02 (m, 1H), 2.04 (s, 3H), 2.73 (t, J = 12.4 Hz, 2H), 2.97-3.17 (m, 1H), 3.10 (s, 3H), 3.49-3.55 (m, 2H), 4.09-4.23 (m, 2H), 4.90 (sept, J = 6.3 Hz, 1H), 7.45 (t, J = 8.3 Hz, 1H), 7.76-7.80 (m, 2H), 8.12 (s, 1H). Exact mass calculated for C₂₅H₃₃FN₄O₅S 520.2, found 521.5 (MH⁺).

15

EXAMPLE 13

SYNTHESES OF COMPOUNDS OF THE PRESENT INVENTION

20 **Example 13.1: Preparation of 4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-ylsulfanyl]-piperidine-1-carboxylic acid isopropyl ester (Compound E1).**

Step 1: Preparation of 4-(6-chloro-5-methyl-pyrimidin-4-ylsulfanyl)-piperidine-1-carboxylic acid *tert*-butyl ester.

25 A mixture of 4-mercapto-piperidine-1-carboxylic acid *tert*-butyl ester (1.5352 g, 7.06 mmol) and 4,6-dichloro-5-methyl-pyrimidine (1.1512 g, 7.06 mmol) in 15 mL of THF with sodium *t*-butoxide (1M in THF, 8.3 mL, 8.3 mmol) added dropwise. After 5 min, mixture was concentrated and residue was extracted with CH₂Cl₂ and H₂O. Organic phase was dried over MgSO₄, filtered and concentrated to give 4-(6-chloro-5-methyl-pyrimidin-4-ylsulfanyl)-piperidine-1-carboxylic acid *tert*-butyl ester as a yellowish solid (2.3469 g, 97%). Exact mass calculated for C₁₅H₂₂ClN₃O₂S 343.11, found 344.1 (MH⁺).

30 **Step 2: Preparation of 4-chloro-5-methyl-6-(piperidin-4-ylsulfanyl)-pyrimidine.**

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-ylsulfanyl)-piperidine-1-carboxylic acid *tert*-butyl ester (2.3469 g, 6.82 mmol) and 40 mL of 4M HCl in dioxane was stirred at room temperature over night. Mixture was concentrated to give 4-chloro-5-methyl-6-(piperidin-4-ylsulfanyl)-pyrimidine as a yellowish solid (1.8985 g, 99%). Exact mass calculated for C₁₀H₁₄ClN₃S 243.06, found 244.1 (MH⁺).

35

Step 3: Preparation of 4-(6-chloro-5-methyl-pyrimidin-4-ylsulfanyl)-piperidine-1-carboxylic acid isopropyl ester.

A mixture of 4-chloro-5-methyl-6-(piperidin-4-ylsulfanyl)-pyrimidine (HCl salt, 1.8985 g, 6.77 mmol) and triethylamine (2.825 mL, 0.02 mol) in 50 mL of CH₃CN was stirred under room temperature. After 15 min, isopropyl chloroformate (1M in toluene, 8.13 mL, 8.13 mmol) were added slowly under 0°C. Mixture was stirred under room temperature. After 3 h, mixture was concentrated and residue was extracted with EtOAc and saturated NaHCO₃. Organic phase was dried over MgSO₄, filtered and concentrated to give 4-(6-chloro-5-methyl-pyrimidin-4-ylsulfanyl)-piperidine-1-carboxylic acid isopropyl ester as yellowish oil (1.9143 g, 85%). Exact mass calculated for C₁₄H₂₀ClN₃O₂S 329.1, found 330.3 (MH⁺).

Step 4: Preparation of 4-[6-(2-fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-ylsulfanyl]-piperidine-1-carboxylic acid isopropyl ester (Compound E1).

A mixture of 4-(6-chloro-5-methyl-pyrimidin-4-ylsulfanyl)-piperidine-1-carboxylic acid isopropyl ester (1.2234 g, 3.7 mmol), 2-fluoro-4-methanesulfonyl-phenylamine (702 mg, 3.7 mmol), palladium acetate (84.3 mg, 0.37 mmol), 2-(di-*t*-butylphosphino)biphenyl (11 mg, 0.037 mmol), and sodium *tert*-butoxide (891.8 mg, 9.28 mmol) in 15 mL 1,4-dioxane was heated in microwave for 2 hours at 120°C. The mixture was purified by HPLC to give Compound E1 as a tanned solid (TFA salt, 601.1 mg, 27%). ¹H NMR (MeOH-d₄, 400 MHz) δ 1.16-1.17 (d, J = 6.32 Hz, 6H), 1.49-1.58 (m, 2H), 2.00-2.04 (m, 2H), 2.14 (s, 3H), 3.07 (s, 3H), 3.21-3.23 (m, 2H), 3.90-3.94 (m, 2H), 4.01-4.08 (m, 1H), 4.75-4.81 (m, 1H), 7.66-7.68 (d, J = 8.08 Hz, 2H), 8.01-8.05 (m, 1H), 8.27 (s, 1H). Exact mass calculated for C₂₁H₂₇FN₄O₄S₂ 482.15, found 483.4 (MH⁺).

EXAMPLE 14

Protocol for RUP3 Dose Responses in Melanophores

Melanophores are maintained in culture as reported by Potenza, M. N. and Lerner, M. R., in Pigment Cell Research, Vol. 5, 372-378, 1992 and transfected with the RUP3 expression vector (pCMV) using electroporation. Following electroporation, the transfected cells are plated into 96 well plates for the assay. The cells are then allowed to grow for 48 hours in order to both recover from the electroporation procedure and attain maximal receptor expression levels.

On the assay day, the growth medium on the cells is replaced with serum-free buffer containing 10nM melatonin. The melatonin acts via an endogenous Gi-coupled GPCR in the melanophores to lower intracellular cAMP levels. In response to lowered cAMP levels, the melanophores translocate their pigment to the center of the cell. The net effect of this is a significant decrease in the absorbance reading of the cell monolayer in the well, measured at 600-650nm.

After a 1-hour incubation in melatonin, the cells become completely pigment-aggregated. At this point a baseline absorbance reading is collected. Serial dilutions of test compounds are then added to the plate and compounds that stimulate RUP3 produce increases in intracellular cAMP levels. In response to these increased cAMP levels, the melanophores translocate their pigment back into the

cell periphery. After one hour, stimulated cells are fully pigment-dispersed. The cell monolayer in the dispersed state absorbs much more light in the 600-650nm range. The measured increase in absorbance compared to the baseline reading allows one to quantitate the degree of receptor stimulation and plot a dose-response curve.

- 5 The compounds in the above examples were screened using the melanophore assay. Representative compounds and their corresponding EC₅₀ values are shown in the Table 8 below:

Table 8

Compound	RUP3 (EC ₅₀) (nM)
A11	86
A14	242
A24	185
A27	76.5
A32	43.5
A39	16.9
A90	52
B4	300
C168	28.3

- 10 Other compounds in the Examples showed EC₅₀ activities in the membrane cyclase assay less than about 10 μ M.

EXAMPLE 15

Food intake study

- 15 Male ZDF (Zucker diabetic fatty) rats weighing 350g-400g were dosed independently with two structurally divergent chemotypes exhibiting agonism to the RUP3 receptor. Rats were dosed daily via oral gavage with either vehicle (100% PEG 400), First Compound (30mg/kg, 100mg/kg), or Second Compound (10mg/kg, 30mg/kg) at a volume of 3ml/kg. Body weight and food intake were monitored and recorded daily. The table shown below illustrates the body weight (g) and cumulative
- 20 food intake (g) taken after both seven days and 14 days of dosing.


Substance	Dose (mg/Kg)	Cumulative Food Intake (g)		Body Weight (g)	
		Week 1	Week 2	Week 1	Week 2
First Compound	Vehicle	321	672	390	395
	30 mg/Kg	271	557	383	383

	100 mg/Kg	211	457	361	376
Second Compound	Vehicle	261	563	393	393
	10 mg/Kg	217	459	388	390
	30 mg/Kg	159	307	377	373

Those skilled in the art will recognize that various modifications, additions, substitutions, and variations to the illustrative examples set forth herein can be made without departing from the spirit of the invention and are, therefore, considered within the scope of the invention. All documents referenced
5 above, including, but not limited to, printed publications, and provisional and regular patent applications, are incorporated herein by reference in their entirety.

We claim:

1. A compound of Formula (I):



or a pharmaceutically acceptable salt, hydrate or solvate, or N-oxide thereof;
wherein:

A and B are each independently C₁₋₃ alkylene optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₃ alkyl, C₁₋₄ alkoxy, carboxy, cyano, C₁₋₃ haloalkyl and halogen;

D is O, S, S(O), S(O)₂, CR₂R₃ or N-R₂;

E is N, C or CR₄;

-- is a single bond when E is N or CR₄, or a double bond when E is C;

V₁ is selected from the group consisting of C₁₋₃ alkylene, ethynylene and C₁₋₂ heteroalkylene optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₃ alkyl, C₁₋₄ alkoxy, carboxy, cyano, C₁₋₃ haloalkyl and halogen; or V₁ is a bond;

V₂ is C₃₋₆ cycloalkylene or C₁₋₃ alkylene wherein each are optionally substituted with 1 to 4 substituents selected from the group consisting of C₁₋₃ alkyl, C₁₋₄ alkoxy, carboxy, cyano, C₁₋₃ haloalkyl and halogen; or V₂ is a bond;

W is NR₅, O, S, S(O) or S(O)₂; or W is absent;

Q is NR₆, O, S, S(O) or S(O)₂;


X is N or CR₇;

Y is N or CR₈;

Z is selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, C₁₋₂ alkylamino, C₂₋₄ dialkylamino, carbamimidoyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₄₋₈ diacylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₂₋₆ dialkylsulfonylamino, formyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylcarboxamide, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, halogen, aryl, heterocyclic, heteroaryl, hydroxyl, hydroxycarbamimidoyl, hydroxylamino, nitro and tetrazolyl, wherein C₁₋₈ alkyl, C₃₋₇ cycloalkyl, and heterocyclic are each optionally substituted with 1, 2, 3 or 4 groups selected

from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, amino, C₁₋₂ alkylamino, C₂₋₄ dialkylamino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, formyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro, and wherein said C₁₋₇ alkyl is optionally substituted with amino; or

Z is a group of Formula (A):




wherein:

R₉ is H, C₁₋₈ alkyl or C₃₋₇ cycloalkyl; and

R₁₀ is H, nitro or nitrile;

Ar₁ is aryl or heteroaryl each optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅; wherein R₁₁ is selected from the group consisting of C₁₋₅ acyl, C₁₋₆ acylsulfonamide, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, arylsulfonyl, carbamimidoyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, guanidiny, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, heterocyclic, heterocyclic-oxy, heterocyclicsulfonyl, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, hydroxyl, nitro, C₄₋₇ oxo-cycloalkyl, phenoxy, phenyl, sulfonamide, sulfonic acid, and thiol, and wherein C₁₋₅ acyl, C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, arylsulfonyl, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, heteroaryl, phenoxy and phenyl are optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, heteroaryl, heterocyclic, hydroxyl, nitro, phenyl, and phosphonoxy, wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; or

R₁₁ is a group of Formula (B):



wherein:

“p” and “r” are each independently 0, 1, 2 or 3; and R₁₆ is H, C₁₋₅ acyl, C₂₋₆ alkenyl, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, halogen, heteroaryl or phenyl, and wherein the heteroaryl or phenyl optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₄ alkoxy, amino, C₁₋₄ alkylamino, C₂₋₆ alkynyl, C₂₋₈ dialkylamino, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl and hydroxyl; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, hydroxyl and nitro; or

two adjacent groups selected from the group consisting of R₁₂, R₁₃, R₁₄ and R₁₅ together with the atoms to which they are attached form a 5-, 6- or 7-membered cycloalkyl, cycloalkenyl or heterocyclic group fused with Ar₁, wherein the 5-, 6- or 7-membered group is optionally substituted with halogen;


R₁, R₇ and R₈ are each independently selected from the group consisting of H, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, amino, C₁₋₄ alkylamino, C₂₋₈ dialkylamino, carboxamide, cyano, C₃₋₇ cycloalkyl, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylsulfonamide, halogen, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio and hydroxyl;

R₂ is selected from the group consisting of C₁₋₈ alkyl, amino, aryl, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl and hydroxyl; and wherein C₁₋₈ alkyl, aryl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₃₋₆-cycloalkyl-C₁₋₃-heteroalkylene, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄

haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino and nitro; or

R₂ is -Ar₂-Ar₃ wherein Ar₂ and Ar₃ are each independently aryl or heteroaryl optionally substituted with 1 to 5 substituents selected from the group consisting of H, C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, C₁₋₄ alkylamino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₆-cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, hydroxyl and nitro; or


R₂ is a group of Formula (C):



wherein:

R₁₇ is H, C₁₋₈ alkyl, C₃₋₇ cycloalkyl, aryl, heteroaryl or OR₁₉; and R₁₈ is F, Cl, Br, CN or NR₂₀R₂₁; where R₁₉ is H, C₁₋₈ alkyl or C₃₋₇ cycloalkyl, and R₂₀ and R₂₁ are each independently H, C₁₋₈ alkyl, C₃₋₇ cycloalkyl, aryl or heteroaryl; or

R₂ is a group of Formula (D):



wherein:

G is:

- i) -C(O)-, -C(O)NR₂₃-, -C(O)O-, -OC(O)NR₂₃-, -NR₂₃C(O)O-, -OC(O)-, -C(S)-, -C(S)NR₂₃-, -C(S)O-, -OC(S)-, -CR₂₃R₂₄-, -O-, -S-, -S(O)- or -S(O)₂- when D is CR₂R₃, or
- ii) -CR₂₃R₂₄C(O)-, -C(O)-, -CR₂₃R₂₄C(O)NR₂₅-, -C(O)NR₂₃-, -C(O)O-, -C(S)-, -C(S)NR₂₃-, -C(S)O-, -CR₂₃R₂₄-, -S(O)₂-, or a bond when D is NR₂,

wherein R₂₃, R₂₄ and R₂₅ are each independently H or C₁₋₈ alkyl; and R₂₂ is H, C₁₋₈ alkyl, C₂₋₆ alkynyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio,


halogen, heteroaryl, heterocyclic, hydroxyl, hydroxylamino, nitro, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, heteroaryl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylthiocarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylthiourey, C₁₋₄ alkylureyl, amino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino, and nitro;

R₃ is H, C₁₋₈ alkyl, C₁₋₄ alkoxy or hydroxyl; and

R₄, R₅ and R₆ are each independently H, C₁₋₈ alkyl or C₃₋₇ cycloalkyl, wherein said C₁₋₈ alkyl is optionally substituted with C₁₋₄ alkoxy, C₃₋₇ cycloalkyl, or heteroaryl.

2. The compound according to claim 1 wherein X is N.
3. The compound according to claim 1 wherein Y is N.
4. The compound according to claim 1 wherein X and Y or both N.
5. The compound according to claim 1 wherein --- is a single bond.
6. The compound according to claim 1 wherein W is NH.
7. The compound according to claim 1 wherein W is O.
8. The compound according to claim 1 wherein Q is NR₆.
9. The compound according to claim 5 wherein R₆ is C₁₋₈ alkyl.
10. The compound according to claim 5 wherein R₆ is C₃₋₇ cycloalkyl.
11. The compound according to claim 5 wherein R₆ is H.
12. The compound according to claim 1 wherein Q is O.
13. The compound according to claim 1 wherein Q is S.

14. The compound according to claim 1 wherein both A and B are $-\text{CH}_2-$.
15. The compound according to claim 1 wherein A is $-\text{CH}_2\text{CH}_2-$ and B is $-\text{CH}_2-$.
16. The compound according to claim 1 wherein both A and B are $-\text{CH}_2\text{CH}_2-$.
17. The compound according to claim 1 wherein V_2 is a bond.
18. The compound according to claim 1 wherein V_2 is a $-\text{CH}_2-$ or $-\text{CH}_2\text{CH}_2-$ group.
19. The compound according to claim 1 wherein V_1 is a bond.
20. The compound according to claim 1 wherein D is N-R_2 .
21. The compound according to claim 1 wherein R_2 is a group of Formula (D):



(D)

wherein:

G is $-\text{CR}_{23}\text{R}_{24}\text{C}(\text{O})-$, $-\text{C}(\text{O})-$, $-\text{CR}_{23}\text{R}_{24}\text{C}(\text{O})\text{NR}_{23}-$, $-\text{C}(\text{O})\text{NR}_{23}-$, $-\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{S})-$, $-\text{C}(\text{S})\text{NR}_{23}-$, $-\text{C}(\text{S})\text{O}-$, $-\text{CR}_{23}\text{R}_{24}-$, $-\text{S}(\text{O})_2-$, or a bond; wherein R_{23} and R_{24} are each independently H or C_{1-8} alkyl; and

R_{22} is H, C_{1-8} alkyl, C_{3-7} cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-5} acyl, C_{1-5} acyloxy, C_{2-6} alkenyl, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylamino, C_{1-4} alkylcarboxamide, C_{1-4} alkylthiocarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, C_{1-4} alkylthiourey, C_{1-4} alkylureyl, amino, carbo- C_{1-6} -alkoxy, carboxamide, carboxy, cyano, C_{3-7} cycloalkyl, C_{2-8} dialkylamino, C_{2-6} dialkylcarboxamide, C_{2-6} dialkylthiocarboxamide, C_{2-6} dialkylsulfonamide, C_{1-4} alkylthiourey, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkylsulfinyl, C_{1-4} haloalkylsulfonyl, C_{1-4} haloalkyl, C_{1-4} haloalkylthio, halogen, heteroaryl, heterocyclic, hydroxyl, hydroxylamino, nitro, phenyl, phenoxy, and sulfonic acid, wherein said C_{1-7} alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-5} acyl, C_{1-5} acyloxy, C_{1-4} alkoxy, C_{1-8} alkyl, C_{1-4} alkylamino, C_{1-4} alkylcarboxamide, C_{1-4} alkylthiocarboxamide, C_{1-4} alkylsulfonamide, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-4} alkylthio, C_{1-4} alkylthiourey, C_{1-4} alkylureyl, amino, carbo- C_{1-6} -alkoxy, carboxamide, carboxy, cyano, C_{3-7} cycloalkyl, C_{2-8}

dialkylamino, C₂₋₆ dialkylcarboxamide, C₂₋₆ dialkylthiocarboxamide, C₂₋₆ dialkylsulfonamide, C₁₋₄ alkylthiourey, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkyl, C₁₋₄ haloalkylthio, halogen, heterocyclic, hydroxyl, hydroxylamino, and nitro.

22. The compound according to claim 1 wherein R₂ is -C(O)OR₂₂ and R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic.
23. The compound according to claim 1 wherein R₂ is -C(O)OR₂₂ and R₂₂ is C₁₋₈ alkyl, or C₃₋₇ cycloalkyl each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid.
24. The compound according to claim 1 wherein R₂ is -C(O)OR₂₂ and R₂₂ is C₁₋₈ alkyl, or C₃₋₇ cycloalkyl wherein said C₃₋₇ cycloalkyl is optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy, C₁₋₇ alkyl, carboxy, C₂₋₈ dialkylamino, and halogen.
25. The compound according to claim 1 wherein R₂ is -C(O)OR₂₂ and R₂₂ is C₁₋₈ alkyl, or C₃₋₇ cycloalkyl
26. The compound according to claim 1 wherein R₂ is -C(O)R₂₂ and R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic.

27. The compound according to claim 1 wherein R_2 is $-C(O)R_{22}$ and R_{22} is C_{1-8} alkyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of H, C_{1-4} alkoxy, C_{1-7} alkyl, amino, carboxy, halogen, heteroaryl, hydroxyl, phenoxy, and sulfonic acid, wherein said C_{1-7} alkyl and phenoxy are optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C_{1-4} haloalkoxy, and heterocyclic.
28. The compound according to claim 1 wherein R_2 is $-CH_2R_{22}$, or $-R_{22}$ and R_{22} is C_{1-8} alkyl, C_{3-7} cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-5} acyl, C_{2-6} alkenyl, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylsulfonyl, amino, carbo- C_{1-6} -alkoxy, carboxy, cyano, C_{3-7} cycloalkyl, C_{2-8} dialkylamino, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C_{1-7} alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C_{1-4} haloalkoxy, and heterocyclic.
29. The compound according to claim 1 wherein R_2 is $-CH_2R_{22}$, or $-R_{22}$, and R_{22} is C_{1-8} alkyl, C_{3-7} cycloalkyl, or heteroaryl each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{1-5} acyl, C_{2-6} alkenyl, C_{1-4} alkoxy, carbo- C_{1-6} -alkoxy, carboxy, cyano, C_{3-7} cycloalkyl, and hydroxyl.
30. The compound according to claim 1 wherein R_2 is $-S(O)_2R_{22}$ and R_{22} is C_{1-8} alkyl, C_{3-7} cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{2-6} alkenyl, C_{1-4} alkoxy, C_{1-7} alkyl, C_{1-4} alkylsulfonyl, amino, carbo- C_{1-6} -alkoxy, carboxy, cyano, C_{3-7} cycloalkyl, C_{2-8} dialkylamino, C_{1-4} haloalkoxy, C_{1-4} haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C_{1-7} alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C_{1-4} haloalkoxy, and heterocyclic.
31. The compound according to claim 1 wherein R_2 is $-S(O)_2R_{22}$ and R_{22} is C_{1-8} alkyl, or heteroaryl and said heteroaryl is optionally substituted with 1 to 5 C_{1-7} alkyl.
32. The compound according to claim 1 wherein R_2 is $-CR_{23}R_{24}C(O)R_{22}$ and wherein R_{23} and R_{24} are each independently H or C_{1-8} alkyl; and
 R_{22} is C_{1-8} alkyl, C_{3-7} cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C_{2-6} alkenyl, C_{1-4}

alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic.

33. The compound according to claim 1 wherein R₂ is -CR₂₃R₂₄C(O)R₂₂ and wherein R₂₃ and R₂₄ are each independently H or C₁₋₈ alkyl; and
R₂₂ is phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, cyano, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, and phenyl.
34. The compound according to claim 1 wherein R₂ is -CR₂₃R₂₄C(O)NR₂₅R₂₂ and wherein R₂₃, R₂₄ and R₂₅ are each independently H or C₁₋₈ alkyl; and
R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic.
35. The compound according to claim 1 wherein R₂ is -CH₂C(O)NHR₂₂ and wherein R₂₂ is phenyl optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ haloalkyl, and halogen.
36. The compound according to claim 1 wherein Ar₁ is aryl or heteroaryl each optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅;
wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonamide, alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy,

C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonoxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

37. The compound according to claim 36 wherein Ar₁ is aryl.
38. The compound according to claim 36 wherein Ar₁ is heteroaryl.
39. The compound according to claim 1 wherein Ar₁ is phenyl optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅;

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonoxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

40. The compound according to claim 1 wherein Ar₁ is phenyl optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅;

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide,

C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonoxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

41. The compound according to claim 1 wherein Ar₁ is phenyl optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅;

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonoxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₈ alkyl, and halogen.

42. The compound according to claim 1 wherein Ar₁ is pyridyl optionally substituted with R₁₁, R₁₂, R₁₃, and R₁₄;

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, and sulfonamide, and wherein C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonamide, alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonoxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, and R₁₄ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

43. The compound according to claim 1 wherein Ar₁ is pyridyl optionally substituted with R₁₁, R₁₂, R₁₃, and R₁₄;

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, C₂₋₆ dialkylamino, halogen, heterocyclic, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, alkylsulfonyl, C₁₋₄ alkylthio, C₂₋₆ dialkylamino, and heterocyclic are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₄ alkylsulfonyl, C₃₋₇ cycloalkyloxy, heteroaryl, hydroxyl, phenyl, and phosphonoxy; and

R₁₂, R₁₃, and R₁₄ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

44. The compound according to claim 1 wherein Ar₁ is pyridyl optionally substituted with R₁₁, R₁₂, R₁₃, and R₁₄;

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, C₂₋₆ dialkylamino, halogen, heterocyclic, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋

₄ alkylamino, alkylsulfonyl, C₁₋₄ alkylthio, C₂₋₆ dialkylamino, and heterocyclic are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₄ alkylsulfonyl, C₃₋₇ cycloalkyloxy, heteroaryl, hydroxyl, phenyl, and phosphonoxy; and

R₁₂, R₁₃, and R₁₄ are each independently selected from the group consisting of C₁₋₈ alkyl, and halogen.

45. The compound according to claim 1 wherein Z is selected from the group consisting of C₁₋₅ acyl, C₁₋₈ alkyl, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, amino, carbamimidoyl, cyano, C₃₋₇ cycloalkyl, heterocyclic, and hydroxycarbamimidoyl, wherein C₁₋₈ alkyl, C₃₋₇ cycloalkyl, and heterocyclic are each optionally substituted with 1, 2, 3 or 4 groups selected from the group consisting of C₁₋₅ acyl, C₁₋₅ acyloxy, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfinyl, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₁₋₄ alkylureyl, amino, C₁₋₂ alkylamino, C₂₋₄ dialkylamino, carbo-C₁₋₆-alkoxy, carboxamide, carboxy, cyano, formyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkylsulfinyl, C₁₋₄ haloalkylsulfonyl, C₁₋₄ haloalkylthio, halogen, hydroxyl, hydroxylamino and nitro, and wherein said C₁₋₇ alkyl is optionally substituted with amino.
46. The compound according to claim 1 wherein Z is selected from the group consisting of C₁₋₅ acyl, C₁₋₈ alkyl, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, amino, carbamimidoyl, cyano, C₃₋₇ cycloalkyl, heterocyclic, and hydroxycarbamimidoyl, wherein said heterocyclic is optionally substituted with a -CH₂NH₂ group.
47. The compound according to claim 1 wherein Z is selected from the group consisting of C(O)CH₃, C(O)CH₂CH₃, CH₃, CH₂CH₃, C≡CH, NHS(O)₂CH₃, amino, carbamimidoyl, cyano, cyclopropyl, 4,5-dihydro-1H-imidazol-2-yl, 5-aminomethyl-4,5-dihydro-oxazol-2-yl, and hydroxycarbamimidoyl.
48. The compound according to claim 1 wherein:
 - A and B are each independently -CH₂CH₂- or -CH₂-;
 - D is N-R₂;
 - V₁ is a bond;
 - V₂ is -CH₂-, -CH₂CH₂-, or a bond;
 - W and Q are each independently NH or O;
 - X and Y are each independently N or CH, provided that if either X or Y is CH then the other is N;

Z is selected from the group consisting of nitro, C₁₋₅ acyl, C₁₋₈ alkyl, C₂₋₆ alkynyl, C₁₋₄ alkylsulfonamide, amino, carbamimidoyl, cyano, C₃₋₇ cycloalkyl, heterocyclic, and hydroxycarbamimidoyl, wherein said heterocyclic is optionally substituted with a -CH₂NH₂ group;

R₂ is -C(O)OR₂₂, -C(O)R₂₂, -CH₂R₂₂, -R₂₂, -S(O)₂R₂₂, -CR₂₃R₂₄C(O)R₂₂, or -CR₂₃R₂₄C(O)NR₂₅R₂₂, wherein R₂₂ is C₁₋₈ alkyl, C₂₋₆ alkynyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl, phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic; and R₂₃ and R₂₄ are each independently H or C₁₋₈ alkyl;

Ar₁ is aryl or heteroaryl each optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅; wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, and sulfonamide, and wherein C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylsulfonamide, alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₃₋₇ cycloalkyloxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonoxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxamide, cyano, C₂₋₆ dialkylamino, and halogen.

49. The compound according to claim 1 wherein:

A and B are both -CH₂CH₂-;

D is N-R₂;

V₁ and V₂ are both a bond;

W and Q are each independently NH or O;

X and Y are both N;

Z is selected from the group consisting of nitro, C(O)CH₃, C(O)CH₂CH₃, CH₃, CH₂CH₃, C≡CH, NHS(O)₂CH₃, amino, carbamimidoyl, cyano, cyclopropyl, 4,5-dihydro-1H-imidazol-2-yl, 5-aminomethyl-4,5-dihydro-oxazol-2-yl, and hydroxycarbamimidoyl;

R₂ is -C(O)OR₂₂, wherein R₂₂ is C₁₋₈ alkyl, or C₃₋₇ cycloalkyl each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylsulfonyl, amino, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, and hydroxyl;

Ar₁ is phenyl optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅;

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonooxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₈ alkyl, and halogen.

50. The compound according to claim 1 wherein:

A and B are both -CH₂CH₂-;

D is N-R₂;

V₁ and V₂ are both a bond;

W is NH;

Q is O;

X and Y are both N;

Z is nitro, cyano, C(O)CH₃, amino, CH₃, CH₂CH₃, or C≡CH;

R₂ is -C(O)OR₂₂, -C(O)R₂₂, -R₂₂, or -S(O)₂R₂₂ wherein R₂₂ is C₁₋₈ alkyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₅ acyl, C₂₋₆ alkenyl, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carbo-C₁₋₆-alkoxy, carboxy, cyano, C₃₋₇ cycloalkyl, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, heterocyclic, hydroxyl,

phenyl, phenoxy, and sulfonic acid, wherein said C₁₋₇ alkyl, phenyl and phenoxy are each optionally substituted with 1 to 5 substituents selected from the group consisting of amino, C₁₋₄ haloalkoxy, and heterocyclic;

Ar₁ is phenyl, 3-pyridyl, or 2-pyridyl each optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅,

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, carboxamide, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, heteroarylcarbonyl, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, carbamimidoyl, C₂₋₆ dialkylamino, heterocyclic, heterocyclic-carbonyl, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₆ acylsulfonamide, C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, C₁₋₄ alkylsulfonyl, carboxy, C₂₋₆ dialkylamino, C₂₋₆ dialkylcarboxamide, heteroaryl, heterocyclic, hydroxyl, phenyl, and phosphonooxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently CH₃, or F.

52. The compound according to claim 1 wherein:

A and B are both -CH₂CH₂-;

D is N-R₂;

V₁ and V₂ are both a bond;

W and Q are both O;

X and Y are both N;

Z is selected from the group consisting of CH₃, CH₂CH₃, cyclopropyl, or C≡CH;

R₂ is -C(O)OR₂₂, -C(O)R₂₂, -R₂₂, -CH₂C(O)R₂₂, or -CH₂C(O)NHR₂₂, wherein R₂₂ is C₁₋₈ alkyl, C₂₋₆ alkynyl, C₃₋₇ cycloalkyl, phenyl, heteroaryl, or heterocyclic each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylsulfonyl, amino, carboxy, cyano, C₂₋₈ dialkylamino, C₁₋₄ haloalkoxy, C₁₋₄ haloalkyl, halogen, heteroaryl, hydroxyl, phenyl, and phenoxy, wherein said C₁₋₇ alkyl, is optionally substituted with 1 or 2 substituents selected from the group consisting of C₁₋₄ haloalkoxy, and heterocyclic;

Ar₁ is phenyl, 2-pyridyl, or 3-pyridyl each optionally substituted with R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅,

wherein R₁₁ is selected from the group consisting of C₁₋₆ acylsulfonamide,

C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, amino, carbamimidoyl, carboxy, cyano, C₂₋₆ dialkylamino, halogen, heterocyclic, heterocyclic-oxy, heterocyclic-carbonyl, heteroaryl, and sulfonamide, and wherein C₁₋₄ alkoxy, C₁₋₈ alkyl, C₁₋₄ alkylamino, C₁₋₆ alkylcarboxamide, C₁₋₄ alkylsulfonyl, C₁₋₄ alkylthio, C₂₋₆ dialkylamino, and heteroaryl are each optionally substituted with 1 to 5 substituents selected independently from the group consisting of C₁₋₄ alkoxy, C₁₋₇ alkyl, C₁₋₄ alkylcarboxamide, heteroaryl, hydroxyl, and phosphonooxy wherein said C₁₋₇ alkyl and C₁₋₄ alkylcarboxamide are each optionally substituted with 1 to 5 substituents selected from the group consisting of C₁₋₄ alkoxy and hydroxy; and

R₁₂, R₁₃, R₁₄, and R₁₅ are each independently selected from the group consisting of C₁₋₈ alkyl, and halogen.

53. The compound according to any one of claims 48 to 52 wherein R₂ is selected from the group consisting of methoxycarbonyl, ethoxycarbonyl, *iso*-propoxycarbonyl, *n*-propoxycarbonyl, *n*-butoxycarbonyl, *tert*-butoxycarbonyl, *iso*-butoxycarbonyl, and *n*-pentyloxycarbonyl.
54. The compound according to any one of claims 48 to 52 wherein R₁₁ is selected from the group consisting of sulfamoyl, acetylsulfamoyl, propionylsulfamoyl, butyrylsulfamoyl, pentanoylsulfamoyl, methanesulfonyl, ethanesulfonyl, propane-1-sulfonyl, hydroxymethyl; 2-hydroxyethyl; 3-hydroxypropyl; 4-hydroxy-butyl; phosphonooxymethyl; 2-phosphonooxyethyl; 3-phosphonooxy-propyl; and 4-phosphonooxy-butyl.
55. The compound according to claim 1 wherein said compound is selected from the group consisting of:
 - 3-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxymethyl]-pyrrolidine-1-carboxylic acid *tert*-butyl ester;
 - 4-[5-Cyano-6-(6-methylsulfanyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid *tert*-butyl ester;
 - 4-[5-Cyano-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid *tert*-butyl ester;
 - [6-(1-Hexyl-piperidin-4-yloxy)-5-nitro-pyrimidin-4-yl]-(4-methanesulfonyl-phenyl)-amine;
 - [6-(1-Cyclopropylmethyl-piperidin-4-yloxy)-5-nitro-pyrimidin-4-yl]-(4-methanesulfonyl-phenyl)-amine;
 - 4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid 2-isopropyl-5-methyl-cyclohexyl ester;

{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-pyridin-3-yl-methanone;

(2-Chloro-pyridin-3-yl)-{4-[6-(4-methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-methanone;

{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-pyridin-2-yl-methanone;

(4-Methanesulfonyl-phenyl)-[6-(1-methanesulfonyl-piperidin-4-yloxy)-5-nitro-pyrimidin-4-yl]-amine;

(4-Methanesulfonyl-phenyl)-{5-nitro-6-[1-(propane-1-sulfonyl)-piperidin-4-yloxy]-pyrimidin-4-yl}-amine;

{6-[1-(Butane-1-sulfonyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine;

(4-Methanesulfonyl-phenyl)-{5-nitro-6-[1-(thiophene-2-sulfonyl)-piperidin-4-yloxy]-pyrimidin-4-yl}-amine;

(4-Methanesulfonyl-phenyl)-{6-[1-(1-methyl-1H-imidazole-4-sulfonyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-amine;

{6-[1-(2,4-Dimethyl-thiazole-5-sulfonyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine;

4-[5-Cyano-6-(3-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

4-[6-(6-Methanesulfonyl-pyridin-3-ylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

4-[5-Acetyl-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

4-[5-Amino-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid ethyl ester;

4-[5-Cyano-6-(4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isobutyl ester;

4-(4-Methanesulfonyl-phenylamino)-6-[1-(tetrahydro-furan-2-carbonyl)-piperidin-4-yloxy]-pyrimidine-5-carbonitrile;

4-[1-(3,3-Dimethyl-2-oxo-butyl)-piperidin-4-yloxy]-6-(4-methanesulfonyl-phenylamino)-pyrimidine-5-carbonitrile;

4-(4-Methanesulfonyl-phenylamino)-6-[1-(pyridine-3-carbonyl)-piperidin-4-yloxy]-pyrimidine-5-carbonitrile;

4-(1-Formyl-piperidin-4-yloxy)-6-(4-methanesulfonyl-phenylamino)-pyrimidine-5-carbonitrile and

4-(4-Methanesulfonyl-phenylamino)-6-[1-(pyridine-2-carbonyl)-piperidin-4-yloxy]-pyrimidine-5-carbonitrile;

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

56. The compound according to claim 1 wherein said compound is selected from the group consisting of:

4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

(4-Methanesulfonyl-phenyl)-[5-nitro-6-(piperidin-4-yloxy)-pyrimidin-4-yl]-amine;

1-{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-3,3-dimethyl-butan-1-one;

(4-Methanesulfonyl-phenyl)-[5-nitro-6-(1-pyridin-2-ylmethyl-piperidin-4-yloxy)-pyrimidin-4-yl]-amine;

(4-Methanesulfonyl-phenyl)-[5-nitro-6-(1-pyridin-3-ylmethyl-piperidin-4-yloxy)-pyrimidin-4-yl]-amine;

{6-[1-(3,3-Dimethyl-butyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine;

(4-Methanesulfonyl-phenyl)-{6-[1-(3-methyl-butyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-amine;

(4-Methanesulfonyl-phenyl)-[5-nitro-6-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yloxy)-pyrimidin-4-yl]-amine;

4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid ethyl ester;

1-{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-piperidin-1-yl}-3,3-dimethyl-butan-2-one;

{6-[1-(2-Ethoxy-ethyl)-piperidin-4-yloxy]-5-nitro-pyrimidin-4-yl}-(4-methanesulfonyl-phenyl)-amine;

4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxymethyl]-piperidine-1-carboxylic acid tert-butyl ester;

4-{2-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-ethyl}-piperidine-1-carboxylic acid tert-butyl ester;

3-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester and

3-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-yloxymethyl]-pyrrolidine-1-carboxylic acid tert-butyl ester;

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

57. The compound according to claim 1 wherein said compound is selected from the group consisting of:

4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-ylamino]-piperidine-1-carboxylic acid tert-butyl ester;

N-(4-Methanesulfonyl-phenyl)-5-nitro-N'-piperidin-4-yl-pyrimidine-4,6-diamine;

1-{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-ylamino]-piperidin-1-yl}-ethanone and

1-{4-[6-(4-Methanesulfonyl-phenylamino)-5-nitro-pyrimidin-4-ylamino]-piperidin-1-yl}-2,2-dimethyl-propan-1-one;

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

58. The compound according to claim 1 wherein said compound is selected from the group consisting of:

4-[6-(4-Cyano-2-fluoro-phenylamino)-5-ethynyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Ethynyl-6-(2-fluoro-4-[1,2,4]triazol-1-yl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Ethynyl-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-pyrimidin-4-ylamino}-3-fluoro-benzonitrile;

{5-Ethynyl-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-pyrimidin-4-yl}-(2-fluoro-4-methanesulfonyl-phenyl)-amine;

4-{6-[2,5-Difluoro-4-(2-methanesulfonyl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-sulfamoyl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[6-(2-Fluoro-ethyl)-2-methyl-pyridin-3-ylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{2-[4-Fluoro-6-(2-isopropoxy-ethyl)-pyridin-3-ylamino]-3-methyl-pyridin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2,5-Difluoro-4-(2-[1,2,4]triazol-1-yl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Ethynyl-6-[2-fluoro-4-(4-methoxy-pyridin-2-yl)-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-propionylsulfamoyl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester; and

4-{6-[2,3-Difluoro-4-(2-methanesulfonyl-ethyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

59. The compound according to claim 1 wherein said compound is selected from the group consisting of:

4-[5-Acetyl-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isobutyl ester;

1-[4-(1-Benzyl-azetidin-3-yloxy)-6-(6-methanesulfonyl-pyridin-3-ylamino)-pyrimidin-5-yl]-ethanone;

4-[5-Cyano-6-(6-propylamino-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Cyano-6-(2-fluoro-4-isopropylamino-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Cyano-6-(2-fluoro-4-propylamino-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Cyano-6-(2-fluoro-4-propoxy-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Cyano-6-(6-propyl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Cyano-6-[4-(2-dimethylamino-ethylsulfanyl)-2-fluoro-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Cyano-6-[4-(2-dimethylamino-ethanesulfonyl)-2-fluoro-phenylamino]-3-oxo-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Cyano-6-[2-fluoro-4-(4-methyl-piperazin-1-yl)-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Cyano-6-[2-fluoro-4-(3-methyl-butylamino)-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Cyano-6-(2-fluoro-4-morpholin-4-yl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Cyano-6-[4-(2-dimethylamino-ethylamino)-2-fluoro-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Cyano-6-(4-dimethylamino-2-fluoro-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Cyano-6-[2-fluoro-4-(2-pyrrolidin-1-yl-ethylamino)-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Cyano-6-[2-fluoro-4-(2-morpholin-4-yl-ethylamino)-phenylamino]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-iodo-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Cyano-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-morpholin-4-yl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2,5-Difluoro-4-propoxy-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-propylamino-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-methoxy-ethylamino)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-(6-{2-Fluoro-4-[(tetrahydro-furan-2-yl)methyl]-amino}-phenylamino)-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethylamino)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-(6-{2-Fluoro-4-[(2-methanesulfonyl-ethyl)-methyl-amino]-phenylamino}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Bromo-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Cyano-2-fluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

- 4-[6-(4-Cyano-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-[6-(2,5-Difluoro-4-morpholin-4-yl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-[6-(6-Chloro-2-methyl-pyridin-3-ylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-[5-Methyl-6-(2-methyl-6-morpholin-4-yl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-[5-(4,5-Dihydro-1H-imidazol-2-yl)-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidin-4-yl}-amine;
- 4-[6-(2-Fluoro-4-propoxy-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2-Fluoro-4-(2-methoxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2-Fluoro-4-(2-isopropoxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-[6-(6-Chloro-4-methyl-pyridin-3-ylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-(N-hydroxycarbamimidoyl)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-[5-Carbamimidoyl-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2-Fluoro-4-(tetrahydro-furan-2-ylmethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-[5-Methyl-6-(4-methyl-6-morpholin-4-yl-pyridin-3-ylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[6-(2-Methoxy-ethoxy)-2-methyl-pyridin-3-ylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[6-(2-Methoxy-ethoxy)-4-methyl-pyridin-3-ylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2,5-Difluoro-4-(2-methoxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-isopropoxy-ethylsulfamoyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2,5-Difluoro-4-(N-hydroxycarbamimidoyl)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Carbamoyl-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[(2-Fluoro-4-methanesulfonyl-phenyl)-(2-methoxy-ethyl)-amino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Carbamimidoyl-2,5-difluoro-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[4-(2-Ethoxy-ethoxy)-2-fluoro-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(tetrahydro-pyran-4-yloxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-hydroxy-ethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butan-1-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-pentan-1-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-butan-1-one;

4-{6-[2-Fluoro-4-(pyridin-2-ylmethoxy)-phenylamino]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[2-(2-Fluoro-4-methanesulfonyl-phenylamino)-3-methyl-pyridin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(6-Chloro-4-fluoro-pyridin-3-ylamino)-5-cyano-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester; and

4-[5-Amino-6-(2-fluoro-4-methanesulfonyl-phenylamino)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

60. The compound according to claim 1, of the formula:

4-({6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-yl}-isopropyl-amino)-methyl)-piperidine-1-carboxylic acid tert-butyl ester;

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

61. The compound according to claim 1 wherein said compound is selected from the group consisting of:

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-6-[1-(3-methoxy-propyl)-piperidin-4-yloxy]-5-methyl-pyrimidine;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methoxy-propan-2-ol;

4-{6-[2-Fluoro-4-(5-isopropoxymethyl-[1,2,4]oxadiazol-3-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(5-methoxy-pyridin-2-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[6-(2-Cyclopropoxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(pyridine-2-carbonyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methanesulfonylamino-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Methoxy-6'-methyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-2-(4-trifluoromethoxy-phenoxy)-propan-1-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-2-(4-trifluoromethoxy-phenoxy)-ethanone;

N-(4-Chloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetamide;

N-(3-Chloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetamide;

N-(3,5-Dichloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetamide;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-N-(4-trifluoromethyl-phenyl)-acetamide;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-N-phenyl-acetamide;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-N-(4-isopropyl-phenyl)-acetamide;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-N-(4-methoxy-phenyl)-acetamide;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-N-(3-trifluoromethyl-phenyl)-acetamide;

4-{6-[2-Fluoro-4-(3-methoxy-propane-1-sulfonyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[6-(2-Isopropoxy-ethyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Methyl-6-[2-methyl-6-(2-pyridin-2-yl-ethoxy)-pyridin-3-yloxy]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(thiophene-2-carbonyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-(6-{6-[(2-Isopropoxy-ethyl)-methyl-amino]-2-methyl-pyridin-3-yloxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[6-(2-Isopropoxy-ethanesulfonyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[6-(2-Hydroxy-ethanesulfonyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(6-Amino-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(3-methyl-butyl)-piperidin-4-yloxy]-pyrimidine;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-morpholin-4-yl-ethanone;

1-(3,4-Dichloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

1-(3-Chloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-thiophen-3-yl-ethanone;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-phenyl-ethanone;

1-(2,4-Dimethoxy-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(4-methyl-pentyl)-piperidin-4-yloxy]-pyrimidine;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-isopropoxy-propan-1-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-isopropoxy-butan-1-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-hydroxy-propan-1-one;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(5-pyridin-2-yl-thiophen-2-yl)-ethanone;

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(5-methyl-hexyl)-piperidin-4-yloxy]-pyrimidine;

3-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-oxo-propane-1-sulfonic acid

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-thiophen-2-yl-ethanone;

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-(1-pentyl-piperidin-4-yloxy)-pyrimidine;

4-(1-Butyl-piperidin-4-yloxy)-6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidine;

4-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-cyclohexanecarboxylic acid;

1-(4-Diethylamino-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(2-methyl-4-phenyl-furan-3-yl)-ethanone;

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-6-(1-hexyl-piperidin-4-yloxy)-5-methyl-pyrimidine;

4-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butyric acid;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-pentan-2-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-hexan-2-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-hexan-2-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-methyl-pentan-2-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-5-methyl-hexan-2-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-6-methyl-heptan-2-one;

5-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-oxo-pentanoic acid;

5-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-oxo-pentanenitrile;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-2-pyridin-2-yl-ethanone;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-pyridin-4-yl-ethanone;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-ylmethyl}-acrylic acid;

1-[1,4]Dioxan-2-yl-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

1-(2,3-Dihydro-[1,4]dioxin-2-yl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-p-tolyl-ethanone;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-methoxy-phenyl)-ethanone;

1-(2-Chloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

3-(2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetyl)-benzonitrile;

1-(2,4-Dimethyl-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

1-(4-Chloro-3-methyl-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

1-(4-Difluoromethoxy-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

1-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(5-phenyl-thiophen-2-yl)-ethanone;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-thiophen-2-yl-ethanone;

{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetic acid ethyl ester;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methoxy-propan-2-ol;

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-6-[1-(4-methoxy-cyclohexyl)-piperidin-4-yloxy]-5-methyl-pyrimidine;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-hexan-1-one;

4-{6-[2-Fluoro-4-(2-isobutoxy-ethoxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[4-(2-Cyclopropoxy-ethoxy)-2-fluoro-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[4-(2-Ethoxy-ethoxy)-2-fluoro-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(3-methoxy-propoxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-pyridin-2-yl-ethoxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(tetrahydro-pyran-4-yloxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[4-(2-tert-Butoxy-ethoxy)-2-fluoro-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-sulfo-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2,5-Difluoro-4-trifluoromethoxy-phenoxy)-5-ethynyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2,5-Difluoro-4-trifluoromethoxy-phenoxy)-5-prop-1-ynyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Ethynyl-6-(2-fluoro-4-methoxy-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Ethynyl-6-(6-methoxy-4-methyl-pyridin-3-yloxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Ethynyl-6-[6-(2-isopropoxy-ethyl)-2-methyl-pyridin-3-yloxy]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Cyano-2-fluoro-phenoxy)-5-ethynyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Ethynyl-6-(2-fluoro-4-[1,2,4]triazol-4-yl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Ethynyl-6-(2-fluoro-4-[1,2,4]triazol-1-yl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

1-{4-[5-Ethynyl-6-(2-fluoro-4-[1,2,4]triazol-1-yl-phenoxy)-pyrimidin-4-yloxy]-piperidin-1-yl}-3-pyridin-2-yl-propan-1-one;

4-{5-Ethynyl-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-pyrimidin-4-yloxy}-3-fluoro-benzonitrile;

5-Ethynyl-4-(2-fluoro-4-methanesulfonyl-phenoxy)-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-pyrimidine;

4-[1-(3-Ethyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-ethynyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidine;

4-[1-(3-Ethyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidine;

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(3-methyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-pyrimidine;

4-[6-(2-Fluoro-4-methanesulfonylamino-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

cis-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-cyclohexyl}-carbamic acid isopropyl ester;

trans-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-cyclohexyl}-carbamic acid isopropyl ester;

N-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-cyclohexyl}-3-methyl-butyramide;

N-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-cyclohexyl}-isobutyramide;

4-{6-[2,5-Difluoro-4-(2-methanesulfonyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[4-Fluoro-6-(2-methanesulfonyl-ethyl)-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Cyclopropyl-6-[2,5-difluoro-4-(2-hydroxy-ethyl)-phenoxy]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-(5-Cyclopropyl-6-{2,5-difluoro-4-[2-(4-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2,5-Difluoro-4-(2-morpholin-4-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

- 4-(6-{2-Fluoro-4-[2-(4-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[6-(2-Fluoro-ethyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2-Fluoro-4-(1-hydroxy-cyclopropylmethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{2-[2,5-Difluoro-4-(2-methanesulfonyl-ethyl)-phenoxy]-3-methyl-pyridin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- (R)-4-(6-{2-Fluoro-4-[2-(3-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;
- (S)-4-(6-{2-Fluoro-4-[2-(3-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;
- (R)-4-(5-Ethynyl-6-{2-fluoro-4-[2-(2-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;
- (S)-4-(2-{2-Fluoro-4-[2-(2-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-3-methyl-pyridin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[4-Fluoro-6-(2-morpholin-4-yl-ethyl)-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{5-Ethynyl-6-[4-fluoro-6-(2-methanesulfonyl-ethyl)-pyridin-3-yloxy]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{2-[2,5-Difluoro-4-(2-isopropoxy-ethyl)-phenoxy]-3-methyl-pyridin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2-Fluoro-4-(2-propionylsulfamoyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2-Fluoro-4-(2-sulfamoyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2,5-Difluoro-4-(2-sulfamoyl-ethyl)-phenoxy]-5-ethynyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2,5-Difluoro-4-(2-[1,2,4]triazol-1-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[2,3-Difluoro-4-(2-methanesulfonyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-(2-{2-Fluoro-4-[2-(6-methoxy-pyridin-2-yl)-ethyl]-phenoxy}-3-methyl-pyridin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;
- 4-(6-{2-Fluoro-4-[2-(3-methoxy-pyridin-2-yl)-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(3-Fluoro-1-oxy-pyridin-4-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(5'-Methoxy-6-methyl-[2,2']bipyridinyl-5-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{5-Ethynyl-6-[2-fluoro-4-(4-methoxy-pyridin-2-yl)-phenoxy]-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(3-methoxy-pyridin-2-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-(6-{2,5-Difluoro-4-[2-(3-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester; and

4-(6-{2,5-Difluoro-4-[2-(3-methoxy-piperidin-1-yl)-ethyl]-phenoxy}-5-ethynyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

62. The compound according to claim 1 wherein said compound is selected from the group consisting of:

4-[6-(2-Fluoro-4-morpholin-4-yl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-[6-(2-pyrrolidin-1-yl-ethyl)-pyridin-3-yl]-methanone;

(6-Amino-pyridin-3-yl)-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-methanone;

4-[5-Ethyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[6-(2-Isopropoxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[6-(2-Hydroxy-ethylsulfanyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Methyl-6-(2-methyl-6-pentyl-pyridin-3-yloxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(3-fluoro-phenyl)-ethanone;

4-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-6-[1-(2-pyridin-3-yl-ethyl)-piperidin-4-yloxy]-pyrimidine;

- 2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-trifluoromethoxy-phenyl)-ethanone;
- 2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-pyridin-2-yl-ethanone;
- 4-{6-[6-(2-Methoxy-ethanesulfonyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-(2-Fluoro-4-methanesulfonyl-phenoxy)-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine;
- 4-(6-{2-Fluoro-4-[(2-hydroxy-ethylcarbamoyl)-methyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;
- 4-[6-(5-Iodo-pyridin-2-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-(6-{2-Fluoro-4-[N-(2-isopropoxy-ethyl)-carbamimidoyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;
- 4-[6-(4-Carboxy-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-(4-Bromo-2-fluoro-phenoxy)-6-[1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yloxy]-5-methyl-pyrimidine;
- 4-[6-(5-Methanesulfonyl-pyridin-2-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[6-(2-Hydroxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-[5-Cyclopropyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 4-{6-[6-(2-Methanesulfonyl-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;
- 4-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-oxo-butyric acid;
- 2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(3-trifluoromethyl-phenyl)-ethanone;
- 4-[6-[6-(2-Methoxy-ethylsulfanyl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;
- 1-(2,5-Dimethoxy-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;
- 2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-pyridin-2-yl-ethanone;

4-[6-(6-Chloro-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-fluoro-phenyl)-ethanone;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-trifluoromethyl-phenyl)-ethanone;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3,3-dimethyl-butan-2-one;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-pyridin-3-yl-ethanone;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butan-2-one;

4-(6-{2-Fluoro-4-[(2-isopropoxy-ethylcarbamoyl)-methyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-(4-methanesulfonyl-phenyl)-ethanone;

1-(4-Chloro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

4-(2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-acetyl)-benzonitrile;

1-(3,4-Difluoro-phenyl)-2-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

4-{6-[2-Fluoro-4-(2-isopropoxy-ethylcarbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-butan-1-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-pentan-1-one;

4-[6-(2,4-Difluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-butan-1-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-methyl-pentan-1-one;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-5-methyl-hexan-1-one;

4-{6-[2-Fluoro-4-(2-methoxy-ethylcarbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Bromo-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(methoxy-methyl-carbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methoxy-propan-1-one;

4-[6-(4-Cyano-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-(5-Aminomethyl-4,5-dihydro-oxazol-2-yl)-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[6-(2-Methoxy-ethylamino)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[6-(3-Methanesulfonyl-pyrrolidin-1-yl)-2-methyl-pyridin-3-yloxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(6-Benzylamino-2-methyl-pyridin-3-yloxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Carbamoyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-isopropoxy-ethylamino)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-(6-{2-Fluoro-4-[(tetrahydro-furan-2-ylmethyl)-amino]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

4-(6-{6-[(2-Methanesulfonyl-ethyl)-methyl-amino]-2-methyl-pyridin-3-yloxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-hydroxycarbamoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-pyrrolidin-1-yl-ethylcarbamoyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(4-isopropyl-piperazine-1-carbonyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-morpholin-4-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-methanesulfonyl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-hydroxy-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Carboxymethyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Dimethylcarbamoylmethyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-sulfamoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-propionylsulfamoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Ethynyl-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-phosphonoxy-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-[5-Bromo-6-(2-fluoro-4-methanesulfonyl-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-(6-{2-Fluoro-4-[2-(2-methanesulfonyl-pyrrolidin-1-yl)-2-oxo-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Carbamoylmethyl-2-fluoro-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-4-[(tetrahydro-furan-2-ylmethyl)-carbamoyl]-methyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(2-Fluoro-3-sulfamoyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

C-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-C-(4-fluoro-phenyl)-methylethylamine;

3-tert-Butoxy-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-propan-1-one;

2-Ethoxy-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-ethanone;

{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-(tetrahydro-furan-2-yl)-methanone;;

(S)-1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-2-methylamino-butan-1-one;

4-(6-{2-Fluoro-4-[2-(3-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-phenoxy}-5-methyl-pyrimidin-4-yloxy)-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-morpholin-4-yl-2-oxo-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-imidazol-1-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

4-{6-[2-Fluoro-4-(2-[1,2,3]triazol-1-yl-ethyl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

(R)-1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-2-methylamino-butan-1-one;

(S)-1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-hydroxy-butan-1-one;

(R)-N-(1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carbonyl}-2-methyl-propyl)-acetamide;

(S)-N-(1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carbonyl}-2-methyl-propyl)-acetamide;

(R)-N-(2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-methyl-2-oxo-ethyl)-acetamide;

(S)-N-(2-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-1-methyl-2-oxo-ethyl)-acetamide;

4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid (S)-tetrahydro-furan-3-yl ester;

4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid (R)-tetrahydro-furan-3-yl ester;

4-[6-(2-Amino-4-ethanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

4-[6-(4-Methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

(1-{4-[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidine-1-carbonyl}-2-methyl-propyl)-carbamic acid tert-butyl ester;

4-{6-[2-Fluoro-4-(6-methoxy-pyridin-3-yl)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester;

3-Amino-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-4-methyl-pentan-1-one;

2-Amino-1-{4-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yloxy]-piperidin-1-yl}-3-methyl-butan-1-one;

4-{6-[2-Fluoro-4-(2-isopropoxy-ethoxy)-phenoxy]-5-methyl-pyrimidin-4-yloxy}-piperidine-1-carboxylic acid isopropyl ester; and

4-[5-Methyl-6-(4-sulfo-phenoxy)-pyrimidin-4-yloxy]-piperidine-1-carboxylic acid isopropyl ester;

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

63. The compound according to claim 1 wherein said compound is selected from the group consisting of:

4-({Cyclopropyl-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-amino}-methyl)-piperidine-1-carboxylic acid tert-butyl ester;

4-({Cyclopropyl-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-amino}-methyl)-piperidine-1-carboxylic acid isopropyl ester;

4-({[6-(2-Fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-isopropyl-amino}-methyl)-piperidine-1-carboxylic acid isopropyl ester; and

4-({Cyclopropylmethyl-[6-(2-fluoro-4-methanesulfonyl-phenoxy)-5-methyl-pyrimidin-4-yl]-amino}-methyl)-piperidine-1-carboxylic acid isopropyl ester;

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

64. The compound according to claim 1 of the formula:

4-[6-(2-Fluoro-4-methanesulfonyl-phenylamino)-5-methyl-pyrimidin-4-ylsulfanyl]-piperidine-1-carboxylic acid isopropyl ester;

or a pharmaceutically acceptable salt, hydrate or solvate thereof.

65. A pharmaceutical composition comprising at least one compound according to any one of claims 1 to 64 and a pharmaceutically acceptable carrier.
66. A method for treatment of a metabolic-related disorder in an individual comprising administering to said individual in need of such treatment a therapeutically effective amount of a compound according to any one of claims 1 to 64 or a pharmaceutical composition of claim 65.
67. The method according to claim 66 wherein said metabolic-related disorder is selected from the group consisting of type I diabetes, type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia and syndrome X.
68. The method according to claim 67 wherein said metabolic-related disorder is type II diabetes.

69. The method according to claim 67 wherein said metabolic-related disorder is hyperglycemia.
70. The method according to claim 67 wherein said metabolic-related disorder is hyperlipidemia.
71. The method according to claim 67 wherein said metabolic-related disorder is hypertriglyceridemia.
72. The method according to claim 67 wherein said metabolic-related disorder is type I diabetes.
73. The method according to claim 67 wherein said metabolic-related disorder is dyslipidemia.
74. The method according to claim 67 wherein said metabolic-related disorder is syndrome X.
75. The method according to any one of claims 66 to 74 wherein said individual is a mammal.
76. The method according to claim 75 wherein said mammal is a human.
77. A method of decreasing food intake of an individual comprising administering to said individual in need thereof a therapeutically effective amount of a compound according to any one of claims 1 to 64 or a pharmaceutical composition of claim 65.
78. A method of inducing satiety in an individual comprising administering to said individual in need thereof a therapeutically effective amount of a compound according to any one of claims 1 to 64 or a pharmaceutical composition of claim 65.
79. A method of controlling or decreasing weight gain of an individual comprising administering to said individual in need thereof a therapeutically effective amount of a compound according to any one of claims 1 to 64 or a pharmaceutical composition of claim 65.
80. The method according to any one of claims 77 to 79 wherein said individual is a mammal.
81. The method according to claim 80 wherein said mammal is a human.
82. The method according to claim 81 wherein said human has a body mass index of about 18.5 to about 45.

83. The method according to claim 81 wherein said human has a body mass index of about 25 to about 45.
84. The method according to claim 81 wherein said human has a body mass index of about 30 to about 45.
85. The method according to claim 81 wherein said human has a body mass index of about 35 to about 45.
86. A method of modulating a **RUP3** receptor in an individual comprising contacting the receptor with a compound according to any one of claims 1 to 64.
87. The method of modulating the **RUP3** receptor according to claim 86 wherein said compound is an agonist.
88. The method of modulating the **RUP3** receptor according to claim 86 or 87 wherein said modulation of the **RUP3** receptor is treatment of a metabolic-related disorder.
89. The method of modulating the **RUP3** receptor according to claim 88 wherein said metabolic-related disorder is selected from the group consisting of type I diabetes, type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia or syndrome X.
90. The method of modulating the **RUP3** receptor according to claim 89 wherein said metabolic-related disorder is type II diabetes.
91. The method of modulating the **RUP3** receptor according to claim 89 wherein said metabolic-related disorder is insulin resistance.
92. The method of modulating the **RUP3** receptor according to claim 89 wherein said metabolic-related disorder is hyperglycemia.
93. The method of modulating the **RUP3** receptor according to claim 89 wherein said metabolic-related disorder is hyperlipidemia.
94. The method of modulating the **RUP3** receptor according to claim 89 wherein said metabolic-related disorder is hypertriglyceridemia.

95. The method of modulating the **RUP3** receptor according to claim 89 wherein said metabolic-related disorder is type I diabetes.
96. The method of modulating the **RUP3** receptor according to claim 89 wherein said metabolic-related disorder is dyslipidemia.
97. The method of modulating the **RUP3** receptor according to claim 89 wherein said metabolic-related disorder is syndrome X.
98. The method of modulating the **RUP3** receptor according to any one of claims 86 to 97 wherein said individual is a mammal.
99. The method of modulating the **RUP3** receptor according to claim 98 wherein said mammal is a human.
100. The method of modulating the **RUP3** receptor according to claim 86 or 87 wherein said modulation of the **RUP3** receptor reduces food intake of said individual.
101. The method of modulating the **RUP3** receptor according to claim 86 or 87 wherein said modulation of the **RUP3** receptor induces satiety in said individual.
102. The method of modulating the **RUP3** receptor according to claim 86 or 87 wherein said modulation of the **RUP3** receptor controls or reduces weight gain of said individual.
103. The method of modulating the **RUP3** receptor according to any one of claims 100 to 102 wherein said individual is a mammal.
104. The method of modulating the **RUP3** receptor according to claim 103 wherein said mammal is a human.
105. The method according to claim 104 wherein said human has a body mass index of about 18.5 to about 45.
106. The method according to claim 104 wherein said human has a body mass index of about 25 to about 45.

107. The method according to claim 104 wherein said human has a body mass index of about 30 to about 45.
108. The method according to claim 104 wherein said human has a body mass index of about 35 to about 45.
109. Use of a compound according to any one of claims 1 to 64 for production of a medicament for use in treatment of a metabolic-related disorder.
110. The use of a compound according to claim 109 wherein said metabolic-related disorder is selected from the group consisting of type I diabetes, type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia or syndrome X.
111. Use of a compound according to any one of claims 1 to 64 for production of a medicament for use in decreasing food intake in an individual.
112. Use of a compound according to any one of claims 1 to 64 for production of a medicament for use of inducing satiety in an individual.
113. Use of a compound according to any one of claims 1 to 64 for production of a medicament for use in controlling or decreasing weight gain in an individual.
114. The use according to any one of claims 111 to 113 wherein said individual is a mammal.
115. The use according to claim 114 wherein said mammal is a human.
116. The use according to claim 115 wherein said human has a body mass index of about 18.5 to about 45.
117. The use according to claim 115 wherein said human has a body mass index of about 25 to about 45.
118. The use according to claim 115 wherein said human has a body mass index of about 30 to about 45.

119. The use according to claim 115 wherein said human has a body mass index of about 35 to about 45.
120. A compound according to any one of claims 1 to 64 for use in a method of treatment of the human or animal body by therapy.
121. A compound according to any one of claims 1 to 64 for use in a method of prophylaxis or treatment of a metabolic-related disorder of the human or animal body by therapy.
122. A compound according to any one of claims 1 to 64 for use in a method of decreasing food intake of the human or animal body by therapy.
123. A compound according to any one of claims 1 to 64 for use in a method of inducing satiety of the human or animal body by therapy.
124. A compound according to any one of claims 1 to 64 for use in a method of controlling or decreasing weight gain of the human or animal body by therapy.
122. The method of producing a pharmaceutical composition comprising admixing at least one compound of any one of claims 1 to 64 and a pharmaceutically acceptable carrier.




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4

SEQUENCE LISTING

<110> Arena Pharmaceuticals, Inc.
 Jones, Robert M.
 Semple, Graeme
 Xiong, Yifeng
 Shin, Young-Jun
 Ren, Albert S.
 Calderon, Imelda
 Fioravanti, Beatriz
 Choi, Jin Sun Karoline
 Lehmann, Juerg
 Bruce, Marc A.

<120> TRISUBSTITUTED ARYL AND HETEROARYL DERIVATIVES AS
 MODULATORS OF METABOLISM AND THE PROPHYLAXIS AND TREATMENT OF
 DISORDERS RELATED THERETO

<130> 69.W01

<150> 60/487,370
 <151> 2003-07-14

<150> 60/486,728
 <151> 2003-07-11

<160> 7

<170> PatentIn version 3.2

<210> 1
 <211> 1191
 <212> DNA
 <213> Homo sapiens

<400> 1
 atgtacccat acgacgtccc agactacgct ggaagcttgg aatcatcttt ctcatattgga 60
 jtgatccttg ctgtcctggc ctccctcatc attgctacta acacactagt ggctgtggct 120
 jtgtgtgtgt tgatccacaa gaatgatggt gtcagtctct gcttcacctt gaatctggct 180
 jtggctgaca ccttgattgg tgtggccatc tctggcctac tcacagacca gctctccagc 240
 :cttctcggc ccacacagaa gaccctgtgc agcctgcgga tggcatttgt cacttcctcc 300
 |cagctgcct ctgtcctcac ggtcatgctg atcaccttg acaggtacct tgccatcaag 360
 :agcccttcc gctacttgaa gatcatgagt gggttcgtgg ccggggcctg cattgccggg 420
 tgtgggttag tgtcttacct cattggcttc ctccactcg gaatcccat gttccagcag 480
 ctgcctaca aagggcagtg cagcttcttt gctgtatttc accctcactt cgtgctgacc 540
 tctcctgcg ttggcttctt ccagccatg ctctctttg tcttcttcta ctgcgacatg 600
 tcaagattg cctccatgca cagccagcag attcgaaaga tggaacatgc aggagccatg 660
 ctggagggtt atcgatcccc acggactccc agcgacttca aagctctccg tactgtgtct 720
 ttctcattg ggagctttgc tctatcctgg acccccttcc ttatcactgg cattgtgcag 780
 tggcctgcc aggagtgtca cctctaccta gtgctggaac ggtacctgtg gctgctcggc 840

```

gtgggcaact ccctgctcaa cccactcatc tatgcctatt ggcagaagga ggtgcgactg    900
cagctctacc acatggccct aggagtgaag aaggtgctca cctcattcct cctctttctc    960
tcggccagga attgtggccc agagaggccc agggaaagtt cctgtcacat cgtcactatc   1020
tccagctcag agtttgatgg cgaattcgga tccaagggca attctgcaga tatccagcac   1080
agtggcggcc gctcgagtct agagggcccg cggttcgaag gtaagcctat ccctaaccct   1140
ctcctcggtc tcgattctac gcgtaccggt catcatcacc atcaccattg a             1191

```

```

<210> 2
<211> 396
<212> PRT
<213> Homo sapiens

```

```

<400> 2

```

```

Met Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Gly Ser Leu Glu Ser Ser
1      5      10      15

```

```

Phe Ser Phe Gly Val Ile Leu Ala Val Leu Ala Ser Leu Ile Ile Ala
20      25      30

```

```

Thr Asn Thr Leu Val Ala Val Ala Val Leu Leu Leu Ile His Lys Asn
35      40      45

```

```

Asp Gly Val Ser Leu Cys Phe Thr Leu Asn Leu Ala Val Ala Asp Thr
50      55      60

```

```

Leu Ile Gly Val Ala Ile Ser Gly Leu Leu Thr Asp Gln Leu Ser Ser
65      70      75      80

```

```

Pro Ser Arg Pro Thr Gln Lys Thr Leu Cys Ser Leu Arg Met Ala Phe
85      90      95

```

```

Val Thr Ser Ser Ala Ala Ala Ser Val Leu Thr Val Met Leu Ile Thr
100     105     110

```

```

Phe Asp Arg Tyr Leu Ala Ile Lys Gln Pro Phe Arg Tyr Leu Lys Ile
115     120     125

```

```

Met Ser Gly Phe Val Ala Gly Ala Cys Ile Ala Gly Leu Trp Leu Val
130     135     140

```

```

Ser Tyr Leu Ile Gly Phe Leu Pro Leu Gly Ile Pro Met Phe Gln Gln
145     150     155     160

```

```

Thr Ala Tyr Lys Gly Gln Cys Ser Phe Phe Ala Val Phe His Pro His
165     170     175

```

Phe Val Leu Thr Leu Ser Cys Val Gly Phe Phe Pro Ala Met Leu Leu
 180 185 190
 Phe Val Phe Phe Tyr Cys Asp Met Leu Lys Ile Ala Ser Met His Ser
 195 200 205
 Gln Gln Ile Arg Lys Met Glu His Ala Gly Ala Met Ala Gly Gly Tyr
 210 215 220
 Arg Ser Pro Arg Thr Pro Ser Asp Phe Lys Ala Leu Arg Thr Val Ser
 225 230 235 240
 Val Leu Ile Gly Ser Phe Ala Leu Ser Trp Thr Pro Phe Leu Ile Thr
 245 250 255
 Gly Ile Val Gln Val Ala Cys Gln Glu Cys His Leu Tyr Leu Val Leu
 260 265 270
 Glu Arg Tyr Leu Trp Leu Leu Gly Val Gly Asn Ser Leu Leu Asn Pro
 275 280 285
 Leu Ile Tyr Ala Tyr Trp Gln Lys Glu Val Arg Leu Gln Leu Tyr His
 290 295 300
 Met Ala Leu Gly Val Lys Lys Val Leu Thr Ser Phe Leu Leu Phe Leu
 305 310 315 320
 Ser Ala Arg Asn Cys Gly Pro Glu Arg Pro Arg Glu Ser Ser Cys His
 325 330 335
 Ile Val Thr Ile Ser Ser Ser Glu Phe Asp Gly Glu Phe Gly Ser Lys
 340 345 350
 Gly Asn Ser Ala Asp Ile Gln His Ser Gly Gly Arg Ser Ser Leu Glu
 355 360 365
 Gly Pro Arg Phe Glu Gly Lys Pro Ile Pro Asn Pro Leu Leu Gly Leu
 370 375 380
 Asp Ser Thr Arg Thr Gly His His His His His His
 385 390 395

:210> 3
 :211> 24
 :212> DNA
 :213> Artificial

:220>
 :223> Homo sapiens Primer

<400> 3
cattgccggg ctgtggtttag tgtc

24

<210> 4
<211> 24
<212> DNA
<213> Artificial

<220>
<223> Homo sapiens Primer

<400> 4
ggcatagatg agtggggtga gcag

24

<210> 5
<211> 22
<212> DNA
<213> Artificial

<220>
<223> Rat Primer

<400> 5
catgggccct gcaccttctt tg

22

<210> 6
<211> 24
<212> DNA
<213> Artificial

<220>
<223> Rat Primer

<400> 6
gctccggatg gctgatgata gtga

24

<210> 7
<211> 23
<212> PRT
<213> Artificial

<220>
<223> Chemically synthesized peptide

<400> 7

Arg Gly Pro Glu Arg Thr Arg Glu Ser Ala Tyr His Ile Val Thr Ile
1 5 10 15

Ier His Pro Glu Leu Asp Gly
20

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12 C07D401/14 C07D403/12 C07D405/14 C07D409/14
 C07D413/14 C07D417/14 C07D239/52 A61K31/506 A61K31/5377
 A61K31/505 A61P3/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 03/002544 A (BRISTOL MYERS SQUIBB) 9 January 2003 (2003-01-09)	1-6, 8, 11, 15, 17, 19-21, 36, 37, 39-41, 45-47, 65-67, 75, 76, 109, 110, 120, 121, 125
Y	page 49 paragraph 2 pages 36, 38, ; claims; example 60 ----- -/--	1-125

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

*** Special categories of cited documents :**

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

29 November 2004

Date of mailing of the international search report

10/12/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Helps, I

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 948 786 A (FUJIWARA ET. AL.) 7 September 1999 (1999-09-07) column 3, line 39 - line 67; claims; examples	1-125
A	WO 02/098864 A (F. HOFFMANN-LA ROCHE) 12 December 2002 (2002-12-12) claims; examples	1-125
A	US 6 414 002 B1 (CHENG ET. AL.) 2 July 2002 (2002-07-02) claims; examples	1-125
A	G.B. PHILLIPS ET. AL.: "Discovery of N-'2-'5-'Amino(imino)methyl-2-hydroxyphen oxy-3,5-difluoro-6-'3-(4,5-dihydro-1-meth yl-1H-imidazol-2-yl)-phenoxy!pyridin-4-yl! N-methylglycine (ZK-807834). A Potent, Selective, and Orally Active Inhibitor of the Blood Coagulation Enzyme Factor Xa." JOURNAL OF MEDICINAL CHEMISTRY, vol. 41, 1998, pages 3557-3562, XP002307980 page 3560; table 3	1-125
E	WO 2004/076413 A (ARENA PHARMACEUTICALS) 10 September 2004 (2004-09-10) page 1 - page 4; claims; examples	1-125
E	WO 2004/065380 A (ARENA PHARMACEUTICALS) 5 August 2004 (2004-08-05) page 1 - page 4; claims; examples	1-125

INTERNATIONAL SEARCH REPORT

International application no.
PCT/US2004/022327

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 66-108(part)
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 66-108 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03002544	A	09-01-2003	CA 2451128 A1	09-01-2003
			EP 1406875 A1	14-04-2004
			WO 03002544 A1	09-01-2003
			US 2003139435 A1	24-07-2003
US 5948786	A	07-09-1999	AT 230740 T	15-01-2003
			AU 2521597 A	07-11-1997
			CA 2250943 A1	23-10-1997
			DE 69718306 D1	13-02-2003
			EP 0892795 A1	27-01-1999
			WO 9738992 A1	23-10-1997
			JP 2001511764 T	14-08-2001
WO 02098864	A	12-12-2002	BR 0210102 A	08-06-2004
			CA 2448602 A1	12-12-2002
			WO 02098864 A1	12-12-2002
			EP 1397351 A1	17-03-2004
			US 2003060466 A1	27-03-2003
US 6414002	B1	02-07-2002	US 2003069275 A1	10-04-2003
			US 2003096846 A1	22-05-2003
			US 2003087935 A1	08-05-2003
			US 2004171644 A1	02-09-2004
			US 2004147560 A1	29-07-2004
			AU 7593500 A	24-04-2001
			BR 0014189 A	20-08-2002
			CA 2388452 A1	29-03-2001
			CN 1374955 T	16-10-2002
			EP 1218361 A1	03-07-2002
			HU 0204416 A2	28-04-2003
			JP 2003509503 T	11-03-2003
			NO 20021408 A	14-05-2002
			TR 200200732 T2	21-10-2002
			WO 0121602 A1	29-03-2001
			ZA 200200937 A	02-05-2003
WO 2004076413	A	10-09-2004	WO 2004076413 A2	10-09-2004
WO 2004065380	A	05-08-2004	WO 2004065380 A1	05-08-2004